



Contact Information  
w.bader@ulg.ac.be  
http://girpas.astro.ulg.ac.be


Université  
de Liège



# Retrievals of ethane from ground-based high-resolution FTIR solar observations with updated line parameters: determination of the optimum strategy for the Jungfraujoch station.

W. Bader (1), A. Perrin (2), D. Jacquemart (3), K. Sudo (4), H. Yashiro (5), M. Gauss (6), P. Demoulin (1), C. Servais (1), and E. Mahieu (1).

(1) University of Liège, Liège, Belgium (w.bader@ulg.ac.be), (2) Laboratoire Interuniversitaire des Systèmes Atmosphériques, Paris, France, (3) Laboratoire de Dynamique, Interactions et Réactivité, Université Pierre et Marie Curie, Paris, France, (4) Graduate School of Environmental Studies, Nagoya University, Nagoya, Aichi, Japan, (5) Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, Japan, (6) Norwegian Meteorological Institute, Research and Development Department, Oslo, Norway.



Poster presented at the EGU General Assembly 2012  
Vienna, Austria, 22-27 April 2012.  
Permalink : <http://hdl.handle.net/2268/117067>

## 1. Introduction

The Ethane (C<sub>2</sub>H<sub>6</sub>) is the most abundant Non-Methane HydroCarbon (NMHC) in the Earth's atmosphere, with a lifetime of approximately 2 months. C<sub>2</sub>H<sub>6</sub> main sources are biomass burning, natural gas loss and biofuel consumption. Oxidation by the hydroxyl radical is by far the major C<sub>2</sub>H<sub>6</sub> sink as the seasonally changing OH concentration controls the strong modulation of the ethane abundance throughout the year. Ethane reduces Cl atom concentrations in the lower stratosphere and is a major source of peroxyacetyl nitrate (PAN) and carbon monoxide (by reaction with OH). Involved in the formation of tropospheric ozone and in the destruction of atmospheric methane through changes in OH, C<sub>2</sub>H<sub>6</sub> is an indirect greenhouse gas with a net-global warming potential of 5.5 (100-yr horizon).

All the spectra analyzed here have been recorded at the International Scientific Station of the Jungfraujoch (46.5°N, 8°E, 3580 m asl). This high altitude station offers excellent conditions to perform solar observations, in particular in the infrared (IR), because of weak local pollution (no major industries within 20km) and very high dryness. Indeed, the amount of water vapor, a strong interference in the IR, is at least twenty times lower than at sea level.

Two Fourier Transform Infrared (FTIR) instruments (a commercial Bruker IFS-120HR and a homemade spectrometer) have been put into regular operation in 1984 and 1991, respectively. They allow to record wide-band high-resolution IR solar spectra either with a MCT or InSb detector. Since 1991, the two FTIR instruments are affiliated to the framework of the Network for the Detection of Atmospheric Composition Change (NDACC, visit <http://www.ndacc.org>).

## 2. Retrieval Strategy

Parameters have been settled down on the basis of tests on a full year minimizing residuals and maximizing DOFS.

### Microwindows Selection




Figure 1 - Synthetic spectra of our three fitted  $\mu$ -windows (see limits on Table I) for all gases (in black) as well as for the individual contributors (C<sub>2</sub>H<sub>6</sub> in blue, H<sub>2</sub>O in cyan, CH<sub>4</sub> in green, O<sub>3</sub> in dark pink, CH<sub>3</sub>Cl in pink and solar lines in orange) to the absorption in this spectral interval. For clarity, the contributions of each species have been vertically displaced. Simulated solar zenith angle: 60°.

### A priori Volume Mixing Ratio (VMR) profiles




Figure 2 - The *a priori* VMR profiles tested (left panel) and their associated relative standard deviation (right panel) are illustrated. The first adopted *a priori* C<sub>2</sub>H<sub>6</sub> profile (grey crosses) is a zonal mean (for the 41-51°N latitude band) of 771 occultations recorded by the ACE-FTS instrument between the 2nd of November in 2004 and the 8th of February in 2011 extending from 8.5 to 20 km. The profile extension down to 3.58 km is based on EMEP in situ measurements (bottom panel) while the upper extension to 100 km is based on the WACCM model climatology (Whole Atmosphere Community Climate, <http://waccm.acd.ucar.edu>). Pink crosses is the *a priori* profile issued from the CHASER 3-D Chemical Transport Model developed at the Center for Climate System Research (CCSR), University of Tokyo/National Institute for Environmental Studies (NIES), Chemical Transport Model v.2 and v.3 from the University of Oslo (Berntsen *et al.*, 1997) are plotted in dark and light green crosses, respectively.

## References

- Berntsen T., and I. S. A. Saksen (1997), A global 3-D chemical transport model for the troposphere, 1, Model description and CO and Ozone results., *J. Geophys. Res.*, 102, 21239-21280.
- Bray C., A. Perrin, D. Jacquemart *et al.* (2011), The v1, v4 and 3v6 bands of methyl chloride in the 3.4  $\mu$  region: Line positions and intensities, *J. Quant. Spectrosc. Radiat. Transfer*, 112, 2446-2462.
- Gardiner T., A. Forbes, M. de Mazière *et al.* (2008), Trend analysis of greenhouse gases over Europe measured by a network of ground-based remote FTIR instruments., *Atmos. Chem. Phys.*, 8, 6719-6727.
- Harrison J. J., N. D. C. Allen and P. F. Bernath (2010), Infrared cross sections for ethane (C<sub>2</sub>H<sub>6</sub>) in the 3  $\mu$  region., *J. Quant. Spectrosc. Radiat. Transfer*, 111, 357-363.
- Rinsland C. P., N. B. Jones, and B. J. Connor (1998), Northern and southern hemisphere ground-based infrared spectroscopic measurements of tropospheric carbon monoxide and ethane, *J. Geophys. Res.*, 103 (D21), 28197.
- Rothman L.S., D. Jacquemart, A. Barbe *et al.* (2009), The HITRAN 2008 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transfer*, 110, 533-572.
- Sudo K., M. Takahashi, J. Kurokawa *et al.* (2002), CHASER : A global chemical model of the troposphere. 1. Model Description, *J. Geophys. Res.*, 107, 4339.

## Spectroscopic Linelists

As the current state of ethane parameters in HITRAN (e.g. : Rothman *et al.*, 2009, <http://www.hitran.com>) was rather unsatisfactory in the 3  $\mu$  region, new ethane absorption cross sections recorded at the Molecular Spectroscopy Facility of the Rutherford Appleton Laboratory (Harrison *et al.*, 2010) were combined with HITRAN 2004 line parameters (including all 2006 updates) and therefore used in our retrievals, after conversion into pseudolines by G. C. Toon (personal communication, 2011). These cross sections were calibrated in intensity by using reference low-resolution spectra from the Pacific Northwest National Laboratory (PNNL, Washington, USA, <http://www.pnl.gov/>) IR database.

We quantified the impact of two updates of the spectroscopic parameters for both Hitran 2004 and 2008. We first reckoned changes on spectral residuals (see Table I) induced by the update of two O<sub>3</sub> lines (encompassed in the 1-<sup>P</sup>Q<sub>3</sub>  $\mu$ -window) corrected by P. Chelin (LPMA, Paris, France) in the framework of the UFTIR project. Also, the improvement brought by the update of the line positions and intensities of methyl chloride (CH<sub>3</sub>Cl) in the 3.4  $\mu$  region (Bray *et al.*, 2011) is quantified. Improvements brought by Hitran-08 over the 2004 edition are illustrated on Figure 3.

$\mu$ -windows	1- <sup>P</sup> Q <sub>3</sub>	u2	u3	Global
1- Hitran-08	0.431	0.206	0.471	0.424
2- Harrison + 1	0.171	0.158	0.173	0.179
3- Chelin + 2	0.169	0.158	0.172	0.179
4- Bray + 3	0.153	0.151	0.152	0.163

Table I - RMS Values (in %) for each  $\mu$ -window (see Table II) according to each linelist tested.




Figure 3 - Mean observed (blue curve)/simulated (red curve) spectra and residuals (green curve) for 187 spectral fits (4.96 mK), using Hitran-04 (left panel) and Hitran-08 (right panel). Harrison's, Chelin's and Bray's updates are used for both.

## Summary

All retrievals have been performed with the SFIT-2 algorithm (v 3.91) (Rinsland *et al.*, 1998) in order to retrieve three ethane vertical columns (see next frame) on a series of about 13 000 spectra recorded between 1994 and 2011. The fitting has been narrowed down to 3 micro-windows described on Table II.

The C<sub>2</sub>H<sub>6</sub> a priori VMR profile adopted in all our retrievals, as well as its uncertainties, are based on synthetic profiles produced by the CHASER model for the 2007-2009 time period. A priori profiles for the interfering gases are based on the WACCM model climatology.

In the present runs, HITRAN 2008 line parameters including Harrison's pseudo lines, the two ozone lines update provided by P. Chelin (Personal Communication, 2004) and the updated CH<sub>3</sub>Cl lines (Bray *et al.*, 2011) as well as the solar line compilation provided by F. Hase (KIT) have been assumed for target and interfering absorptions. Adopted temperature and geopotential height data sets are provided by the National Centers for Environmental Prediction (NCEP, Washington, USA).

$\mu$ -windows	Limits (cm <sup>-1</sup> )	Interfering species
1 - <sup>P</sup> Q <sub>3</sub>	2976.66 - 2977.059	C <sub>2</sub> H <sub>6</sub> , H <sub>2</sub> O, CH <sub>4</sub> , O <sub>3</sub> , CH <sub>3</sub> Cl
2	2983.2 - 2983.5	C <sub>2</sub> H <sub>6</sub> , H <sub>2</sub> O, CH <sub>4</sub> , O <sub>3</sub> , CH <sub>3</sub> Cl
3	2986.43 - 2986.85	C <sub>2</sub> H <sub>6</sub> , H <sub>2</sub> O, CH <sub>4</sub> , O <sub>3</sub> , CH <sub>3</sub> Cl

Table II - List of microwindows used for our C<sub>2</sub>H<sub>6</sub> inversions, for each of them, the third column provides interfering gases adjusted during the retrieval.

## 3. Information Content and Error Budget




Figure 4 - Information content calculated for typical C<sub>2</sub>H<sub>6</sub> retrievals at the Jungfraujoch station. The three first eigenvectors are reproduced in the left frame. Right frame gives the corresponding error budget, with identification of the main error components, together with the assumed variability.

Information content and error budget have been carefully evaluated. Figure 4 displays typical results computed for VMR. The first eigen vector and corresponding eigenvalues (see left frame, in red) show that information on both selected C<sub>2</sub>H<sub>6</sub> partial columns, namely 3.58-6.79 km (low-tropospheric) and 8.45-14.3 km (Upper Tropospheric-Lower Stratospheric, UTLS), is mainly coming from the retrieval (99 %).

The error budget affecting the retrieved VMRs below 18 km is given in the right panel of Figure 4. Total and individual error profiles are provided and identified (see color codes).

We are still working on establishing a complete error budget. Nevertheless, we have already quantified the impact on our retrieved C<sub>2</sub>H<sub>6</sub> of the NCEP temperature profiles. We computed respectively for our total, low-tropospheric and UTLS columns an error of 0.88, 3.61 and 3.69 %. A maximum error during Spring (10 %) and a minimum (< 0.6 %) during Autumn have been observed.

## 4. Time Series




Figure 5 - Time series of C<sub>2</sub>H<sub>6</sub> total column (in blue), low-tropospheric (3.58-6.79 km, in green) and UTLS (8.45-14.3 km, in pink) partial columns above Jungfraujoch. Red lines are linear trends.

The Figure 5 displays our retrieved C<sub>2</sub>H<sub>6</sub> total column and both partial columns (low-tropospheric and UTLS) above Jungfraujoch. We computed an overall decrease in ethane concentrations since 1994 of -14, -9 and -39 % resp. for our three columns. Trends have been determined using the bootstrap resampling tool developed by Gardiner (2008) (see Table III).

The decrease of ethane remains smaller than the seasonal amplitude. Our measurements allow to characterize strong seasonal variations of C<sub>2</sub>H<sub>6</sub> total and low-tropospheric columns with a maximum generally observed around mid-February. On average, the peak-to-peak amplitudes respectively amount to 50 and 76 % of the 1994 reference column.

Whereas the seasonal change of ethane UTLS column is less obvious with a peak-to-peak amplitude of 25 % (Ref : 1994 column). Therefore, the observed overall decrease of 39 % prevails in the UTLS region.

Time Series	Total Column	3.58 - 6.79 km	8.45 - 14.3 km
Jungfraujoch (1994-2011)	-0.47 ± 0.35 % (1994)	-0.92 ± 0.30 % (1994)	-0.69 ± 0.24 % (1994)
Jungfraujoch (1994-2008)	-1.06 ± 0.31 % (1994)	-	-
CHASER (1994-2008)	0.55 ± 0.18 % (1994)	-	-
Jungfraujoch (1998-2005)	-1.58 ± 0.68 % (1998)	-0.79 ± 0.98 % (1998)	-2.59 ± 0.89 % (1998)
Oslo's CTM (1998-2005)	-0.89 ± 0.56 % (1998)	-0.72 ± 0.57 % (1998)	-1.26 ± 1.23 % (1998)

Table III - Annual Change (in %), its 2- $\sigma$  uncertainties and its reference year for Jungfraujoch, CHASER Model and Oslo's CTM v.3 time series for our three columns.

On Table III, we notice a good agreement, significant within 2- $\sigma$  between Oslo's CTM computed trends and the trends of our retrieved columns.

On Figure 6, we compare monthly means of an averaged year of our Jungfraujoch's observations and model data issued from the CHASER 3D-Model and Oslo's CTM v.3. We notice both models underestimate the amount of ethane in the atmosphere. In the case of Oslo's model, it may be explained by the use of undervalued atmospheric pressures.

C<sub>2</sub>H<sub>6</sub> seasonal change has been evaluated for both CHASER and Oslo's CTM data with a peak-to-peak amplitude of 35 % (Ref. : 1994) and of 44 % (Ref. : 1998), respectively for ethane total columns. While our low-tropospheric peak-to-peak amplitude amounts to 67 % for the CHASER Model and for 51 % for Oslo's CTM.

## 5. Model Comparison




Figure 6 - C<sub>2</sub>H<sub>6</sub> three columns typical year for Jungfraujoch observations (in blue), results from CHASER Model (in pink) and from Oslo's CTM v.3 (in green). Total columns are respectively averaged on the 1994-2011, 1970-2008 and 1998-2005 time periods while CHASER's partial columns are computed over the 2007-2009 time period.

## 6. Conclusion

Harrison's new ethane parameters coupled to Hitran 2008 compilation improve the retrieval of ethane in terms of spectral residuals and information content ; as well as Chelin's O<sub>3</sub> and Bray's CH<sub>3</sub>Cl updates.

The selected a priori VMR profiles issued from the CHASER Model gives the least negative profiles with best residuals and DOFS.

Concerning the long-term trend of C<sub>2</sub>H<sub>6</sub>, we determined a significant decrease in its concentration over the 1994-2011 time period. We also characterized a seasonal change in total and tropospheric columns of ethane greater than the previously mentioned decrease. In the UTLS the long-term decrease is as the same order of magnitude than the seasonal modulation.

We expect to put more investigations into the study of the interannual variations as well as of the seasonal change of ethane in the lower-troposphere and in the UTLS region.

We still have to evaluate the impact of spectroscopy, instrumental line shape, geometry in order to refine our error budget.

## Acknowledgments

The University of Liège involvement has primarily been supported by the PRODEX program funded by the Belgian Federal Science Policy Office Brussels and by the Swiss GAW-CH program. E. Mahieu is Research Associate with the F.R.S. - FNRS. The FRS-FNRS and the Fédération Wallonie-Bruxelles are further acknowledged for observational activities support. We thank the International Foundation High Altitude Research Stations Jungfraujoch and Gornergrat (HFSJG, Bern) for supporting the facilities needed to perform the observations. We further acknowledge the vital contribution from all the Belgian colleagues in performing the Jungfraujoch observations used here. We further thank G.C. Toon (NASA-JPL, Pasadena) for the conversion of the ethane cross sections into pseudolines which can be used by our retrieval algorithm. We finally thank K. Sudo and O. A. Søvde for providing synthetic data.