

Mechanism of lipolysis in milk : a modelistic approach using a Langmuir film balance.

Authors : Sabine Braipson-Danthine, Christophe Blecker.

Affiliation : Gembloux agricultural University, Department of Food Technology (Head : Prof. C.Deroanne)

Support : Gembloux Agricultural University, Passage des déportés 2 B-5030 Gembloux Belgium

Abstract : Lipases are enzymes that specifically act on aggregated lipids in an aqueous environment and more accurately at interfaces. In order to explain the pathway of lipolysis, a reversible adsorption or penetration of the enzyme into the interface has been proposed. This step is supposed to precede the formation of the enzyme-substrate complex. Studies using monolayers have shown that activity of lipolytic enzymes are greatly influenced by "the interfacial quality". For reasons of simplification, most studies of model interfaces have involved only 1 or 2 type of lipid and/or protein. Generally all biological interfacial regions are however composed of a complex mixture of lipids and proteins. In milk and dairy cream, the interfacial region between milk fat and the aqueous phase where lipases may be present is the milk fat globule membrane (MFGM). This MFGM is a complex combination of proteins and phospholipids. The main purpose of this work was to describe the interactions between a MFGM monolayer and 3 kind of lipases in order to investigate the mechanism of lipolysis in milk. The monolayer technique used in this work has the advantage that the arrangement of the molecules can be controlled by changing the molecular area and the surface pressure of the monolayer. So this monolayer technique was used to provide a model system where the quality of the interface can be controlled. The lipases studied in this work are representative of those which are responsible for lipolysis in milk products : pancreatic lipase (PPL) and milk lipoprotein lipase (MLPL) ; these two enzymes are close in term of structure and activity and also a lipase from *Pseudomonas fluorescens* (PFL) (a typical bacterium of refrigerated milk). MFGM was first extracted from raw cream and compositional analyses were performed to confirm that material was representative of that which is really present around milk fat globules. Interaction of the 3 lipases with MFGM depends on surface pressure of MFGM monolayer, and thus on its organisation. PFL was able to interact with MFGM monolayer compressed at surface pressure up to 30 mN/m while PPL and MLPL are unable to interact with MFGM monolayer compressed at surface pressure higher than 25mN/m. This can be related to the fact that lipolysis in milk by endogenous lipase occurs only if MFGM has been deteriorated, on the other hand microbial lipases could be efficient even if the fat globules are intact.