

Abelian symmetries in multi-Higgs-doublet models

Igor P. Ivanov^{1,2}, Venus Keus¹, Evgeny Vdovin²

¹ IFPA, Université de Liège, Allée du 6 Août 17, bâtiment B5a, 4000 Liège, Belgium

² Sobolev Institute of Mathematics, Koptyug avenue 4, 630090, Novosibirsk, Russia

December 8, 2011

Abstract

N -Higgs doublet models (NHDM) are a popular framework to construct electroweak symmetry breaking mechanisms beyond the Standard model. Usually, one builds an NHDM scalar sector which is invariant under a certain symmetry group. Although several such groups have been used, no general analysis of symmetries possible in the NHDM scalar sector exists. Here, we describe a strategy that identifies all abelian groups which are realizable as symmetry groups of the NHDM Higgs potential. We consider both the groups of Higgs-family transformations only and the groups which also contain generalized CP transformations. We illustrate this strategy with the examples of 3HDM and 4HDM and prove several statements for arbitrary N .

1 Introduction

The Higgs mechanism of the electroweak symmetry breaking (EWSB) has not yet been discovered experimentally, although the LHC is expected to provide soon first definite clues about the nature of the EWSB. Many different variants of the Higgs mechanism have been proposed so far [1]. Moreover, even if one focuses on a specific Higgs mechanism beyond the Standard Model (SM), one has many free parameters describing the scalar and Yukawa sectors, all of them having phenomenological consequences. Ideally, one would want to have a formalism that reconstructs the phenomenology of a given model in its most general form, that is, in the entire space of the allowed free parameters. Unfortunately, such a construction seems impossible for most non-minimal Higgs mechanisms. In this situation the standard approach is to narrow down the freedom by imposing certain symmetries both on the Higgs sector and on the Yukawa interactions. The problem is that in most cases it is not known a priori which symmetries can be used. Many specific symmetry groups have been analyzed but their complete lists for every class of non-minimal Higgs model are still lacking.

The general task of classifying symmetries in models beyond SM can be, roughly, separated into two subtasks. First, one needs to know which symmetries can be implemented in the scalar sector of the model and how these symmetries are broken under EWSB. Second, one has to understand how these symmetry patterns can be encoded in the Yukawa sector of the model and what are the resulting properties of the fermions. In this paper we exclusively focus on the first task.

1.1 Realizable symmetries

When searching for the symmetry group which can be implemented in the scalar sector of a non-minimal Higgs model, one must distinguish between *realizable* and *non-realizable* groups. If it is possible to write a potential which is symmetric under a group G but not symmetric under a larger symmetry group containing it, we call G a realizable group (the exact definition will be given in Section 2.3). A non-realizable group, on the contrary, automatically leads to a larger symmetry group of the potential. In other words, the true symmetry properties of the potentials are reflected in realizable groups.

Just to give an example, cyclic groups such as \mathbb{Z}_7 and \mathbb{Z}_{13} have been used in the context of the flavor symmetry problem, see e.g. [2] and references therein. These symmetry groups are usually implemented in the space of fermion families, but sometimes they are transferred to the Higgs sector of the model. Anticipating the results of this paper, we can state that these symmetry groups are not realizable in the three-Higgs-doublet model potential: trying to impose such a symmetry, one ends up with a potential that has a continuous symmetry.

Therefore, if one focuses on the scalar sector of the model and aims at classifying the possible groups which can be implemented in the scalar potential, it is natural to restrict one's choice to the realizable symmetry groups.

1.2 Multi-Higgs-doublet models

One particular non-minimal Higgs sector where the classification of possible symmetries of the scalar sector has been established is the two-Higgs-doublet (2HDM), [3–5]. Although the general Higgs potential in 2HDM cannot be minimized explicitly with the straightforward algebra, it can still be analyzed fairly completely with several basis-invariant tools that were developed in the last years. The basic idea is that reparametrization transformations, or the Higgs-basis changes, link differently looking but physically equivalent potentials to each other. This idea can be implemented via the tensorial formalism at the level of Higgs fields, [6–9], or via geometric constructions in the space of gauge-invariant bilinears, [10–15]. In particular, all possible groups of Higgs-family symmetries and generalized CP -transformations of the 2HDM potential have been found. In addition, it was noted that if one disregards the $U(1)$ gauge transformations the structure of the Higgs potential of 2HDM is invariant under a wider class of transformations, [16].

It is natural to attempt to extend these successful techniques to N Higgs doublets. Some properties of the general NHDM potential were analyzed in [12, 17–20]. In particular, the bilinear-space geometric approach was recently adapted to the NHDM in [23, 24]. In what concerns the symmetries of the scalar sector of NHDM, several groups have attacked various issues of this problem, [20–22, 24–27]. In particular, in [20] an accurate and lengthy analysis of possible textures of the quartic interaction in the 3HDM led to the list of one-parametric symmetries of the potential. Although the method of [20] could be extended to $N > 3$, its practical implementation relies on the set of all possible textures for the quartic coupling constants, which is not known *a priori*.

In [24] an attempt was made to describe possible symmetry groups of NHDM in the adjoint representation. The symmetry properties of the vectors M_i and Λ_{0i} and the symmetric tensor Λ_{ij} in the expression (4) written below could be easily identified, but the main difficulty was to take into account the non-trivial shape of the NHDM orbit space in the adjoint space.

Mathematically, this led to the problem of classifying intersections of two orthogonal subgroups of $SO(N^2 - 1)$, which was not solved in [24].

It is therefore fair to state that despite several partial successes the task of classifying all possible Higgs-family and generalized- CP symmetries in NHDM is far from being accomplished.

In this paper we make a step towards solving this problem. We focus on abelian symmetries of the NHDM scalar potential and describe a strategy that algorithmically identifies realizable abelian symmetry groups for any given N . The strategy first addresses groups of unitary transformations in the space of Higgs families and then it is extended to groups that include antiunitary (generalized CP) transformations. The strategy also yields explicit examples of the potentials symmetric under a given realizable group.

The structure of the paper is the following. In Section 2 we describe the Higgs potential of the general NHDM, discuss the group of reparametrization transformations and introduce the notion of realizable symmetries. In Section 3 we describe the strategy that identifies all realizable abelian symmetry groups. Then in Section 4 we illustrate the general approach with the examples of 3HDM and 4HDM. Section 5 contains some further general results valid for all N . Finally, in Section 6 we make several remarks and draw our conclusions. In the Appendices we prove the theorem about maximal abelian groups of $PSU(N)$ and work out in detail the groups containing antiunitary transformations in the case of 3HDM.

2 Scalar sector of the N -Higgs doublet model

2.1 NHDM potential

In the NHDM we introduce N complex Higgs doublets with electroweak isospin $Y = 1/2$:

$$\phi_a = \begin{pmatrix} \phi_a^+ \\ \phi_a^0 \end{pmatrix}, \quad a = 1, \dots, N. \quad (1)$$

Hermitian conjugation is denoted by dagger: ϕ_a^\dagger . The generic renormalizable Higgs potential in NHDM can be written in the tensorial form [1, 4]:

$$V = Y_{ab}(\phi_a^\dagger \phi_b) + Z_{abcd}(\phi_a^\dagger \phi_b)(\phi_c^\dagger \phi_d), \quad (2)$$

where all indices run from 1 to N . It is constructed from N^2 gauge-invariant bilinears $(\phi_a^\dagger \phi_b)$, which uniquely parametrize the gauge orbits, and therefore it can be viewed as defined in the orbit space. Coefficients in the quadratic and quartic parts of the potential are grouped into components of tensors Y_{ab} and Z_{abcd} , respectively; there are N^2 independent components in Y and $N^2(N^2 + 1)/2$ independent components in Z . Thus, every conceivable Higgs sector in NHDM can be viewed as a point in the $N^2(N^2 + 3)/2$ -dimensional parameter space.

A convenient way to parametrize N^2 bilinears is to introduce the real vector $r^\mu = (r_0, r_i)$ with

$$r_0 = \sqrt{\frac{N-1}{2N}} \phi_a^\dagger \phi_a, \quad r_i = \frac{1}{2} \phi_a^\dagger (\lambda_i)_{ab} \phi_b, \quad (3)$$

where λ_i are the standard $SU(N)$ generators. The Higgs potential can be rewritten as

$$V = -M_0 r_0 - M_i r_i + \frac{1}{2} \Lambda_{00} r_0^2 + \Lambda_{0i} r_0 r_i + \frac{1}{2} \Lambda_{ij} r_i r_j, \quad (4)$$

which, if needed, can be represented even more compactly as $V = -M_\mu r^\mu + \Lambda_{\mu\nu} r^\mu r^\nu / 2$. All the coefficients Y_{ab} and Z_{abcd} become now components of the vector M_μ and symmetric tensor $\Lambda_{\mu\nu}$, which transform under adjoint representation of the group $GL(N, \mathbb{C})$. For more details about the properties of the Higgs potential in the space of bilinears, see [23, 24].

2.2 The group of reparametrization transformations

When discussing symmetries of the potential, we focus on the reparametrization transformations, which are non-degenerate linear transformations mixing different doublets ϕ_a but keeping invariant the kinetic term (which includes interaction of the Higgs fields with the gauge sector of the model). Alternatively, they can be defined as norm-preserving transformations of doublets that do not change the intradoublet structure. In this work we do not use the more general transformations in the spirit of [16] but focus on the ‘‘classically defined’’ reparametrization transformations.

A reparametrization transformation must be unitary (a Higgs-family transformation) or antiunitary (a generalized CP -transformation):

$$U : \phi_a \mapsto U_{ab} \phi_b \quad \text{or} \quad U_{CP} = U \cdot J : \phi_a \mapsto U_{ab} \phi_b^\dagger, \quad (5)$$

with a unitary matrix U_{ab} . The transformation $J \equiv CP$ acts on doublets by complex conjugation and satisfies $J^2 = 1$.

Let us focus first on the unitary transformations U . A priori, such transformations form the group $U(N)$. However, the overall phase factor multiplication is already taken into account by the $U(1)_Y$ from the gauge group. This leaves us with the $SU(N)$ group of reparametrization transformations. Then, this group has a non-trivial center $Z(SU(N)) = \mathbb{Z}_N$ generated by the diagonal matrix $\exp(2\pi i/N) \cdot 1_N$, where 1_N is the identity matrix. Therefore, the group of *physically distinct* unitary reparametrization transformations is

$$G_u = PSU(N) \simeq SU(N)/\mathbb{Z}_N. \quad (6)$$

Consider now antiunitary transformations $U_{CP} = U \cdot J$, $U \in U(N)$. One can define an action of J on the group $U(N)$ by

$$J \cdot U \cdot J = U^*, \quad (7)$$

where asterisk denoted complex conjugation. This action leaves invariant both the overall phase subgroup $U(1)_Y$ and the center of the $SU(N)$. Therefore, we again can consider only such $U_{CP} = U \cdot J$ that $U \in PSU(N)$. Once this action is defined, we can represent all distinct reparametrization transformations as a semidirect product

$$G = PSU(N) \rtimes \mathbb{Z}_2. \quad (8)$$

2.3 Realizable symmetry groups

A (reparametrization) *symmetry* of the Higgs lagrangian is such a reparametrization transformation that leaves the potential invariant: $V(U_{(CP)}\phi) = V(\phi)$. For any given potential all such transformations form a group, the automorphism group of the potential, which we denote as $G_V = Aut(V) \cap G$ and which is obviously a subgroup of G in (8).

Let us stress that when we say a group G_V is a symmetry group of the potential V , we mean that it is equal to, not just contained in, $Aut(V) \cap G$. In [24] it was suggested to call such groups *realizable* symmetry groups. In simple words, a symmetry group G_V is realizable if one can write a potential symmetric under G_V but not symmetric under any overgroup of G_V . For example, a potential which depends on the first doublet only via $(\phi_1^\dagger \phi_1)$ is obviously symmetric under the cyclic group \mathbb{Z}_n of discrete phase rotations of this doublet for any n . However, these \mathbb{Z}_n 's have no interest on their own because they trivially arise as subgroups of the larger symmetry group of this potential $U(1)$ describing arbitrary phase rotations. It is this $U(1)$, not its individual subgroups \mathbb{Z}_n , which has a chance to be the realizable symmetry group of this potential.

Since in this paper we deal with groups A_u containing only unitary transformations of Higgs families and with groups A that can contain antiunitary transformations, let us adopt the following convention: a group $A_u \in G_u$ is called realizable if $A_u = Aut(V) \cap G_u$, and a group $A \in G$ is called realizable if $A = Aut(V) \cap G$.

Note that in the case of 2HDM, where all possible reparametrization symmetries are known, \mathbb{Z}_2 , $(\mathbb{Z}_2)^2$, $(\mathbb{Z}_2)^3$, $U(1)$, $U(1) \times \mathbb{Z}_2$, $SU(2)$, each of these symmetry groups is realizable, which is most easily proved in the space of bilinears.

3 Finding abelian groups

We now focus on abelian groups for NHDM and describe in this Section the strategy that identifies all realizable abelian groups of the NHDM potential for any N . This strategy can be outlined as follows: we first describe maximal abelian subgroups of $PSU(N)$, then we explore their realizable subgroups, and finally we check which of these groups can be extended by including antiunitary transformations.

3.1 Maximal abelian subgroups of $PSU(N)$

A maximal abelian subgroup of G_u (6) is an abelian group that is not contained in a larger abelian subgroup of G_u . A priori, there can be several maximal abelian subgroups in a given group. Any subgroup of G_u must be either a maximal one, or lie inside a maximal one. Therefore, we first need to identify all maximal abelian subgroups of G_u and then study their realizable subgroups.

If G_u were $SU(N)$, then the situation would be simple. As we describe in Appendix A all maximal abelian subgroups of $SU(N)$ are the so called *maximal tori*,

$$[U(1)]^{N-1} = U(1) \times U(1) \times \cdots \times U(1), \quad (9)$$

and all such maximal tori are conjugate inside $SU(N)$. It means that without loss of generality one could pick up a specific maximal torus, for example, the one that is represented by phase rotations of individual doublets

$$\text{diag}[e^{i\alpha_1}, e^{i\alpha_2}, \dots, e^{i\alpha_{N-1}}, e^{-i\sum \alpha_i}], \quad (10)$$

and study its subgroups.

However, the group of distinct reparametrizations is $G_u = PSU(N)$, and it has a richer structure. Consider the canonical epimorphism

$$\overline{} : SU(N) \rightarrow SU(N)/Z(SU(N)) = \overline{SU(N)} \simeq PSU(N), \quad (11)$$

where $Z(SU(N)) \simeq \mathbb{Z}_N$ is the center of $SU(N)$. Given subgroup \overline{A} of $PSU(N)$ we denote by $A \in SU(N)$ the complete preimage of \overline{A} . If \overline{A} is abelian, its complete preimage A *does not have to be abelian*. Specifically, in Appendix A we prove the following Theorem:

Theorem 1. *Let \overline{A} be a maximal abelian subgroup of $PSU(N)$, denote by A a complete preimage of \overline{A} in $SU(N)$. Then one of the following statements holds.*

- (1) *A is abelian and A is conjugate to the subgroup of all diagonal matrices in $SU(N)$.*
- (2) *A is a finite nilpotent group of class 2, the exponent of A is divisible by N and divides N^2 . Moreover, $Z(A) = Z(SU(N))$ and the commutator $[A, A] \leq Z(A)$.*

This theorem states that there are two sorts of maximal abelian groups in $PSU(N)$: (1) maximal tori, which will be constructed below, (2) certain finite abelian groups, which are not subgroups of maximal tori and which must be treated separately.

3.2 Maximal tori in $PSU(N)$

Let us explicitly construct the maximal torus in $PSU(N)$ referred to in Theorem 1. We first introduce some convenient notation. A diagonal unitary matrix acting in the space of Higgs doublets and performing phase rotations of individual doublets, such as (10), will be written as a vector of phases:

$$\left(\alpha_1, \alpha_2, \dots, \alpha_{N-1}, -\sum \alpha_i \right). \quad (12)$$

In addition, if M_1, \dots, M_k are subsets of a group G , then $\langle M_1, \dots, M_k \rangle$ denotes the subgroup generated by $M_1 \cup \dots \cup M_k$ of G . Then we construct a maximal torus in $SU(N)$

$$T_0 = \langle U(1)_1, U(1)_2, \dots, U(1)_{N-1} \rangle, \quad (13)$$

where

$$\begin{aligned} U(1)_1 &= \alpha_1(-1, 1, 0, 0, \dots, 0), \\ U(1)_2 &= \alpha_2(-2, 1, 1, 0, \dots, 0), \\ U(1)_3 &= \alpha_3(-3, 1, 1, 1, \dots, 0), \\ &\vdots \\ U(1)_{N-1} &= \alpha_{N-1}(-N+1, 1, 1, 1, \dots, 1), \end{aligned} \quad (14)$$

with all $\alpha_i \in [0, 2\pi)$. Clearly, for every j we have $\langle U(1)_1, \dots, U(1)_{j-1} \rangle \cap U(1)_j = \{e\}$, whence T_0 is a direct product of $U(1)_1, \dots, U(1)_{N-1}$. In particular, any element $u \in SU(N)$ can be uniquely written as

$$u = u_1(\alpha_1)u_2(\alpha_2) \cdots u_{N-1}(\alpha_{N-1}), \quad u_i \in U(1)_i. \quad (15)$$

Moreover, the center $Z(SU(N))$ is contained in the last group and is generated by $\alpha_{N-1} = 2\pi/N$. One can therefore introduce $\overline{U(1)}_{N-1} = U(1)_{N-1}/Z(SU(N))$, which can be parametrized as

$$\overline{U(1)}_{N-1} = \alpha_{N-1} \left(-\frac{N-1}{N}, \frac{1}{N}, \dots, \frac{1}{N} \right), \quad (16)$$

where $\alpha_{N-1} \in [0, 2\pi)$. Since $Z(SU(N)) \cap U(1)_j = 1$ for $j = 1, \dots, N-2$, we identify these subgroups with their homomorphic images and omit the bar. Thus the homomorphic image T of T_0 can be written as

$$T = U(1)_1 \times U(1)_2 \times \dots \times \overline{U(1)}_{N-1}. \quad (17)$$

3.3 Identifying symmetries of the potential

Next we study which subgroups of the maximal torus T can be realizable in the scalar sector of NHDM.

We start from the most general T -symmetric potential:

$$V(T) = - \sum_a m_a^2 (\phi_a^\dagger \phi_a) + \sum_{a,b} \lambda_{ab} (\phi_a^\dagger \phi_a) (\phi_b^\dagger \phi_b) + \sum_{a \neq b} \lambda'_{ab} (\phi_a^\dagger \phi_b) (\phi_b^\dagger \phi_a), \quad (18)$$

where among the $N(N-1)/2$ terms in the last sum there are only $2N-3$ algebraically independent ones, [23]. Each term in this potential transforms trivially under the entire T . The important fact now is that a sufficiently general potential of this form has no other unitary symmetry. In fact, an even stronger statement is true:

Theorem 2. *Consider potential $V(T)$, (18), defined by the set of coefficients m_a^2 , λ_{ab} , λ'_{ab} .*

1. *There exist coefficients such that the only unitary symmetries of the potential are the phase rotations from T .*
2. *If one constructs $V = V(T) + V_1$ by adding further terms that were absent in (18), then for any V_1 there still exist coefficients of $V(T)$ such that any unitary symmetry of V belongs to T .*

Proof. The first assertion is a particular case of the second, which we now prove. Consider a generic potential V in the adjoint space (4). Any unitary transformation of the Higgs families corresponds to an orthogonal transformation in the r_i -space. Therefore, any unitary symmetry of the potential must leave invariant each of the five terms in (4). Consider, specifically, the second and the fourth terms:

$$V = -Y_{ab} (\phi_a^\dagger \phi_b) + Z_{ab} (\phi_a^\dagger \phi_b) \sum_c (\phi_c^\dagger \phi_c) + \dots, \quad (19)$$

where we omitted the remaining terms and introduced the tensor

$$Z_{ab} = \frac{2}{N} \sum_d Z_{abdd} - \frac{2}{N^2} \delta_{ab} \sum_{c,d} Z_{ccdd}, \quad (20)$$

which is basically the $Z_{ab}^{(2)}$ of [7] made traceless. Both Y_{ab} and Z_{ab} are hermitean matrices, and their diagonal values can be arbitrarily changed by changing m_a^2 and λ_{ab} in (18). In particular,

one can adjust them so that the spectra of Y_{ab} and Z_{ab} are non-degenerate. Their eigenvectors can be then used to construct two orthonormal bases in the space of Higgs doublets, $\{e_i^{(Y)}\}$ and $\{e_i^{(Z)}\}$, respectively ($i = 1, \dots, N$). Let us also introduce the canonical basis $\{e_i^{(0)}\}$ consisting of vectors of the form $(0, \dots, 1, \dots, 0)$, so that the maximal torus T consists of phase rotations of these vectors.

The symmetry group of the first term in (19) is the maximal torus T_Y of phase rotations of individual eigenvectors $e_i^{(Y)}$. The symmetry group of the second term in (19) is the maximal torus T_Z of phase rotations of individual eigenvectors $e_i^{(Z)}$. The symmetry group of (19) is, therefore, $T_Y \cap T_Z$, the phase rotations of common eigenvectors of the two matrices (and the symmetry group of the entire V is its subgroup).

Take an eigenvector of Y_{ab} . Its orientation can be changed by readjusting coefficients m_a^2 if and only if it does not belong to the canonical basis. The same applies to Z_{ab} . Therefore, even if Y_{ab} and Z_{ab} had common eigenvectors not belonging to the canonical basis, one could avoid this coincidence by readjusting the coefficients m_a^2 and λ_{ab} . After this procedure, the only common eigenvectors of Y_{ab} and Z_{ab} will be those belonging to the canonical basis. Therefore, the symmetry group of (19), and consequently, of the full potential will be a subgroup of T . \square

The theorem just proved guarantees that when we start from the T -symmetric potential (18) and add more terms, we will never generate any new unitary symmetry that was not already present in T . This is the crucial step in proving that the groups described below are realizable. Our task now is to find which subgroups of T can be obtained in this way.

Consider a bilinear $\phi_a^\dagger \phi_b$, $a \neq b$. It gains a phase change under T (17) which linearly depends on the angles α_i :

$$\phi_a^\dagger \phi_b \rightarrow \exp[i(p_{ab}\alpha_1 + q_{ab}\alpha_2 + \dots + t_{ab}\alpha_{N-1})] \cdot \phi_a^\dagger \phi_b \quad (21)$$

with some integer coefficients $p_{ab}, q_{ab}, \dots, t_{ab}$. Note that all coefficients are antisymmetric in their indices: $p_{ba} = -p_{ab}$, etc. These coefficients can be represented by real antisymmetric matrices with integer values, or graphically, as labels of the edges of $N - 1$ *oriented graphs*, one for each $U(1)$ group. Each such graph has N vertices, corresponding to doublets ϕ_a ; all vertices are joined with arbitrarily oriented edges, orientation indicated by an arrow. An edge oriented from ϕ_b towards ϕ_a (edge $b \rightarrow a$) is associated with the bilinear $\phi_a^\dagger \phi_b$ and is labeled by p_{ba} in the first graph, q_{ba} in the second graph, etc. Examples of these graphs will be shown later for 3HDM and 4HDM.

The Higgs potential is a sum of monomial terms which are linear or quadratic in $\phi_a^\dagger \phi_b$. Consider one such term and calculate its coefficients p, \dots, t . Let us first focus on how this term depends on any single $U(1)_i$ subgroup of T . There are two possibilities depending on the value of the i -th coefficient:

- If the coefficient k in front of α_i is zero, this term is $U(1)_i$ -symmetric.
- If the coefficient $k \neq 0$, then this term is symmetric under the $\mathbb{Z}_{|k|}$ subgroup of $U(1)_i$ -group generated by phase rotations by $2\pi/|k|$.

However, even if a given monomial happens to have finite symmetry groups with respect to each single $U(1)_i$, its symmetry group under the entire T is still continuous: for any set of coefficients one can adjust angles α_i in such a way that $p_{ab}\alpha_1 + \dots + t_{ab}\alpha_{N-1} = 0$. Therefore,

when studying symmetries of a given term or a sum of terms, we cannot limit ourselves to individual $U(1)_i$ groups but must consider the full maximal torus.

The strategy presented below guarantees that we find all possible realizable subgroups of the maximal torus T , both finite and infinite.

Consider a Higgs potential V which, in addition to the T -symmetric part (18) contains $k \geq 1$ additional terms, with coefficients p_1, q_1, \dots, t_1 to p_k, q_k, \dots, t_k . This potential defines the following $(N - 1) \times k$ matrix of coefficients:

$$X(V) = \begin{pmatrix} p_1 & q_1 & \cdots & t_1 \\ p_2 & q_2 & \cdots & t_2 \\ \vdots & \vdots & & \vdots \\ p_k & q_k & \cdots & t_k \end{pmatrix}. \quad (22)$$

The symmetry group of this potential can be derived from the set of non-trivial solutions for α_i of the following equations:

$$X(V) \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_{N-1} \end{pmatrix} = \begin{pmatrix} 2\pi n_1 \\ \vdots \\ 2\pi n_k \end{pmatrix}, \quad (23)$$

There are two major possibilities depending on the rank of this matrix.

- **Finite symmetry group.** If $\text{rank} X(V) = N - 1$, then there is no non-trivial solution of the equation (23) with the trivial right-hand side (i.e. all $n_i = 0$). Instead, there exists a unique solution for any non-trivial set of n_i , and all such solutions form the finite group of phase rotations of the given potential.

To find which symmetry groups can be obtained in this way, we take exactly $N - 1$ monomials, so that the matrix $X(V)$ becomes a square matrix with a non-zero determinant. It is known that any square matrix with integer entries can be diagonalized by a sequence of the following operations: swapping two rows or two columns, adding a row (a column) to another row (column) and multiplying a row (a column) by -1 . After diagonalization, the matrix $X(V)$ becomes $\text{diag}(d_1, \dots, d_{N-1})$, where d_i are positive non-zero integers. This matrix still defines the equation (23) for α'_i , which are linear combinations of α_i 's and with $n'_i \in \mathbb{Z}$. Therefore, the finite symmetry group of this matrix is $\mathbb{Z}_{d_1} \times \cdots \times \mathbb{Z}_{d_{N-1}}$ (where \mathbb{Z}_1 means no non-trivial symmetry).

Note also that each of the allowed manipulations conserves the absolute value of the determinant of $X(V)$. Therefore, even before diagonalization one can calculate the order of the finite symmetry group as $|\det X(V)|$.

This derivation leads us to the strategy that identifies all finite subgroups of torus realizable as symmetry groups of the Higgs potential in NHDM: write down all possible monomials with N -doublets, consider all possible subsets with exactly $N - 1$ distinct monomials, construct the matrix X for this subset and find its symmetry group following the above scheme. Although this strategy is far from being optimal, its algorithmic nature allows it to be easily implemented in a machine code.

- **Continuous symmetry group.** If $\text{rank}X(V) < N-1$, so that $D = N-1-\text{rank}X(V) > 0$, then there exists a D -dimensional subspace in the space of angles α_i , which solves the equation (23) for the trivial right-hand side. One can then focus on the orthogonal complement of this subspace, where no non-trivial solution of the homogeneous equation is possible, and repeat the above strategy to find the finite symmetry group G_D in this subspace. The symmetry group of the potential is then $[U(1)]^D \times G_D$.

3.4 Antiunitary transformations

Having found the list of realizable subgroups of torus T , we can ask whether these groups can be extended to larger abelian groups that would include not only unitary but also antiunitary transformations. Here we describe the strategy that solves this problem.

As described above, one can define the action of the CP -transformation J on $PSU(N)$ given by (7): $J \cdot U \cdot J = U^*$. This action can be restricted to the maximal torus T (17) and even further, to any subgroup $A \subset T$ (indeed, if $a \in A$, then $a^* \in A$). However, J does not commute with a generic element of T , therefore the group $\langle A, J \rangle$ is not, in general, abelian.

In order to embed A in a larger abelian group, we must find an antiunitary transformation $J' = b \cdot J$ which commutes with any element $a \in A$:

$$J'a(J')^{-1} = a \quad \Leftrightarrow \quad bJaJb^{-1} = a \quad \Leftrightarrow \quad ba^{-1}b^{-1} = a \quad \Leftrightarrow \quad b = aba. \quad (24)$$

The last expression here is a linear matrix equation for the matrix b at any given a . Since a is diagonal, $(aba)_{ij} = a_{ii}a_{jj}b_{ij}$, so whenever $a_{ii}a_{jj} \neq 1$, one must set b_{ij} to zero.

If at least one matrix b satisfying (24) is found, all the other matrices can be constructed with the help of the last form of this equation. Suppose $b' = xb$ also satisfies this equation; then, $axa^{-1} = x$. Therefore, x can be any unitary matrix from $PSU(N)$ commuting with a , that is, x centralizes the chosen abelian group A .

In summary, the strategy for embedding a given abelian group $A \subset T$ into a larger abelian group A_{CP} which includes antiunitary transformations proceeds in five steps:

- Find one b which solves the matrix equation (24) for all $a \in A$.
- Find all $x \in PSU(N)$ which commute with each element of A , that is, construct the centralizer of A in $PSU(N)$.
- Find restrictions placed on x 's by the requirement that product of two different antiunitary transformations (x_1J') and (x_2J') belongs to A . This procedure can result in several distinct groups A_{CP} .
- Find which among the A -symmetric terms in the potential are also symmetric under some of A_{CP} ; drop terms which are not.
- Check that the resulting collection of terms is not symmetric under a larger group of *unitary* transformations, that is, A is still a realizable symmetry group of this collection of terms.

As an exercise, let us apply this strategy to the full maximal torus T . A generic $a \in T$ acting on a doublet ϕ_i generates a non-trivial phase rotation $\psi_i(\alpha_1, \dots, \alpha_{N-1})$ which can be reconstructed from (14). The equation (24) then becomes

$$e^{i(\psi_i+\psi_j)}b_{ij} = b_{ij}. \quad (25)$$

Since $\psi_i + \psi_j \neq 0$ for any i, j , the only solution to this equation is $b_{ij} = 0$ for all i and j . This means that there is no J' that would commute with every element of the torus T . Thus, T cannot be embedded into a larger abelian group T_{CP} that would include antiunitary transformations.

In order to avoid possible confusion, we note that in fact J is a symmetry of the T -symmetric potential. However, the symmetry group, generated by T and J , is a semidirect product $T \rtimes J$, and it is nonabelian.

4 Examples of 3HDM and 4HDM

In this Section we illustrate the general strategy presented above with the particular cases of 3HDM and 4HDM. For 3HDM we give the full list of abelian groups realizable as symmetry groups of the Higgs potential and show explicit examples of such potentials; for 4HDM we do it only for the finite groups of unitary transformations.

4.1 Maximal torus for 3HDM and its subgroups

In the 3HDM the representative maximal torus $T \subset PSU(3)$ is parametrized as

$$T = U(1)_1 \times U(1)_2, \quad U(1)_1 = \alpha(-1, 1, 0), \quad U(1)_2 = \beta \left(-\frac{2}{3}, \frac{1}{3}, \frac{1}{3} \right), \quad (26)$$

where $\alpha, \beta \in [0, 2\pi)$. The most general Higgs potential symmetric under T is

$$\begin{aligned} V = & -m_1^2(\phi_1^\dagger\phi_1) - m_2^2(\phi_2^\dagger\phi_2) - m_3^2(\phi_3^\dagger\phi_3) \\ & + \lambda_{11}(\phi_1^\dagger\phi_1)^2 + \lambda_{22}(\phi_2^\dagger\phi_2)^2 + \lambda_{33}(\phi_3^\dagger\phi_3)^2 \\ & + \lambda_{12}(\phi_1^\dagger\phi_1)(\phi_2^\dagger\phi_2) + \lambda_{23}(\phi_2^\dagger\phi_2)(\phi_3^\dagger\phi_3) + \lambda_{13}(\phi_3^\dagger\phi_3)(\phi_1^\dagger\phi_1) \\ & + \lambda'_{12}|\phi_1^\dagger\phi_2|^2 + \lambda'_{23}|\phi_2^\dagger\phi_3|^2 + \lambda'_{13}|\phi_3^\dagger\phi_1|^2. \end{aligned} \quad (27)$$

There are six bilinear combinations of doublets transforming non-trivially under T with coefficients

	p	q	
$(\phi_2^\dagger\phi_1)$	-2	-1	(28)
$(\phi_3^\dagger\phi_2)$	1	0	
$(\phi_1^\dagger\phi_3)$	1	1	

and their conjugates whose coefficients p and q carry opposite signed with respect to (28). These coefficients are shown graphically as labels of the edges of two oriented graphs shown in Fig. 1.

Before applying the general strategy described in the previous Section, let us check which finite symmetry groups can arise as subgroups of either $U(1)_1$ or $U(1)_2$ individually.

Consider first $U(1)_1$. The coefficient p of any monomial can be obtained either by picking the labels from the first graph in Fig. 1 directly or by summing two such labels, multiplied by -1 when needed. In this way we can obtain any $|p|$ from 0 to 4. For a monomial with $p = 0$, the symmetry group is the entire $U(1)_1$. For $|p| = 1$, there is no non-trivial symmetry. For $|p| = 2, 3, 4$, we obtain the cyclic group $\mathbb{Z}_{|p|}$. In each case it is straightforward to construct




Figure 1: Coefficients p and q as labels of triangles representing bilinears in 3HDM

the monomials with a given symmetry; for example, $(\phi_1^\dagger\phi_3)(\phi_2^\dagger\phi_3)$ and its conjugate are $U(1)_1$ -symmetric, while $(\phi_1^\dagger\phi_2)(\phi_1^\dagger\phi_3)$, $(\phi_1^\dagger\phi_2)(\phi_3^\dagger\phi_2)$ and their conjugates are symmetric under the \mathbb{Z}_3 -group with a generator

$$a = (\omega, 1, \omega^{-1}), \quad \omega \equiv \exp(2\pi i/3). \quad (29)$$

In the case of the group $U(1)_2$, the labels can sum up to $|q| = 0, 1, 2$. Thus, the largest finite group here is \mathbb{Z}_2 .

As we mentioned in the previous Section, one cannot limit oneself to subgroups of individual $U(1)_i$ factors or to direct products of such subgroups. In order to find all realizable groups, one has to write the full list of possible monomials and then calculate the symmetry group of all distinct pairs (for $N = 3$) of monomials. For example, if the two monomials are $v_1 = \lambda_1(\phi_1^\dagger\phi_2)(\phi_1^\dagger\phi_3)$ and $v_2 = \lambda_2(\phi_2^\dagger\phi_1)(\phi_2^\dagger\phi_3)$, then the matrix $X(v_1 + v_2)$ has form

$$X(v_1 + v_2) = \begin{pmatrix} 3 & 2 \\ -3 & -1 \end{pmatrix} \rightarrow \begin{pmatrix} 3 & 2 \\ 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}, \quad (30)$$

and it produces the symmetry group \mathbb{Z}_3 . The solution of the equation

$$X(v_1 + v_2) \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} 2\pi n_1 \\ 2\pi n_2 \end{pmatrix} \quad (31)$$

yields $\alpha = 2\pi/3 \cdot k$, $\beta = 0$, which implies the transformation matrix of the doublets (29).

In 3HDM there are, up to complex conjugation, three bilinears and nine products of two bilinears transforming non-trivially under T . It is a straightforward exercise to check all possible pairs of monomials; in fact, by using permutations of the doublets the number of truly distinct cases is rather small. This procedure reveals just one additional finite group $\mathbb{Z}_2 \times \mathbb{Z}_2$, which arises when at least two terms among $(\phi_1^\dagger\phi_2)^2$, $(\phi_2^\dagger\phi_3)^2$, $(\phi_3^\dagger\phi_1)^2$ are present. This group is simply the group of sign flips of individual doublets.

Thus, we arrive at the full list of subgroups of the maximal torus realizable as the symmetry groups of the Higgs potential in 3HDM:

$$\mathbb{Z}_2, \quad \mathbb{Z}_3, \quad \mathbb{Z}_4, \quad \mathbb{Z}_2 \times \mathbb{Z}_2, \quad U(1), \quad U(1) \times \mathbb{Z}_2, \quad U(1) \times U(1). \quad (32)$$

Most of these groups were identified in [20] in the search of “simple” symmetries of 3HDM scalar potential. In that work a symmetry were characterized not by its group, as in our paper, but by providing a single unitary transformation S and then reconstructing the pattern in the Higgs potential which arises after requiring that it is S -symmetric. In certain cases the authors of [20] found that the potential is symmetric under a larger group than $\langle S \rangle$, in accordance with the notion of realizable symmetry discussed above.

The explicit correspondence between the seven symmetries S_1, \dots, S_7 of [20] and the list (32) is the following:

$$\begin{aligned} S_1 &\rightarrow \mathbb{Z}_2, & S_2 &\rightarrow U(1) \text{ realized as } U(1)_2, & S_3 &\rightarrow \mathbb{Z}_3, \\ S_4 &\rightarrow \mathbb{Z}_4, & S_5 &\rightarrow U(1) \text{ realized as } U(1)_1, & S_6 &\rightarrow U(1) \times \mathbb{Z}_2, & S_7 &\rightarrow U(1) \times U(1). \end{aligned} \quad (33)$$

In addition to these symmetries, our list (32) contains one more group $\mathbb{Z}_2 \times \mathbb{Z}_2$, which was not found in [20] because it does not correspond to a ‘‘simple’’ symmetry.

Let us also explicitly write the potentials which are symmetric under each group in (32).

- $\mathbf{U}(1) \times \mathbf{U}(1) = \mathbf{T}$. The most general T -symmetric potential of 3HDM is given by (27).
- $\mathbf{U}(1)$. This group can be realized in two non-equivalent ways, which are conjugate either to $U(1)_1$ or $U(1)_2$ in (26). The distinction between the two realizations lies in the eigenspace decomposition: a $U(1)_2$ -type group has a two-(complex)-dimensional space of eigenvectors, while eigenspaces of a $U(1)_1$ -type group are all one-dimensional. This leaves more freedom in constructing a $U(1)_2$ -symmetric potential than a $U(1)_1$ -symmetric one. Specifically, the general $U(1)_1$ -invariant potential contains, in addition to (27), the following terms:

$$\lambda_{1323}(\phi_1^\dagger \phi_3)(\phi_2^\dagger \phi_3) + h.c. \quad (34)$$

while the general $U(1)_2$ -invariant potential contains

$$\begin{aligned} &-m_{23}^2(\phi_2^\dagger \phi_3) + \left[\lambda_{1123}(\phi_1^\dagger \phi_1) + \lambda_{2223}(\phi_2^\dagger \phi_2) + \lambda_{3323}(\phi_3^\dagger \phi_3) \right] (\phi_2^\dagger \phi_3) \\ &+ \lambda_{2323}(\phi_2^\dagger \phi_3)^2 + \lambda_{2323}(\phi_2^\dagger \phi_1)(\phi_1^\dagger \phi_3) + h.c. \end{aligned} \quad (35)$$

It must be stressed that these potentials are written for the specific convention of groups $U(1)_1$ and $U(1)_2$ used in (26). This convention reflected a specific basis used to describe the structure of the torus T . In other bases, the explicit terms symmetric under a $U(1)_1$ or $U(1)_2$ -type groups will look differently. For example, the term $(\phi_2^\dagger \phi_1)(\phi_3^\dagger \phi_1)$ is symmetric under a $U(1)_1$ -type transformation with phases $(0, \alpha, -\alpha)$. To this end, we note that reconstructing the symmetry group of any given potential is a separate and difficult task which is not addressed in the present paper (here we only classify realizable symmetries and give examples of their realizations).

- $\mathbf{U}(1) \times \mathbb{Z}_2$. Looking back to (34), one might be tempted to think that the true symmetry group of this term is not $U(1)$ but $U(1) \times \mathbb{Z}_2$, because this term is also invariant under sign flip of ϕ_3 . However, inside $SU(3)$, a transformation with phases $(0, 0, \pi)$ is equivalent to $(-\pi, \pi, 0)$ which is already included in $U(1)_1$.

A potential whose true symmetry group is $U(1) \times \mathbb{Z}_2$ is given by (35) with all coefficients set to zero except for λ_{2323} . The corresponding term, $(\phi_2^\dagger \phi_3)^2$, is symmetric not only under the full $U(1)_2$, but also under $(-\pi, \pi, 0)$, which generates a \mathbb{Z}_2 subgroups inside $U(1)_1$.

- \mathbb{Z}_4 . The potential symmetric under \mathbb{Z}_4 contains, in addition to (27), the following terms:

$$\lambda_{1323}(\phi_1^\dagger \phi_3)(\phi_2^\dagger \phi_3) + \lambda_{1212}(\phi_1^\dagger \phi_2)^2 + h.c. \quad (36)$$

In accordance with the general discussion, it must contain $N - 1 = 2$ distinct terms, as required for a finite symmetry group. Again, this set of terms is specific for the particular choice of the $(U(1)_1, U(1)_2)$ -basis on the maximal torus.

- \mathbb{Z}_3 . The potential symmetric under \mathbb{Z}_3 contains

$$\lambda_{1232}(\phi_1^\dagger\phi_2)(\phi_3^\dagger\phi_2) + \lambda_{2313}(\phi_2^\dagger\phi_3)(\phi_1^\dagger\phi_3) + \lambda_{3121}(\phi_3^\dagger\phi_1)(\phi_2^\dagger\phi_1) + h.c. \quad (37)$$

In fact, any pair of the three terms is already sufficient to define a \mathbb{Z}_3 -symmetric potential. Note also that different $(U(1)_1, U(1)_2)$ -basis on the maximal torus lead to the same \mathbb{Z}_3 -group, because inside $PSU(3)$ the following groups are isomorphic:

$$\langle(1, \omega, \omega^2)\rangle \simeq \langle(\omega, \omega^2, 1)\rangle \simeq \langle(\omega^2, 1, \omega)\rangle, \quad \omega \equiv \exp(2\pi i/3). \quad (38)$$

- $\mathbb{Z}_2 \times \mathbb{Z}_2$. This group can be realized simply as a group of independent sign flips of the three doublets (up to the overall sign flip). Every term in the potential must contain each doublet in pairs. In addition to (27), the $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetric potential can contain

$$\lambda_{1212}(\phi_1^\dagger\phi_2)^2 + \lambda_{2323}(\phi_2^\dagger\phi_3)^2 + \lambda_{3131}(\phi_3^\dagger\phi_1)^2 + h.c. \quad (39)$$

- \mathbb{Z}_2 . This group can be realized, for example, as a group of sign flips of the third doublet. The potential can contain any term where ϕ_3 appears in pairs.

4.2 The $\mathbb{Z}_3 \times \mathbb{Z}_3$ -group

The only finite abelian group that is not contained in any maximal torus in $PSU(3)$ is $\mathbb{Z}_3 \times \mathbb{Z}_3$. Although there are many such groups inside $PSU(3)$, all of them are conjugate. Thus, only one representative case can be considered, and we choose the following two generators

$$a = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \quad b = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad \omega = \exp\left(\frac{2\pi i}{3}\right). \quad (40)$$

A generic potential that stays invariant under this group of transformations is

$$\begin{aligned} V = & -m^2 \left[(\phi_1^\dagger\phi_1) + (\phi_2^\dagger\phi_2) + (\phi_3^\dagger\phi_3) \right] + \lambda_0 \left[(\phi_1^\dagger\phi_1) + (\phi_2^\dagger\phi_2) + (\phi_3^\dagger\phi_3) \right]^2 \\ & + \lambda_1 \left[(\phi_1^\dagger\phi_1)^2 + (\phi_2^\dagger\phi_2)^2 + (\phi_3^\dagger\phi_3)^2 \right] + \lambda_2 \left[|\phi_1^\dagger\phi_2|^2 + |\phi_2^\dagger\phi_3|^2 + |\phi_3^\dagger\phi_1|^2 \right] \\ & + \lambda_3 \left[(\phi_1^\dagger\phi_2)(\phi_1^\dagger\phi_3) + (\phi_2^\dagger\phi_3)(\phi_2^\dagger\phi_1) + (\phi_3^\dagger\phi_1)(\phi_3^\dagger\phi_2) \right] + h.c. \end{aligned} \quad (41)$$

with real $M_0, \lambda_0, \lambda_1, \lambda_2$ and complex λ_3 . It is also interesting to note that this group describes an abelian frustrated symmetry of the 3HDM potential, see details in [28].

The potential (41) is not symmetric under any continuous Higgs-family transformation, which can be proved, for example, using the adjoint representation of the bilinears described in [23] and briefly outlined in section 2.1. However it is symmetric under the exchange of any two doublets, e.g. $\phi_1 \leftrightarrow \phi_2$. Together with b , such transformations generate the group S_3 of permutations of the three doublets. Its action on elements of the \mathbb{Z}_3 generated by a is also naturally defined, therefore, the potential (41) is automatically symmetric under the group $\mathbb{Z}_3 \rtimes S_3$, which is non-abelian.

We can conclude that the symmetry group $\mathbb{Z}_3 \times \mathbb{Z}_3$ is not realizable for 3HDM.

4.3 Including antiunitary symmetries in 3HDM

In Section 3.4 we described the strategy of embedding an abelian group of unitary transformations into a larger abelian group which includes anti-unitary transformations. Applying this strategy to 3HDM for all groups from the list (32), we obtain the following additional realizable abelian groups:

$$\mathbb{Z}_2^*, \quad \mathbb{Z}_2 \times \mathbb{Z}_2^*, \quad \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2^*, \quad \mathbb{Z}_4^*, \quad (42)$$

where the asterisk indicates that the generator of the corresponding cyclic group is an anti-unitary transformation. Details of the derivation are presented in Appendix B. Here we just briefly explain why groups such as \mathbb{Z}_6^* , \mathbb{Z}_8^* , $U(1) \times \mathbb{Z}_2^*$ do not appear in this list. When we search for anti-unitary transformations commuting with the selected subgroup of T , we can indeed construct such groups at the price of imposing certain restrictions on the coefficients of the T -symmetric part (27). This makes the potential symmetric under a larger group of *unitary* transformations, which is nonabelian. According to our definition, this means that the groups such as \mathbb{Z}_6^* are not realizable, although they are subgroups of larger realizable non-abelian symmetry groups.

4.4 Abelian symmetries of the 4HDM potential

In the case of four Higgs doublets the representative maximal torus in $PSU(4)$ is $T = U(1)_1 \times U(1)_2 \times U(1)_3$, where

$$U(1)_1 = \alpha(-1, 1, 0, 0), \quad U(1)_2 = \beta(-2, 1, 1, 0), \quad U(1)_3 = \gamma\left(-\frac{3}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right). \quad (43)$$

The phase rotations of a generic bilinear combination of doublets under T is characterized by three integers p, q, r ,

$$(\phi_a^\dagger \phi_b) \rightarrow \exp[i(p_{ab}\alpha + q_{ab}\beta + r_{ab}\gamma)](\phi_a^\dagger \phi_b). \quad (44)$$

These coefficients can again be represented graphically as labels of edges of three simplices shown in Fig. 2.




Figure 2: Coefficients p and q as labels of triangles representing bilinears in 4HDM

With four doublets, one can construct (up to conjugation) 42 monomials transforming non-trivially under T . When we search for realizable finite groups, we pick up all possible

combinations of three distinct monomials v_1, v_2, v_3 , construct the 3×3 matrix $X(v_1 + v_2 + v_3)$, check that it is non-degenerate and then diagonalize it to obtain $\text{diag}(d_1, d_2, d_3)$. The symmetry group is then $\mathbb{Z}_{d_1} \times \mathbb{Z}_{d_2} \times \mathbb{Z}_{d_3}$. Although this brute force algorithm can be easily implemented in a machine code, we can in fact apply the results of the next Section to find the list of all realizable finite subgroups of the maximal torus in 4HDM:

$$\mathbb{Z}_k \text{ with } k = 2, \dots, 8; \quad \mathbb{Z}_2 \times \mathbb{Z}_k \text{ with } k = 2, 3, 4; \quad \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2. \quad (45)$$

In short, these are all finite abelian groups of order ≤ 8 . Perhaps, the most surprising of these groups is \mathbb{Z}_7 , because it does not appear as a realizable subgroup of any single $U(1)_i$ factor (no two labels on Fig. 2 sum up to 7). An example of terms that possess this symmetry group is

$$\lambda(\phi_1^\dagger \phi_3)(\phi_1^\dagger \phi_4) + \lambda'(\phi_2^\dagger \phi_1)(\phi_2^\dagger \phi_4) + \lambda''(\phi_3^\dagger \phi_2)(\phi_3^\dagger \phi_4), \quad (46)$$

and the \mathbb{Z}_7 group is generated by the transformation with phases

$$a = \frac{2\pi}{28}(-9, -1, 3, 7). \quad (47)$$

One can immediately check that a^7 lies in the center $Z(SU(4))$ and is, therefore, equivalent to the trivial transformation.

Realizable continuous abelian symmetry groups as well as abelian groups containing antiunitary transformations can also be found following the general strategy.

5 Abelian symmetries in general NHDM

The algorithm described above can be used to find all abelian groups realizable as the symmetry groups of the Higgs potential for any N . We do not yet have the full list of finite abelian groups for a generic N presented in a compact form, although we put forth a conjecture concerning this issue, see Conjectures 3 and 6 below. However, several statements can be proved about the order and possible structure of finite realizable subgroups of the maximal torus.

Throughout this Section we will often use $n := N - 1$. Also, whenever we mention in this Section a finite abelian group we actually imply a finite realizable subgroup of the maximal torus.

5.1 Upper bound on the order of finite abelian groups

It can be expected from the general construction that for any given N there exists an upper bound on the order of finite realizable subgroups of the maximal torus in NHDM.

Let us first prove that the order of any such group must be $\leq 4^{N-1}$. With the choice of the maximal torus (17), let us construct $n = N - 1$ bilinears $(\phi_1^\dagger \phi_{i+1})$, $i = 1, \dots, n$. The vectors of coefficients $a_i = (p_i, q_i, \dots, t_i)$ can be easily written:

$$\begin{aligned} a_1 &= (2, 3, 4, \dots, n, 1), \\ a_2 &= (1, 3, 4, \dots, n, 1), \\ a_3 &= (1, 2, 4, \dots, n, 1), \\ &\vdots \\ a_n &= (1, 2, 3, \dots, n-1, 1). \end{aligned} \quad (48)$$

One can use these vectors to construct the $n \times n$ matrix A :

$$A = \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}, \quad \det A = 1. \quad (49)$$

From the unit determinant we can also conclude that after diagonalization the matrix A becomes the unit matrix.

Now note that the vector of coefficients for $(\phi_i^\dagger \phi_j)$ with $i, j \neq 1$ can be represented as $a_j - a_i$ and it contains only 0's and 1's. Therefore, any monomial in the potential has the vector of coefficients in the form at most $a_j - a_i + a_l - a_k$ (it can contain fewer terms if ϕ_1 enters this monomial).

Let us now pick up n different monomials and construct the matrix X . Suppose it is non-degenerate in order for the symmetry group to be finite. The order of this symmetry group is then

$$\det X \equiv \det \begin{pmatrix} a_{j_1} - a_{i_1} + a_{l_1} - a_{k_1} \\ a_{j_2} - a_{i_2} + a_{l_2} - a_{k_2} \\ \vdots \\ a_{j_n} - a_{i_n} + a_{l_n} - a_{k_n} \end{pmatrix}. \quad (50)$$

Again, if ϕ_1 enters any of these monomials, its vector of coefficients will contain fewer a 's. The determinant is a polylinear combination of the rows: therefore it can be represented as a sum of four determinants

$$\det \begin{pmatrix} a_{j_1} \\ \vdots \end{pmatrix} - \det \begin{pmatrix} a_{i_1} \\ \vdots \end{pmatrix} + \det \begin{pmatrix} a_{l_1} \\ \vdots \end{pmatrix} - \det \begin{pmatrix} a_{k_1} \\ \vdots \end{pmatrix}. \quad (51)$$

One can then repeat this decomposition for the second row, etc, arriving at the end at the sum of 4^n determinants, each being equal to ± 1 . In the worst possible case all 1's sum up with the same signs, thus providing an upper bound on the order of the group:

$$|G| \leq 4^{N-1}. \quad (52)$$

This bound can be improved. Note first that the vector of coefficients for any monomial v_i can be expressed as $\sum_{j=1}^n c_{ij} a_j$, so that the X -matrix is

$$X_{ik} = c_{ij} A_{jk}. \quad (53)$$

The matrix c_{ij} contains only 0's, ± 1 or ± 2 . If ϕ_1 does not enter the monomial v_i , then the corresponding row c_{ij} can be, up to permutations and overall sign change, of the following type:

$$(1, 1, -1, -1, 0, \dots, 0), \quad (1, 1, -2, 0, \dots, 0), \quad \text{or} \quad (2, -2, 0, \dots, 0). \quad (54)$$

If a monomial v_i contains ϕ_1 , the row c_{ij} can be of the form

$$(1, 1, -1, 0, \dots, 0), \quad (2, -1, 0, \dots, 0), \quad (1, 1, 0, \dots, 0), \quad \text{or} \quad (2, 0, \dots, 0). \quad (55)$$

It follows from (53) that $\det X = \det c \cdot \det A = \det c$. But geometrically, the determinant of n vectors represents the volume of the n -dimensional parallelogram spanned by these vectors,

which is equal to the product of absolute values of these vectors and the sines of the angles. The absolute values of vectors c_{ij} for any fixed i is at most $\sqrt{8}$. This leads to the following upper bound:

$$|G| \leq 2^{\frac{3(N-1)}{2}}. \quad (56)$$

It is clear from the construction that this upper bound can be improved further. Here is our conjecture on the outcome of this improvement:

Conjecture 3. *The exact upper bound on the order of the realizable finite subgroup of maximal torus in NHDM is*

$$|G| \leq 2^{N-1}. \quad (57)$$

5.2 Cyclic groups and their products

Here we prove two propositions which show that a rather broad class of finite abelian groups are indeed realizable as the symmetry groups of the Higgs potential in the NHDM. First we show which cyclic groups are realizable and then consider direct products of cyclic groups.

Before we proceed to the main results, we need to prove that whenever we have the decomposition (53), it is indeed sufficient to diagonalize the matrix c_{ij} instead of full matrix X in order to reconstruct the finite symmetry group. Let us recall that the matrix X defines a system of linear equations on angles α_i :

$$X_{ij}\alpha_j = c_{ik}A_{kj}\alpha_j = 2\pi r_i. \quad (58)$$

This system remains the same upon

- simultaneous sign change of the k -th column in c and k -th row in A ;
- simultaneous exchange of two columns in c and two rows in A ;
- simultaneous summation of two columns in c and subtraction of two rows in A .
- sign changes of the k -th column in A and α_k ;
- exchange of columns i and j in A and exchange $\alpha_i \leftrightarrow \alpha_j$;
- summation of two columns in A and subtraction of two α 's;
- sign changes of the k -th row in c and of the integer parameters r_k ;
- exchange of two rows in c and of two r 's;
- summation of two rows in c and of two parameters r .

In short, it means that all allowed manipulations with integer matrices described earlier can be used for A and c .

Then we proceed in the following way. We first diagonalize A ; the matrix c becomes modified by a series of allowed transformations. But diagonal A is equal to the unit matrix. Therefore, we are left only with c in the system of equations. We then diagonalize c and construct the symmetry group from its diagonal values.

Now we move to two propositions.

Proposition 4. *The cyclic group \mathbb{Z}_p is realizable for any positive integer $p \leq 2^n$.*

Proof. We start with the following $n \times n$ matrix c_{ij} :

$$c_{2^n} = \begin{pmatrix} 2 & -1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 2 & -1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 2 & -1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 2 & -1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 2 \end{pmatrix}. \quad (59)$$

By straightforward manipulation with columns, starting from the last one, we arrive at

$$c_{2^n} = \begin{pmatrix} 0 & -1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & -1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & -1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 & -1 \\ 2^n & 2^{n-1} & 2^{n-2} & 2^{n-3} & \cdots & 2^2 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 2^n & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix}, \quad (60)$$

which produces the finite group \mathbb{Z}_{2^n} . Now, let us modify c_{2^n} by replacing the zero at the left lower corner by -1 . Then the same transformation leads us to the group $\mathbb{Z}_{2^{n-1}}$. If, instead, we replace any zero in the first column by -1 , $c_{k1} = -1$, then the diagonalization procedure leads us to the group \mathbb{Z}_p with $p = 2^n - 2^{n-k}$.

Now, generally, take any integer $0 \leq q < 2^n$ and write it in the binary form. This form uses at most n digits. Write this binary form as a vector with n components and subtract it from the first column of the matrix c_{2^n} . Then, after diagonalization, we obtain the symmetry group \mathbb{Z}_p with $p = 2^n - q$. This completes the proof. \square

Just to illustrate this construction, for $n = 5$ and $q = 23$ we write

$$q = 23_{10} = 10111_2 = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 1 \\ 1 \end{pmatrix}; \quad c_{32} - q \equiv \begin{pmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 2 & -1 & 0 & 0 \\ -1 & 0 & 2 & -1 & 0 \\ -1 & 0 & 0 & 2 & -1 \\ -1 & 0 & 0 & 0 & 2 \end{pmatrix}.$$

Repeating the above procedure we obtain at the lowest left corner $2^5 - 2^4 - 2^2 - 2^1 - 2^0 = 32 - 23 = 9$, which gives the \mathbb{Z}_9 group.

Now we show that not only cyclic groups but many of their products can be realized as symmetry groups of some potential.

Proposition 5. *Let $n = \sum_{i=1}^k n_i$ be a partitioning of n into a sum of non-negative integers n_i . Then, the finite group*

$$G = \mathbb{Z}_{p_1} \times \mathbb{Z}_{p_2} \times \cdots \times \mathbb{Z}_{p_k} \quad (61)$$

is realizable for any $0 < p_i < 2^{n_i}$.

Proof. Let us start again from the matrix (59). For any partitioning of n one can turn this matrix into a block-diagonal matrix by replacing some of the -1 's by zeros. The matrix is then represented by the smaller square blocks of size n_1, n_2, \dots, n_k . The i -th blocks has exactly the form of (59), with n replaced by n_i . Therefore, within this block one can encode any cyclic group \mathbb{Z}_{p_i} , with $0 < p_i \leq 2^{n_i}$. Since each block can be treated independently, we can encode any group of the form (61). \square

Let us note that this proposition does not yet exhaust all possible finite abelian groups with order $\leq 2^n$. For example, for $n = 5$, we can think of the group $\mathbb{Z}_5 \times \mathbb{Z}_5$ whose order is smaller than $2^5 = 32$. However, there exists no partitioning of 5 that would lead to this group by applying the proposition just proved. At this moment, we cannot prove whether such groups are also realizable. However, we conjecture that they are. Together with the earlier conjecture on the upper bound on the order of the finite group, we can now propose the following conjecture:

Conjecture 6. *Any finite abelian group with order $\leq 2^{N-1}$ is realizable in NHDM.*

If proven true, this conjecture will give the complete classification of realizable subgroups of the maximal torus in NHDM.

6 Conclusions

Models with several Higgs doublets have become a popular framework to introduce physics beyond the Standard Model. When building the scalar sector of an N -doublet model, one usually imposes various discrete symmetries, and many specific symmetry groups have been used. However, the complete lists of groups that can be implemented for a given N are unknown for $N > 2$.

In this paper we made a step towards classification of possible symmetries of the scalar sector of the NHDM. Namely, we studied which abelian groups can be realized as symmetry groups of the NHDM potential. We proved that they can be either subgroups of the maximal torus, or the image under the canonical homomorphism $SU(N) \rightarrow PSU(N)$ of a finite nilpotent group of class 2. For the subgroups of the maximal torus, we developed an algorithmic strategy that gives full list of possible realizable abelian symmetries for any given N . We illustrated how the strategy works with two small- N examples. For 3HDM we gave the full list of realizable abelian symmetries, including the ones with generalized CP -transformations, while for the 4HDM we just listed all finite realizable subgroups of the maximal torus. We also proved several statements concerning the order and structure of finite groups which apply for arbitrary N . Finally, we conjectured that, in the NHDM, any finite abelian group with order $\leq 2^{N-1}$ is realizable.

It is worth stressing that the Higgs potential taken in the usual quadratic plus quartic form plays a major factor in restricting possible symmetry groups from *all* subgroups of $SU(N)$. If the potential could contain higher-order terms, the list of realizable group would be larger. It means, in particular, that the recent impressive works on the beyond-SM applications of the finite subgroups of $SU(3)$, [21], cannot be directly applied to the classification of the Higgs symmetries in 3HDM because no criterion was given which of the groups are realizable and which are not.

An obvious direction of future research is to understand phenomenological consequences of the symmetries found. This includes, in particular, a study of how the symmetries of the potential are broken, and what is the effect of these symmetries on the physical Higgs boson spectrum.

Acknowledgements

We are grateful to Pedro Ferreira and João Silva for valuable comments and suggestions. This work was supported by the Belgian Fund F.R.S.-FNRS via the contract of Chargé de recherches, and in part by grants RFBR No.11-02-00242-a and No.10-01-0391, and NSH-3810.2010.2.

A Maximal abelian subgroups in $PSU(N)$

Let $V = \mathbb{C}^N$ be a unitary space of dimension N equipped with unitary form (u, v) . If L is a subspace of V , then L^\dagger denotes the orthogonal complements to L in V . Now $SU(N)$ is the group of all transformations of determinant 1 preserving the form, i.e.,

$$SU(N) = \{x \mid \det x = 1 \text{ and } x^\dagger \cdot x = 1\}.$$

We denote the group of all transformations preserving the form by $GU(N)$. Clearly $SU(N)$ is a normal subgroup of $GU(N)$ ($SU(N) \trianglelefteq GU(N)$).

In the following lemma we collect known facts and we provide the proofs for the reader's convenience.

Lemma 7. *Let L be a subspace of V and x be a linear transformation of V . Then the following statements hold.*

- (1) *If L is x -invariant, then L^\dagger is x^\dagger -invariant.*
- (2) *If $x \in GU(N)$, then $x(v) = \lambda v$ implies $x^\dagger(v) = \bar{\lambda}v$.*
- (3) *If $x \in GU(N)$, then there exists an orthonormal basis consisting of eigenvectors of x . Moreover, if $\text{Spec}(x) = \{\lambda_1, \dots, \lambda_k\}$ is the set of all eigenvalues of x and $V_\lambda = \{v \in V \mid x(v) = \lambda v\}$, then there exists the orthogonal decomposition $V = V_{\lambda_1} \oplus^\dagger \dots \oplus^\dagger V_{\lambda_k}$.*
- (4) *If $x, y \in GU(N)$ commute, λ is an eigenvalue of x , then V_λ is y -invariant.*
- (5) *If $A \leq GU(N)$ is abelian, then A is conjugate to a subgroup of $D(N)$ in $GU(N)$, where $D(N)$ is a group of all diagonal matrices in $GU(N)$. In particular, if $A \leq SU(N)$ is abelian, then A is conjugate to a subgroup of $D(N) \cap SU(N)$ in $SU(N)$.*
- (6) *Suppose H is the stabilizer in $SU(N)$ of L and L^\dagger , i.e., $H = (GU(L) \times GU(L^\dagger)) \cap SU(N)$. Assume that $L, L^\dagger \neq 0$. Let $\bar{\cdot} : SU(N) \rightarrow PSU(N)$ be the reduction modulo scalars. Then the natural projection $\pi : \bar{H} \rightarrow GU(L)$ is surjective. Moreover, if T is a maximal torus of H , then the projection of T into $GU(L)$ is a maximal torus of $GU(L)$.*

Proof. By definition of \mathbf{x}^\dagger , for every $u, v \in V$ we have $(\mathbf{x}(u), v) = (u, \mathbf{x}^\dagger(v))$, whence (1).

(2) Assume that $\mathbf{x}(v) = \lambda v$. Then

$$\begin{aligned} 0 &= (\mathbf{x}(v) - \lambda v, \mathbf{x}(v) - \lambda v) = ((\mathbf{x} - \lambda 1)(v), (\mathbf{x} - \lambda 1)(v)) = (v, (\mathbf{x}^\dagger - \bar{\lambda} 1)(\mathbf{x} - \lambda 1)(v)) = \\ &= (v, (\mathbf{x} - \lambda 1)(\mathbf{x}^\dagger - \bar{\lambda} 1)(v)) = ((\mathbf{x}^\dagger - \bar{\lambda} 1)(v), (\mathbf{x}^\dagger - \bar{\lambda} 1)(v)), \end{aligned}$$

whence $\mathbf{x}^\dagger(v) - \bar{\lambda}v = 0$.

(3) Since \mathbb{C} is algebraically closed, there exists an eigenvalue λ of \mathbf{x} . Therefore there exists an eigenvector v such that $\mathbf{x}(v) = \lambda v$. Clearly we may assume that $\|v\| = 1$. Consider the line $L(v)$ spanned by v . In view of item (2) of the lemma, $L(v)$ is \mathbf{x}^\dagger -invariant, hence item (1) of the lemma implies that $L(v)^\dagger$ is \mathbf{x} -invariant. Therefore we obtain a decomposition $V = L(v) \oplus L(v)^\dagger$ of V into an orthogonal sum of \mathbf{x} -invariant subspaces. Induction on $\dim(V)$ completes the proof of this item.

(4) For every $v \in V_\lambda$ we have $\mathbf{x}(\mathbf{y}(v)) = \mathbf{y}(\mathbf{x}(v)) = \lambda \mathbf{y}(v)$, i.e., $\mathbf{y}(v) \in V_\lambda$. Therefore V_λ is a \mathbf{y} -invariant subspace.

(5) Assume that $A \leq GU(N)$ is abelian. Clearly we may assume that A is nonscalar. Choose $x \in A \setminus Z(GU(N))$. Given $\lambda \in \mathbb{C}$ consider $V_\lambda = \{v \in V \mid \mathbf{x}(v) = \lambda v\}$. Denote by $Spec(\mathbf{x})$ the set of all eigenvalues of \mathbf{x} . Since \mathbb{C} is algebraically closed, we obtain that $Spec(\mathbf{x})$ is nonempty. Moreover, since $\mathbf{x} \in A \setminus Z(GU(N))$, it follows that $Spec(\mathbf{x})$ contains at least two elements. In view of the proof of item (3) we obtain the orthogonal decomposition

$$V = \bigoplus_{\lambda \in Spec(\mathbf{x})} V_\lambda.$$

Choose $y \in A$. Then, for every $\lambda \in Spec(\mathbf{x})$ the subspace V_λ is y -invariant, hence V_λ is A -invariant. Induction on $\dim(V)$ completes the proof of this item.

(6) A matrix from $(GU(L) \times GU(L^\dagger)) \cap SU(N)$ has the following form:

$$\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$$

and $|A| \cdot |B| = 1$. Assume that $A \in GU(L)$ and $|A| = \mu$. Then $|\mu| = 1$, i.e. $\mu \cdot \mu^* = 1$, so if $D = \text{diag}(\mu^*, 1, \dots, 1)$, then

$$\mathbf{x} = \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} \in H.$$

Moreover $\pi(\mathbf{x}) = A$, whence (6). □

Consider the canonical epimorphism

$$\overline{\quad} : SU(N) \rightarrow SU(N)/Z(SU(N)) = \overline{SU(N)} \simeq PSU(N), \quad (62)$$

where $Z(SU(N)) \simeq \mathbb{Z}_N$ is the center of $SU(N)$. Given subgroup \overline{A} of $PSU(N)$ we denote by A the complete preimage of \overline{A} in $SU(N)$. The following statement is evident.

Proposition 8. *Let \overline{A} be an abelian subgroup of $PSU(N)$. Then A is either an abelian group or a nilpotent group of class 2 and $Z(SU(N)) \leq Z(A)$.*

Proof. Since $Z(SU(N)) \leq Z(A)$ for every $A \leq SU(N)$, the statement is evident. □

If \overline{A} is a maximal abelian subgroup of $PSU(N)$ such that A is abelian, then A must be a maximal abelian subgroup of $SU(N)$, and item (5) of Lemma 7 implies that A is a maximal torus.

Now we find the structure of A in case, when A is nonabelian. We assume that \overline{A} is a maximal abelian subgroup of $PSU(N)$.

Lemma 9. *If A is nonabelian, then $Z(A) = Z(SU(N))$.*

Proof. Inclusion $Z(SU(N)) \leq Z(A)$ is evident. Assume that there exists $\mathbf{x} \in Z(A) \setminus Z(SU(N))$. Then \mathbf{x} is nonscalar, so $\text{Spec}(\mathbf{x}) = \{\lambda_1, \dots, \lambda_k\}$ and $k \geq 2$. In view of Lemma 7 items (3) and (4), $A \leq C_{SU(N)}(\mathbf{x}) = GU(N_1) \times \dots \times GU(N_k)$, where $N_i = \dim(V_{\lambda_i})$. Now Lemma 7 items (5) and (6) implies that the projection of \overline{A} into $GU(N_i)$ is contained in a maximal torus of $GU(N_i)$. Since \overline{A} is a maximal abelian subgroup of $PSU(N)$, we obtain that the projection of A coincides with a maximal torus of $GU(N_i)$ for every i . Hence \overline{A} is a maximal torus of $PSU(N)$, and so A is a maximal torus of $SU(N)$. In particular, A is abelian, a contradiction. \square

Since $Z(A) = Z(SU(N)) \simeq \mathbb{Z}_N$, we obtain that the exponent of $Z(A)$ equals N . In view of [29, 5.2.22(i)], the exponent of A divides N^2 , in particular, $\pi(A) = \pi(N)$, where $\pi(A)$ denotes the set of prime divisors of $|A|$ and $\pi(N)$ denotes the set of prime divisors of N .

Lemma 10. *A is finite.*

Proof. Let H be a maximal normal abelian subgroup of A . Then $C_A(H) = H$ (see [29, 5.2.3]) and H is contained in a maximal torus of $SU(N)$. In particular, H is generated by a finite number of elements. Since $\pi(H) \subseteq \pi(A) = \pi(N)$ is finite and the exponent of H divides N^2 , we obtain that H is finite. Since $H = C_A(H)$, it follows that A/H is isomorphic to a subgroup of $\text{Aut}(H)$, in particular A/H is finite. Therefore A is finite. \square

All the information about abelian subgroups of $PSU(N)$ obtained above was assembled in Theorem 1 in the main text.

We finish by giving an example of subgroups satisfying statement (2) of Theorem 1. Let ω be the primitive N -th root of 1 in \mathbb{C} . Consider $\mathbf{x} = \text{diag}(1, \omega, \omega^2, \dots, \omega^{N-1})$, $\mathbf{y} = e_{1,2} + e_{2,3} + \dots + e_{N-1,N} + e_{N,1}$, where $e_{i,j}$ is the matrix with 1 on the (i, j) -place and 0 on the remaining places. Then $M = \langle \mathbf{x}, \mathbf{y} \rangle$ is a nilpotent subgroup of class 2 of $SU(N)$ for N odd, $Z(N) = Z(SU(N))$ and the exponent of M equals N .

B Abelian symmetries in 3HDM with antiunitary transformations

Here we explicitly describe embedding of each of the unitary abelian groups (32) into a larger abelian group that contains antiunitary transformations. Throughout this section we will use the following notation: if a cyclic group \mathbb{Z}_q is generated by an antiunitary transformation, we indicate it by an asterisk: \mathbb{Z}_q^* .

B.1 Embedding $U(1)_1 \times U(1)_2$

We proved in the general case that the maximal torus cannot be embedded in a larger abelian group. For illustration, let us repeat the argument in the specific case of 3HDM. At the first step of the strategy we search for a matrix $b \in PSU(3)$ that satisfies $aba = b$ for all $a \in T$. Since a is diagonal, each element $(aba)_{ij}$ is equal to b_{ij} up to a phase rotation ψ_{ij} :

$$\psi_{ij} = \begin{pmatrix} -2\alpha - 4\beta/3 & -\beta/3 & -\alpha - \beta/3 \\ -\beta/3 & 2\alpha + 2\beta/3 & \alpha + 2\beta/3 \\ -\alpha - \beta/3 & \alpha + 2\beta/3 & 2\beta/3 \end{pmatrix}. \quad (63)$$

In order for an element b_{ij} to be non-zero, the corresponding phase rotation must be zero or multiple of 2π , modulo to $2\pi k/3$ along the diagonal. It is impossible to find such b for arbitrary α and β . Therefore, $T = U(1)_1 \times U(1)_2$ cannot be embedded into a larger abelian group.

B.2 Embedding $U(1)$

There are two distinct sorts of $U(1)$ groups inside T : $U(1)_1$ -type and $U(1)_2$ -type. Consider first $U(1)_1$. The phase rotations for the group $U(1)_1$ ($\beta = 0$, arbitrary α) allow b to have non-zero entries b_{12} , b_{21} and b_{33} only. We are free to choose $b_{12} = b_{21} = 1$, $b_{33} = 1$ (the fact that $\det b = -1$ instead of 1 is inessential; we can always multiply all the transformations by the overall -1 factor). Then, the transformation J' which commutes with any $a \in U(1)_1$ is

$$J' = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot J, \quad aJ' = J'a \quad \forall a \in U(1)_1. \quad (64)$$

Next, we search for transformations $x \in PSU(3)$ such that xJ' also commutes with any element $a \in U(1)_1$; such transformations x form the centralizer of $U(1)_1$. From the equation $axa^{-1} = x$, analyzed with the same technique of phase rotations, we find that x must be diagonal. By checking the production of x_1J' and x_2J' stays inside $U(1)_1$, we can conclude that x must belong to one of the $U(1)_1$ -type groups X_ξ :

$$x = \begin{pmatrix} e^{-i\gamma} & 0 & 0 \\ 0 & e^{i\gamma} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{i\xi} & 0 & 0 \\ 0 & e^{i\xi} & 0 \\ 0 & 0 & e^{-2i\xi} \end{pmatrix}. \quad (65)$$

Here $\xi \in [0, 2\pi/3)$ is an arbitrary but fixed parameter specifying which X_ξ group we take, while γ is the running angle parametrizing the elements of this group. At this stage, any choice of ξ is acceptable. Thus, we embedded $U(1)_1$ into an abelian group generated by $\langle U(1)_1, J''_\xi \rangle \simeq U(1) \times \mathbb{Z}_2^*$, where

$$J''_\xi = \begin{pmatrix} e^{i\xi} & 0 & 0 \\ 0 & e^{i\xi} & 0 \\ 0 & 0 & e^{-2i\xi} \end{pmatrix} \cdot J' = \begin{pmatrix} 0 & e^{i\xi} & 0 \\ e^{i\xi} & 0 & 0 \\ 0 & 0 & e^{-2i\xi} \end{pmatrix} \cdot J. \quad (66)$$

Let us now see whether this group is realizable and which ξ must be chosen. The only $U(1)_1$ symmetric terms in the potential are

$$\lambda(\phi_1^\dagger \phi_3)(\phi_2^\dagger \phi_3) + \lambda^*(\phi_3^\dagger \phi_1)(\phi_3^\dagger \phi_2). \quad (67)$$

This sum of two terms is indeed invariant under J_ξ'' provided $\xi = \psi_\lambda/3$, where ψ_λ is the phase of λ . This prescription uniquely specifies which group $X_\xi \subset PSU(3)$. Besides, in order to guarantee that the T -symmetric terms (27) are invariant under J'' , we must set

$$m_{11}^2 = m_{22}^2, \quad \lambda_{11} = \lambda_{22}, \quad \lambda_{13} = \lambda_{23}, \quad \lambda'_{13} = \lambda'_{23}. \quad (68)$$

However upon this reduction of free parameters we observe that the resulting potential acquires an additional *unitary* symmetry: the exchange of the field two doublets $\phi_1 \leftrightarrow \phi_2$. This transformation does not commute with $U(1)_1$. Therefore, the true symmetry group of such potential is non-abelian, though it contains the desired abelian subgroup $U(1) \times \mathbb{Z}_2^*$. According to our definition, we conclude that $U(1) \times \mathbb{Z}_2^*$ is not realizable in 3HDM.

Consider now $U(1)_2$: $\alpha = 0$, arbitrary β . It follows from (63) that no non-trivial matrix $b \in PSU(3)$ can satisfy $aba = b$. Therefore, $U(1)_2$ cannot be embedded into a larger abelian group with antiunitary transformations. Note also that selecting various diagonal $U(1)$ subgroups of T , for example by setting $\alpha + \beta/3 = 0$, will result just in another version of $U(1)_1$. Finally, we note that the unitary $U(1) \times \mathbb{Z}_2$ emerges only when the continuous group is of the $U(1)_2$ -type. Therefore, $U(1) \times \mathbb{Z}_2$ cannot be embedded into a larger realizable abelian group as well.

The overall result of the last two subsections is that in 3HDM all realizable continuous abelian groups are necessarily unitary and cannot contain antiunitary transformations.

B.3 Embedding \mathbb{Z}_4

The \mathbb{Z}_4 symmetry group with generator a arises as a subgroup of $U(1)_1$. Therefore, b and J' can be chosen as before, (64), but the new conditions for x should now be checked. This transformation can still be represented as in (65), but the angle γ can take discrete values:

$$\gamma = \frac{\pi}{2}k \quad \text{or} \quad \gamma = \frac{\pi}{4} + \frac{\pi}{2}k, \quad k \in \mathbb{Z}. \quad (69)$$

Correspondingly, two possibilities for embedding the \mathbb{Z}_4 group arise: $\mathbb{Z}_4 \times \mathbb{Z}_2^*$ and \mathbb{Z}_8^* .

Consider first the $\mathbb{Z}_4 \times \mathbb{Z}_2^*$ embedding. The \mathbb{Z}_2^* subgroup is generated J'' in (66), where, as before, $\xi = \psi_\lambda/3$. As before, the potential satisfies (68), includes (67), but in addition it now contains

$$\lambda'(\phi_1^\dagger\phi_2)^2 + \lambda^*(\phi_2^\dagger\phi_1)^2, \quad (70)$$

Note that these terms are invariant under J'' , and therefore, under the full $\mathbb{Z}_4 \times \mathbb{Z}_2^*$. which is also J'' -symmetric when $\xi_1 = \xi_2$. However, just as in the previous case, restrictions (68) result in a potential which is symmetric under another \mathbb{Z}_2 group generated by the unitary transformation

$$d = \begin{pmatrix} 0 & e^{i\delta} & 0 \\ e^{-i\delta} & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{with} \quad \delta = \frac{\psi_{\lambda'}}{2}, \quad (71)$$

where $\psi_{\lambda'}$ is the phase of λ' in (70). Since d does not commute with phase rotations, therefore, the resulting symmetry group of the potential is non-abelian. Hence, the $\mathbb{Z}_4 \times \mathbb{Z}_2^*$ symmetry group is not realizable.

Consider now the \mathbb{Z}_8^* group generated by

$$J_\xi''' = \begin{pmatrix} 0 & e^{i\xi - i\pi/4} & 0 \\ e^{i\xi + i\pi/4} & 0 & 0 \\ 0 & 0 & e^{-2i\xi} \end{pmatrix} \cdot J \quad (72)$$

with the property $(J''')^2 = a$, the generator of the \mathbb{Z}_4 . Note that upon action of J''' the term $(\phi_1^\dagger \phi_2)^2$ just changes its sign; therefore, such term cannot appear in the potential. But in this case we are left with only one type of \mathbb{Z}_4 -symmetric terms. Then, according to the general discussion, $\text{rank}X(V) = 1$ and, therefore, the potential becomes symmetric under a continuous symmetry group. Thus, \mathbb{Z}_8^* is not realizable as a symmetry group of a 3HDM potential.

B.4 Embedding \mathbb{Z}_3

The \mathbb{Z}_3 subgroup of the maximal torus T is generated by the transformation a which rotates the phases of the doublets by $(-2\pi/3, 2\pi/3, 0)$. This group also arises from the $U(1)_1$ -type group, therefore the representation for J' in (64) and x in (65) are still valid. Closing the group under product requires that γ in (65) is a multiple of $\pi/3$. If we introduce

$$J''_\xi = \begin{pmatrix} 0 & e^{i\xi - i\pi/3} & 0 \\ e^{i\xi + i\pi/3} & 0 & 0 \\ 0 & 0 & e^{-2i\xi} \end{pmatrix} \cdot J. \quad (73)$$

Since $(J'')^2 = a$, this transformation generates the group \mathbb{Z}_6^* .

The \mathbb{Z}_3 -symmetric terms in the potential are

$$\lambda_1(\phi_1^\dagger \phi_2)(\phi_1^\dagger \phi_3) + \lambda_2(\phi_2^\dagger \phi_3)(\phi_2^\dagger \phi_1) + \lambda_3(\phi_3^\dagger \phi_1)(\phi_3^\dagger \phi_2) + h.c. \quad (74)$$

with complex λ_i . These terms become symmetric under \mathbb{Z}_6^* , if the following conditions are satisfied:

$$|\lambda_1| = |\lambda_2| \quad \text{and} \quad \psi_1 + \psi_2 + \psi_3 = \pi, \quad (75)$$

where ψ_i are phases of λ_i . Indeed, one can check that in this case the sum (74) is left invariant under J''_ξ with $\xi = -\psi_3/3$. It is, therefore, possible to have a \mathbb{Z}_6^* -symmetric potential by requiring (75) and applying the usual restrictions (68).

However, just as in the previous cases, the resulting potential becomes symmetric under a larger unitary symmetry group, and therefore \mathbb{Z}_6^* is not realizable as well. The transformation that leaves the potential symmetric is of type (71) with $\delta = (\psi_1 - \psi_2)/3$.

B.5 Embedding $\mathbb{Z}_2 \times \mathbb{Z}_2$

The $\mathbb{Z}_2 \times \mathbb{Z}_2 \subset T$ symmetry can be realized as a group of simultaneous sign flips of any pair of doublets (or alternatively of independent sign flips of any of the three doublets). If the simultaneous sign flip of doublets i and j are denoted as R_{ij} , then the group is $1, R_{12}, R_{13}, R_{23}$, with $R_{12}R_{13} = R_{23}$ and any $(R_{ij})^2 = 1$. This group can be embedded into $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2^*$, where \mathbb{Z}_2^* is generated by

$$J'_{\xi_1, \xi_2} = \begin{pmatrix} e^{i\xi_1} & 0 & 0 \\ 0 & e^{i\xi_2} & 0 \\ 0 & 0 & e^{-i\xi_1 - i\xi_2} \end{pmatrix} \cdot J. \quad (76)$$

At this moment, any pair ξ_1, ξ_2 can be used to construct the $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2^*$ group. Note also that this group can be also written as $\mathbb{Z}_2 \times \mathbb{Z}_2^* \times \mathbb{Z}_2^*$ or $\mathbb{Z}_2^* \times \mathbb{Z}_2^* \times \mathbb{Z}_2^*$ by redefining the generators.

The $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetric potential contains the following terms

$$\tilde{\lambda}_{12}(\phi_1^\dagger \phi_2)^2 + \tilde{\lambda}_{23}(\phi_2^\dagger \phi_3)^2 + \tilde{\lambda}_{31}(\phi_3^\dagger \phi_1)^2 + h.c. \quad (77)$$

with complex $\tilde{\lambda}_{ij}$. If there phases are denoted as ψ_{ij} , then the condition when (77) are symmetric under some \mathbb{Z}_2^* with an appropriate choice of ξ_1, ξ_2 is

$$\psi_{12} + \psi_{23} + \psi_{31} = 0, \quad (78)$$

or alternatively, when the product $\tilde{\lambda}_{12}\tilde{\lambda}_{23}\tilde{\lambda}_{31}$ is real.

Note that in contrast to the previous cases, the full T -symmetric potential (27) is invariant under $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2^*$. This guarantees that no other unitary symmetry arises in this case, and therefore this symmetry group is realizable.

B.6 Embedding \mathbb{Z}_2

The unitary abelian symmetry group \mathbb{Z}_2 can be implemented in a variety of ways, all of which are equivalent. One can take, for example, the \mathbb{Z}_2 group generated by sign flips of the first and second doublets, R_{12} . Again, the usual CP -transformation J commutes with R_{12} . An analysis similar to the previous case shows that the $\mathbb{Z}_2 \times \mathbb{Z}_2^*$ group generated by R_{12} and J is realizable.

There is also a possibility to embed the \mathbb{Z}_2 group into \mathbb{Z}_4^* . Consider the following transformation

$$J'_\xi = \begin{pmatrix} 0 & e^{i\xi} & 0 \\ -e^{i\xi} & 0 & 0 \\ 0 & 0 & e^{-2i\xi} \end{pmatrix} \cdot J, \quad (J')^2 = R_{12}, \quad (79)$$

where ξ is an arbitrary but fixed parameter. The potential with the usual restrictions (68) and with extra terms

$$\lambda_5(\phi_1^\dagger\phi_2)^2 + \lambda_6(\phi_1^\dagger\phi_2) \left[(\phi_1^\dagger\phi_1) - (\phi_2^\dagger\phi_2) \right] + \lambda_7(\phi_1^\dagger\phi_3)(\phi_2^\dagger\phi_3) + \lambda_8(\phi_1^\dagger\phi_3)^2 + \lambda_9(\phi_2^\dagger\phi_3)^2 + h.c. \quad (80)$$

satisfying conditions $|\lambda_8| = |\lambda_9|$ and $\psi_8 + \psi_9 = 2\psi_7 + \pi$ is invariant under J'_ξ with $6\xi = \psi_8 + \psi_9$. Adjusting λ_5 and λ_6 , one can guarantee that no other unitary symmetry appears as a result of (68). Therefore, this \mathbb{Z}_4^* group is realizable.

References

- [1] E. Accomando *et al.*, “Workshop on CP studies and non-standard Higgs physics,” arXiv:hep-ph/0608079.
- [2] S. F. King, C. Luhn, JHEP **0910**, 093 (2009); Q. -H. Cao, S. Khalil, E. Ma, H. Okada, Phys. Rev. Lett. **106**, 131801 (2011).
- [3] T. D. Lee, Phys. Rev. D **8**, 1226 (1973).
- [4] J.F. Gunion, H.E. Haber, G. Kane, S. Dawson, *The Higgs Hunter’s Guide* (Addison-Wesley, Reading, 1990).
- [5] G. C. Branco et al, arXiv:1106.0034 [hep-ph].
- [6] F. J. Botella and J. P. Silva, Phys. Rev. D **51**, 3870 (1995); G. C. Branco, L. Lavoura and J .P. Silva, “CP-violation”, Oxford University Press, Oxford, England (1999).

- [7] S. Davidson and H. E. Haber, Phys. Rev. D **72**, 035004 (2005) [Erratum-ibid. D **72**, 099902 (2005)]; H. E. Haber and D. O’Neil, Phys. Rev. D **74**, 015018 (2006) [Erratum-ibid. D **74**, 059905 (2006)].
- [8] J. F. Gunion and H. E. Haber, Phys. Rev. D **72**, 095002 (2005).
- [9] D. O’Neil, “Phenomenology of the Basis-Independent CP-Violating Two-Higgs Doublet Model”, Ph.D. thesis, University of California Santa Cruz (2009), arXiv:0908.1363 [hep-ph].
- [10] G. Sartori and G. Valente, arXiv:hep-ph/0304026.
- [11] F. Nagel, “New aspects of gauge-boson couplings and the Higgs sector”, Ph.D. thesis, University Heidelberg (2004), [<http://www.ub.uni-heidelberg.de/archiv/4803>].
- [12] M. Maniatis, A. von Manteuffel, O. Nachtmann and F. Nagel, Eur. Phys. J. C **48**, 805 (2006); M. Maniatis, A. von Manteuffel and O. Nachtmann, Eur. Phys. J. C **57**, 719 (2008).
- [13] C. C. Nishi, Phys. Rev. D **74**, 036003 (2006) [Erratum-ibid.D **76**, 119901 (2007)].
- [14] I. P. Ivanov, Phys. Lett. B **632**, 360 (2006); Phys. Rev. D **75**, 035001 (2007) [Erratum-ibid. D **76**, 039902 (2007)]; Phys. Rev. D **77**, 015017 (2008).
- [15] P. M. Ferreira, H. E. Haber, M. Maniatis, O. Nachtmann, J. P. Silva, Int. J. Mod. Phys. **A26**, 769-808 (2011). [arXiv:1010.0935 [hep-ph]].
- [16] A. Pilaftsis, arXiv:1109.3787 [hep-ph].
- [17] R. Erdem, Phys. Lett. B **355**, 222 (1995).
- [18] A. Barroso, P. M. Ferreira, R. Santos and J. P. Silva, Phys. Rev. D **74**, 085016 (2006).
- [19] C. C. Nishi, Phys. Rev. D **76**, 055013 (2007).
- [20] P. M. Ferreira and J. P. Silva, Phys. Rev. D **78**, 116007 (2008).
- [21] W. Grimus and P. O. Ludl, J. Phys. A **43**, 445209 (2010);
W. Grimus and P. O. Ludl, arXiv:1110.6376 [hep-ph].
- [22] K. Olausen, P. Osland and M. A. Solberg, JHEP **1107**, 020 (2011).
- [23] I. P. Ivanov, C. C. Nishi, Phys. Rev. D **82**, 015014 (2010).
- [24] I. P. Ivanov, JHEP **1007**, 020 (2010).
- [25] A. C. B. Machado, J. C. Montero, V. Pleitez, Phys. Lett. B **697**, 318-322 (2011)
- [26] T. Fukuyama, H. Sugiyama, K. Tsumura, Phys. Rev. D **82**, 036004 (2010)
- [27] S. Adler, Phys. Rev. D **60**,015002 (1999).
- [28] I. P. Ivanov, V. Keus, Phys. Lett. **B695**, 459-462 (2011).
- [29] D. J. S. Robinson, “A Course in the Theory of Groups”, Springer, 1996, Second Edition.