

High-precision CoRoT space photometry and fundamental parameter determination of the B2.5V star HD 48977 (Research Note)

A. Thoul^{1,6,*}, P. Degroote^{2,6,**}, C. Catala³, C. Aerts^{2,4,6}, T. Morel¹, M. Briquet^{1,2,***}, M. Hillen², G. Raskin², H. Van Winckel², M. Auvergne³, A. Baglin³, F. Baudin⁵, E. Michel³

¹ Institut d'Astrophysique et de Géophysique, Université de Liège, 17 allée du 6 août, B4000, Liège, Belgium
e-mail: anne.thoul@ulg.ac.be

² Instituut voor Sterrenkunde, K.U.Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium

³ LESIA, Observatoire de Paris, CNRS UMR 8109, Université Pierre et Marie Curie, Université Denis Diderot, 5 place J. Janssen, 92105 Meudon, France

⁴ Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands

⁵ Institut d'Astrophysique Spatiale, CNRS/Université Paris XI UMR 8617, F-091405 Orsay, France

⁶ Kavli Institute for Theoretical Physics, Kohn Hall, University of California, Santa Barbara CA 93106-4030, USA

Received ??, 2011; accepted ??, 2011

ABSTRACT

We present the CoRoT light curve of the bright B2.5V star HD 48977 observed during a short run of the mission in 2008, as well as a high-resolution spectrum gathered with the HERMES spectrograph at the Mercator telescope. We use several time series analysis tools to explore the nature of the variations present in the light curve. We perform a detailed analysis of the spectrum of the star to determine its fundamental parameters and its element abundances. We find a large number of high-order g-modes, and one rotationally induced frequency. We find stable low-amplitude frequencies in the p-mode regime as well. We conclude that HD 48977 is a new Slowly Pulsating B star with fundamental parameters found to be $T_{\text{eff}} = 20\,000 \pm 1000\text{K}$ and $\log g = 4.2 \pm 0.1$. The element abundances are similar to those found for other B stars in the solar neighbourhood. HD 48977 was observed during a short run of the CoRoT satellite implying that the frequency precision is insufficient to perform asteroseismic modelling of the star. Nevertheless, we show that a longer time series of this star would be promising for such modelling. Our present study contributes to a detailed mapping of the instability strips of B stars in view of the dominance of g-mode pulsations in the star, several of which occur in the gravito-inertial regime.

Key words. Asteroseismology – Stars: Individual: HD 48977, 16Mon – Stars: abundances – Stars: fundamental parameters – Stars: oscillations – Techniques: photometric

1. Introduction

Asteroseismology, the study of stellar oscillations in order to infer information about the stellar structure, is a very powerful tool to improve our knowledge of the physics of stellar interiors (Aerts et al. 2010). Helioseismology, the seismology of the Sun, has helped tremendously in improving our knowledge of the Sun's structure and interior physics (Christensen-Dalsgaard 2002). Asteroseismology is a younger but very dynamic field in astronomy, and several recent space missions, in particular MOST, CoRoT, and *Kepler*, have provided high-precision light curves for several classes of variable stars, including main sequence B stars (e.g., Aerts et al. (2006a,b); Saio et al. (2007); Cameron et al. (2008); Huat et al. (2009); Neiner et al. (2009); Diago et al. (2009); Gutiérrez-Soto et al. (2009); Degroote et al. (2009, 2010, 2011); Pápics et al. (2011); Balona et al. (2011)). Despite these high-precision data from space, in-depth seismic

modelling of most of these pulsating B stars must await (partial) identification of the wavenumbers of the dominant observed pulsation frequencies. This was only achieved so far for the Slowly Pulsating B (hereafter SPB) star HD 50230, which led to the conclusion that this star has an inhomogeneously mixed region surrounding its core (Degroote et al. 2010), and for the β Cep star HD 180642 (V1449 Aql) which has an unusually high-amplitude dominant radial mode that induces several other lower-amplitude non-radial modes of the star through resonant mode coupling (Briquet et al. 2009; Aerts et al. 2011).

In the last decade, several asteroseismic studies of B stars based on large ground-based campaigns have been performed as well (Aerts et al. 2003a,b, 2004b; Handler et al. 2004; Aerts et al. 2004a; De Ridder et al. 2004; Jerzykiewicz et al. 2005; Dziembowski & Pamyatnykh 2008; Handler et al. 2005; Briquet et al. 2005; Handler et al. 2006; Desmet et al. 2009). These observations are very extensive and complicated, often involving multi-site multi-technique campaigns. Pulsating B stars typically pulsate with low-order p and g modes and some also experience high-order g modes, making it necessary to observe them for a very long time in order to resolve all those modes. In addition, daily aliases are an observational difficulty since these stars often have (beat) periods of the or-

* Chercheur Qualifié, Fond de la Recherche Scientifique (F.R.S.-FNRS)

** Postdoctoral Fellow, Fund for Scientific Research of Flanders (FWO)

*** Postdoctoral Fellow, Fond de la Recherche Scientifique (F.R.S.-FNRS)

der of a day. Nevertheless, ground-based asteroseismic studies of B stars have proven to be very successful and have led to important new insights in their interior physics. For example, it was possible to put limits on the value of the overshooting parameter of the core (Ausseloos et al. 2004; Dupret et al. 2004; Briquet et al. 2007; Desmet et al. 2009). From partially resolved multiplets due to rotational splitting, it was also possible to show that some of those stars have non-rigid rotation, in the sense that their core regions rotate faster than their envelope (Pamyatnykh et al. 2004; Ausseloos et al. 2004; Dziembowski & Pamyatnykh 2008). Finally, the seismic modelling of those B stars provided values for their fundamental parameters, such as the effective temperature, $\log g$ and age.

In this paper we are concerned with data assembled with the CoRoT (Michel et al. (2006)) satellite. Its main asteroseismology mission is to observe a limited number of bright main sequence stars during a long time span of maximum five months, but in between those long runs also shorter ones of some three to five weeks occur. Because of the success of the ground-based asteroseismology of B stars, several B stars with a variety of spectral type were selected as targets for CoRoT, with the aim to map the instability strips of the β Cep and SPB stars.

This paper is devoted to HD 48977, a B2.5V star (V magnitude of 5.92) which was observed by CoRoT as a secondary asteroseismology target during a short run in 2008. Its spectral type of B2.5V places it in the joint region of the SPB and β Cep instability strip. Very little was known about this star prior to the CoRoT observations. It had been flagged as a potential rotationally or pulsationally variable star and it was observed by Hipparcos, which led to a main frequency peak at 0.5218 d^{-1} (Koen & Eyer 2002). Estimates of its fundamental parameters using colour indices were given in Lyubimkov et al. (2002) as $T_{\text{eff}} = 17500 \pm 300$, $\log g = 4.09 \pm 0.08$, from which the authors deduced $M = 5.7 \pm 0.3 M_{\odot}$, $R = 3.6 \pm 0.4 R_{\odot}$, an age of $30.6 \pm 10.7 \times 10^6$ yrs, and $d = 252 \pm 40 \text{ pc}$ using Claret (1995) evolutionary tracks.

2. CoRoT high-precision photometry of HD 48977

HD 48977 was observed by CoRoT with a cadence of 32 s during a short run in 2008, from 2454531.48027JD (5 May) to 2454556.76361 (31 May), i.e., during some 25 days. The detrended light curve is shown in Fig. 1. It was obtained by fitting the light curve with a linear polynomial and dividing the light curve by this polynomial in order to convert to flux units and compensate for the loss of gain in the satellite CCD (Auvergne et al. 2009).

Searching for frequencies in the light curve was done independently by AT using Period04 (Lenz & Breger 2005) and by PD using the method in Degroote et al. (2009). The Fourier transform is shown in Fig. 2. It can be seen that the dominant signal occurs in the range below 3 d^{-1} , which is typical for g-mode pulsations in early B stars. The two methods give the same list of frequencies (within the resolution). The resulting 55 frequencies found having an amplitude more than 4 times the noise level are listed in Table 1. The main peak has a frequency of 0.5167 d^{-1} , very close to the frequency 0.5218 d^{-1} of the Hipparcos light curve (Koen & Eyer 2002). We see that HD 48977 presents a rich spectrum dominated by g modes with frequencies between 0.1 d^{-1} and 1.5 d^{-1} . We thus classify HD 48977 as a new SPB star.

An amplitude drop with a factor more than two occurs for the frequencies 35 to 55 in Table 1, which represent power in the p-mode regime, compared to the lower frequency regime




Fig. 1. CoRoT light curve of HD 48977, with an inset to show the short-time behaviour. The bottom panel represents the residuals after prewhitening with a model based on the 55 frequencies listed in Table 1.




Fig. 2. Fourier transform of the light curve.

(Fig. 2). While these 20 listed frequencies above 10 d^{-1} are significant, we are cautious with their interpretation because they occur in the region where instrumental power also occurs (Auvergne et al. (2009), see also Fig.2). In particular, the two highest frequencies in Table 1 are twice and five times the orbital frequency of the satellite. Four additional peaks in Table 1 coincide with (daily aliases of) the satellite orbital frequency, keeping in mind the relatively large frequency uncertainty. Nevertheless, significant intrinsic stellar frequencies occur in the region $[10, 20] \text{ d}^{-1}$. As is also the case for the pulsators HD 50230 (B3V, Degroote et al. (2010)) and HD 43317 (B3IV, Pápics et al. (2012)), the frequency structure in that regime is rather stable pointing towards p modes excited by the kappa mechanism. The autocorrelation of the power spectrum peaks




Fig. 3. Phase plot of the rotational signal, where all signal not related to the rotational frequency and its harmonics has been filtered out.

at a possible frequency separation of 1.55 d^{-1} , but, in view of the limited frequency precision and the unknown identification of the spherical wavenumbers of the modes, we refrain from further interpretation of the frequencies in this regime with the present data set at hand. The noise level in the residual amplitude spectrum, measured between 5 and 10 d^{-1} , amounts to only 1 ppm .

Given that the harmonics of the fourth frequency $f_4 = 0.637 \text{ d}^{-1}$ are clearly present in the frequency spectrum, while there are no harmonics for any of the other frequencies, and given that the shape of the curve in the phase diagram is not typical of nonlinear pulsations (see Fig. 3) we interpret f_4 as the rotation frequency of the star. Also this particular aspect of the frequency spectrum is similar for the B3IV pulsator HD 43317 (Pápics et al. (2012)), although the data set of the latter star is much more conclusive on this matter, given that this target was observed during a long run of CoRoT.

The frequency resolution deduced from the CoRoT light curve is given by $1.5/\Delta T \simeq 0.06 \text{ d}^{-1}$, where T is the entire duration of the observation run, i.e., 25 days. Unfortunately, this is of the order of the separation between the observed g-mode frequencies (see Table 1). This prevented us from detecting any meaningful period spacings, as deduced for HD 50230 by Degroote et al. (2010), due to the fact that those spacings would be of the same order as the frequency resolution itself.

The frequency spectrum we have at hand is of too low precision to perform a meaningful seismic modelling of this star. This is not surprising, as SPB stars are known to need observations of several months before the frequency spectrum can be deduced with sufficient precision to interpret it in terms of predicted frequencies from theoretical models. The selection of this secondary seismology target for a short run was based on its rough position in the instability strips, which could have implied the detection of dominant high-amplitude p modes. A good strategy for further understanding the target would be to re-observe it with CoRoT during a medium to long run of several months.

3. Spectroscopy of HD 48977

One high resolution high signal-to-noise spectrum of HD 48977 was obtained using the HERMES spectrograph (Raskin et al. 2011) at the Mercator Telescope. The normalised spectrum for

Table 1. Frequencies in the light curve of HD 48977. SN2 is the signal-to-noise in a 6 d^{-1} bin around the peak after each prewhitening step. Phases are given with respect to $t_0 = 0$, and in units of $2\pi \text{ rad}$.

nr	A (ppm)	$\sigma(A)$ (ppm)	f (d^{-1})	$\sigma(f)$ (d^{-1})	ϕ (2π)	$\sigma(\phi)$ (2π)	SN2
1	2052	125	0.5167	0.0011	0.38	0.05	14.7
2	1737	109	0.7933	0.0013	-0.085	0.06	12.6
3	1583	89	0.6832	0.0012	-0.019	0.06	12.8
4	1489	66	0.6370	0.0010	0.26	0.04	13.5
5	1130	44	1.2688	0.0008	0.24	0.04	12.0
6	924	34	0.4542	0.0009	0.37	0.04	9.3
7	753	38	0.3049	0.0012	0.07	0.05	9.1
8	647	27	0.9847	0.0009	0.06	0.04	9.0
9	498	19	0.8958	0.0010	-0.46	0.04	7.1
10	476	21	1.1412	0.0010	-0.42	0.04	7.4
11	430	17.	0.5699	0.0010	0.29	0.04	7.1
12	427	26	0.2100	0.0011	0.40	0.05	7.5
13	393	14	0.3950	0.0009	0.19	0.04	6.7
14	388	17	0.2604	0.0010	-0.28	0.04	7.2
15	330	12	0.3484	0.0009	-0.24	0.04	6.8
16	299	11	1.9182	0.0008	-0.31	0.04	6.7
17	292	9	0.0410	0.0007	0.15	0.03	6.6
18	201	8	1.0766	0.0011	-0.49	0.05	4.5
19	188	9	2.5325	0.0010	-0.33	0.05	4.7
20	182	8	0.1729	0.0010	0.36	0.05	4.3
21	175	7	2.7611	0.0010	-0.49	0.05	4.2
22	152	6	0.6008	0.0010	-0.33	0.05	4.2
23	143	7	2.7110	0.0010	-0.12	0.05	4.1
24	142	6	0.8440	0.0010	-0.31	0.04	4.3
25	127	6	1.3232	0.0010	-0.42	0.05	4.0
26	125	7	2.0672	0.0010	-0.48	0.05	4.1
27	124	6	1.7022	0.0010	-0.36	0.05	4.1
28	123	5	1.8443	0.0010	-0.33	0.05	4.1
29	115	5	3.0379	0.0010	0.36	0.04	4.1
30	112	5	1.0288	0.0010	-0.49	0.05	4.0
31	108	5	0.7579	0.0010	-0.30	0.04	4.1
32	107	4	2.6192	0.0010	-0.45	0.04	4.0
33	104	4	2.9035	0.0010	-0.33	0.04	4.0
34	94	4	2.8083	0.0009	0.05	0.04	4.2
35	44	2.5	14.7599	0.0013	-0.20	0.06	7.1
36	43	2.5	13.4966	0.0013	0.42	0.06	6.4
37	43	2.5	14.4568	0.0013	-0.23	0.06	7.9
38	33	1.9	16.3164	0.0013	-0.36	0.06	6.7
39	29	1.8	11.1543	0.0013	-0.49	0.06	6.4
40	22	1.5	19.4616	0.0014	-0.48	0.06	5.6
41	22	1.5	10.484	0.0014	-0.41	0.06	4.5
42	21	1.4	17.9453	0.0014	-0.25	0.06	4.8
43	21	1.4	15.7987	0.0014	0.069	0.06	5.0
44	20	1.3	14.6173	0.0014	0.20	0.06	5.5
45	19	1.3	19.5002	0.0014	-0.18	0.06	5.8
46	18	1.1	15.6412	0.0014	0.097	0.06	4.7
47	18	1.2	17.6694	0.0014	-0.14	0.06	4.9
48	17	1.2	16.0906	0.0014	0.0027	0.06	4.7
49	17	1.0	12.872	0.0013	0.48	0.06	4.8
50	17	1.1	11.0296	0.0014	0.15	0.06	4.1
51	14	0.9	21.2714	0.0014	0.11	0.07	5.3
52	14	0.9	15.9682	0.0014	0.32	0.06	4.3
53	9.5	0.7	22.9226	0.0017	0.13	0.08	4.7
54	8.5	0.7	27.9437	0.0017	0.47	0.08	7.1
55	3.2	0.5	69.8564	0.0034	0.33	0.15	5.3

the spectral range $4065\text{-}4155\text{\AA}$ is shown in Fig. 4. It has a resolving power of $80\,000$ and a S/N ratio of about 170 in the region near the Si II doublet at $4128, 4130\text{\AA}$ which is a spectral range most often used to study the oscillations of SPB stars (Aerts et al. 1999). From the Fourier transform of the SiIII 4567




Fig. 4. Excerpt of the high resolution high signal-to-noise spectrum (black) of HD 48977, obtained with the HERMES spectrograph attached to the Mercator Telescope. Grey lines indicate the zero-rotation line depths of a template spectrum with $T_{\text{eff}} = 20\,000\text{ K}$ and $\log g = 4.2\text{ dex}$ (solar mixture and solar metallicity).

Table 2. Si lines used to determine the effective temperature in B stars. Lines present in the spectrum of HD 48977 are underlined. Lines that should be avoided according to Simón-Díaz (2010) are listed in *italic*.

Si II	Si III	Si IV
<u>4128.054</u>	<u>4567.840</u>	4212.405
<u>4130.894</u>	<u>4574.757</u>	<i>6701.12</i>
<i>5056.150</i>	<i>4813.33</i>	
<u>6371.371</u>	4829.073	
	5716.287	
	<u>5739.734</u>	

line, which allows the disentangling of non-rotational broadening from other broadening mechanisms (i.e., pulsations, micro- and macroturbulence, Simón-Díaz & Herrero 2007), we deduce $v \sin i = 29 \pm 1\text{ km s}^{-1}$. Combined with the interpretation of f_4 as the rotation frequency and a canonical radius of B-type stars in this part of the Hertzsprung-Russell diagram ($R = 3.5 \pm 0.7 R_{\odot}$, obtained from a grid of models calculated using CLES (Scuflaire et al. 2008)), we derive an inclination angle $i = 15 \pm 3^{\circ}$, or a true equatorial velocity $v_{\text{eq}} = 113 \pm 23\text{ km s}^{-1}$.

We performed a detailed analysis of the spectrum of HD 48977 using the method explained in Morel et al. (2006) to determine the fundamental parameters and the element abundances of the star. The synthetic spectra used have been computed using an updated version of the non-LTE line-formation codes DETAIL/SURFACE (Butler & Giddings (1985); Giddings (1981))

After normalizing the spectrum, we measured the ratios of the equivalent widths (EWs) of the different ionization lines of Si, in order to obtain rough estimates of the effective temperature. Some Si lines appropriate for use in the case of B stars are listed in Table 2. Among those lines, all four Si II lines are present in the spectrum of HD 48977, while only three Si III lines are present and no Si IV lines. The lines present in the spectrum of HD 48977 are underlined in Table 2. Following Simón-Díaz (2010), some of those lines are badly understood from the view-




Fig. 5. Theoretical EW ratios for $\log g = 4.2$ of the suitable Si II and Si III lines in the spectrum with proper atomic data. Theoretical values are plotted for two values of the microturbulence, $\xi = 1\text{ km s}^{-1}$ (full lines) and $\xi = 5\text{ km s}^{-1}$ (dashed lines), and four values of the Si abundance, $\log \epsilon(\text{Si}) = 7.54, 7.34, 7.14, 6.94\text{ dex}$. The two horizontal lines represent the observed values with their error bar.

point of atomic data and are better avoided in the analysis. Those lines are indicated in *italic* in Table 2. Looking at Table 2, we see that we can use only one Si II line and three Si III lines. The ratios of the EWs of the Si II and Si III lines are shown in Fig. 5 for $\log g = 4.2$. The theoretical ratios are shown for two values of the microturbulence, namely $\xi = 1\text{ km s}^{-1}$ and $\xi = 5\text{ km s}^{-1}$ and for four values of the Si abundance, $\log \epsilon(\text{Si}) = 7.54, 7.34, 7.14, 6.94\text{ dex}$. This gives an estimate for the effective temperature of $T_{\text{eff}} = 19\,900 \pm 400\text{ K}$. Given the fact that we have so few suitable Si lines, and for completeness, we did the same analysis using all the observed Si lines. From the lines that should not be used according to Simón-Díaz (2010), we get higher values for the effective temperature, up to $22\,000\text{ K}$. This is consistent with the general trend reported by Simón-Díaz (2010).

The next step is to find the best value for $\log g$ by fitting the wings of the Balmer lines. We use the H_{β} , H_{γ} , H_{δ} , H_{ϵ} , and H_{α} lines. We find the best fit for $\log g = 4.2 \pm 0.1$ assuming $\log T_{\text{eff}} = 20\,000 \pm 1\,000\text{ K}$. In Fig. 6, we show HD 48977 in the B star instability strips. We see that it falls nicely in the SPB instability region.

In principle, the value for the microturbulence parameter can be found by looking at strong and weak lines of a given ion, and require that they yield the same abundances. Unfortunately, all O lines in the spectrum of HD 48977 are weak and we cannot determine the value of the microturbulence. It is, however, expected to be small for this spectral type. We included the unknown value of the microturbulence parameter for the element abundances by increasing the error bars based on two possible values of $\xi = 1\text{ km s}^{-1}$ and $\xi = 5\text{ km s}^{-1}$.

To refine the estimate of the effective temperature, we required the Si II and Si III lines to return on average the same abundance. Using the Si lines underlined in Table 2, we obtain a value consistent with the previous estimate and $\log \epsilon(\text{Si}) = 7.27 \pm 0.20\text{ dex}$.

We can now obtain the abundances of the other elements with lines in the spectrum. The lines used for the analysis and the results obtained for the abundances are summarized in Table 3.




Fig. 6. Instability strips for B stars, as well as observed SPB stars from ground-based surveys (dots) taken from Pamyatnykh (1999). The position of HD 48977 is indicated as square.

The abundance uncertainties take into account both the line-to-line scatter and the errors arising from the uncertainties in the atmospheric parameters. They are in very good agreement with the abundances found for O and B stars in the solar neighbourhood by Morel et al. (2006) using exactly the same methods and tools. The Ne abundance is found to be 7.96 dex, a value intermediate between the solar values recommended by Grevesse & Sauval (1998) and Asplund et al. (2009). Morel & Butler (2008) obtained a very similar Ne abundance for B stars.

The CNO abundance ratios are found to be $[N/O]=-0.92$ and $[N/C]=-0.61$, which are equal to the solar values within the errors. We thus do not find any indication of deep mixing in HD 48977. Such mixing was found for several among the bright β Cep stars (Morel et al. 2006), which are somewhat hotter than HD 48977.

4. Photometric determination of the effective temperature

The visual magnitude of HD 48977 makes it an optimal target for inclusion in wide-field surveys. We collected absolute photometry in the UV (TD1, ANS, Thompson et al. 1978; Wesselius et al. 1980, 1982), the optical (Strømgren, Geneva, Johnson and Tycho Mermilliod et al. 1997; Rufener 1988; Rufener & Nicolet 1988; Ofek 2008; Mermilliod 1986; Høg et al. 2000) and the (near) infrared (Johnson, 2MASS, WISE, AKARI, Cutri et al. 2003; Wright et al. 2010; Cutri et al. 2011; Ishihara et al. 2010). From the WISE bands at $3.4\ \mu\text{m}$ (W1), $4.6\ \mu\text{m}$ (W2), $12.3\ \mu\text{m}$ (W3) and $22\ \mu\text{m}$ (W4) and AKARI photometry at $8.7\ \mu\text{m}$, we see no signs of infrared excess.

Even though it is well known that fundamental parameters determined from SEDs based on photometric data are usually less precise than those deduced from high-resolution spectroscopy, we estimated the effective temperature and gravity via the procedure outlined in Degroote et al. (2011), as a compatibility check. We chose to fit the absolute fluxes in the infrared ($\lambda > 1\ \mu\text{m}$), and only colours at shorter wavelengths (cf. the infrared flux method Blackwell et al. 1979) (Fig. 7). We used both LTE (Castelli & Kurucz 2003) and NLTE model atmospheres (Lanz & Hubeny 2007). We deduced an effective temperature $T_{\text{eff}} = 18\,000 \pm 1\,300\ \text{K}$ (95% confidence interval), surface grav-




Fig. 7. Observed and model photometric colors of HD 48977. Crosses denote the synthetic flux of the best fitting model, circles denote the observations and their errors. Filled circles are included in the fit as colors (optical and UV), open circles are included as absolute photometry (infrared).

ity $\log g > 3.6$, interstellar extinction $E(B-V) < 0.025$, angular diameter $\theta = 0.128 \pm 0.004\ \text{mas}$ and a solar or sub-solar metallicity. We thus find that the photometric method underestimates the effective temperature by roughly 1 000 K compared to the silicon line ratios (Fig. 5) but the values are consistent within 2σ . Given the much higher quality of the HERMES spectrum compared to broad-band photometry, we adopt the spectroscopic estimate of the effective temperature.

5. Conclusions

From 25 d of CoRoT high-precision space photometry, we found the B2.5V star HD 48977 to be a new SPB star. In fact, it is the hottest confirmed SPB star known to date. In agreement with the star's position in the joint part of the β Cep and SPB instability strips, we detected numerous dominant g modes and low-amplitude p modes in its oscillation spectrum. The g modes occur in the frequency range $0.17 - 3.0\ \text{d}^{-1}$ and reach a maximum amplitude of 2000 ppm. The ratio of the p- and g-mode amplitudes of this moderately rotating star of solar metallicity is less than 0.025. In this aspect, the three stars HD 48977 (B2.5V, moderate rotator; this paper), HD 50230 (B3V, slow rotator; Degroote et al. (2010)) and HD 43317 (B3IV, fast rotator; Pápics et al. (2012)) are very comparable.

While the frequency precision is insufficient to perform forward modelling at this stage, an interesting aspect of HD 48977's variability is that we detected rotational modulation, in addition to the g-mode oscillations, in its photometric light curve, again similar to the case of HD 43317. The combination of the rotation frequency we found in the CoRoT light curve and the overall line broadening in the spectrum, implies that HD 48977 is one of the very few moderate rotators known within the class of SPB stars so far, the B3V star HD 43317 being an even faster rotator (Pápics et al. (2012)).

The rotation frequency of $0.6372\ \text{d}^{-1}$ is of the same order as the g-mode frequencies and implies that any future seismic modelling of the star must take into account the effects of rotation in the pulsation description. While this can be done using the traditional approximation for modes whose frequency is above twice

Table 3. Non-LTE elemental abundances of HD 48977 on the scale in which $\log \epsilon[\text{H}]=12$. Mean values obtained using the same techniques for nearby, early B-type stars showing no signs of mixing (i.e., with unaltered CNO abundances) are shown for comparison (Morel & Butler 2008; Morel et al. 2008).

element	ion used	lines used (Å)	HD 48977	Nearby B stars
C	C II	5133.11	8.14±0.10	8.21±0.09
N	N II	3994.00, 4607.16, 4643.09, 5005.15, 5679.55	7.53±0.12	7.67±0.11
O	O II	4069.75, 4414.90, 4641.81, 4649.13, 4661.63, 4705.35	8.45±0.12	8.51±0.09
Ne	Ne I	6143.06, 6266.49, 6334.43, 6382.99, 6402.25, 6506.53	7.96±0.12	7.97±0.07
Mg	Mg II	7877.05	7.61±0.10	7.39±0.11
Al	Al III	4149.96, 4512.54, 4529.04, 5696.6, 5722.7	6.21±0.28	6.13±0.08
Si		see Table 2	7.27±0.20	7.18±0.07
S	S II	4162.48, 4524.81, 4815.55	7.11±0.18	7.20±0.11
Fe	Fe III	4419.60, 5156.11	7.21±0.14	7.28±0.10

the rotation frequency (Aerts & Dupret 2012; Ballot et al. 2012), the lowest-frequency modes of HD 48977 correspond with so-called gravito-inertial waves and require a treatment not relying on perturbative methods (Ballot et al. 2010).

Our findings show that HD 48977 would be an excellent target for a long run during an extension of the CoRoT mission. This would probably allow to decide on the nature of the power in the frequency regime above 10 d^{-1} and to unravel rotational splitting of several of the pulsation modes, and hence would open stellar modelling applications from the mode identification of the multiplet structures.

Acknowledgements. A.T., P.D., and C.A., are grateful to the staff of the Kavli Institute of Theoretical Physics (KITP) of the University of California at Santa Barbara (UCSB) for the hospitality during their stay in the framework of the 2011 Research Programme “Asteroseismology in the Space Age”. T.M. acknowledges financial support from Belspo for contract PRODEX GAIA-DPAC. The research leading to these results has received funding from the European Research Council under the European Community Seventh Framework Programme (FP7/20072013)/ERC grant agreement n°227224 (PROSPERITY), from the Belgian Science Policy Office (Belspo) under PRODEX contract C90309: CoRoT Data Exploitation, and from the National Science Foundation of the United States under grant n°NSF PHY05–51164, and from the FNRS. The CoRoT space mission was developed and is operated by the French space agency CNES, with participation of ESAs RSSD and Science Programmes, Austria, Belgium, Brazil, Germany, and Spain. Based on observations made with the HERMES spectrograph, installed at the Mercator Telescope, operated on the island of La Palma by the Flemish Community, at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias and supported by the Fund for Scientific Research of Flanders (FWO), Belgium, the Research Council of K.U.Leuven, Belgium, the Fonds National de la Recherche Scientifique (F.R.S.–FNRS), Belgium, the Royal Observatory of Belgium, the Observatoire de Genève, Switzerland and the Thüringer Landessternwarte Tautenburg, Germany.

References

Aerts, C., Briquet, M., Degroote, P., Thoul, A., & van Hoolst, T. 2011, *A&A*, 534, A98
Aerts, C., Christensen-Dalsgaard, J., & Kurtz, D. W. 2010, *Asteroseismology*, ed. Springer
Aerts, C., De Cat, P., Handler, G., et al. 2004a, *MNRAS*, 347, 463
Aerts, C., De Cat, P., Kuschnig, R., et al. 2006a, *ApJ*, 642, L165
Aerts, C., De Cat, P., Peeters, E., et al. 1999, *A&A*, 343, 872
Aerts, C. & Dupret, M.-A. 2012, in *Astronomical Society of the Pacific Conference Series*, Vol. 462, *Progress in Solar/Stellar Physics with Helio- and Asteroseismology*, ed. H. Shibahashi, M. Takata, & A. E. Lynas-Gray, 103
Aerts, C., Lehmann, H., Briquet, M., et al. 2003a, *A&A*, 399, 639
Aerts, C., Marchenko, S. V., Matthews, J. M., et al. 2006b, *ApJ*, 642, 470
Aerts, C., Thoul, A., Daszyńska, J., et al. 2003b, *Science*, 300, 1926
Aerts, C., Waelkens, C., Daszyńska-Daszkiewicz, J., et al. 2004b, *A&A*, 415, 241
Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
Ausseloes, M., Scuflaire, R., Thoul, A., & Aerts, C. 2004, *MNRAS*, 355, 352
Auvergne, M., Bodin, P., Boisnard, L., et al. 2009, *A&A*, 506, 411

Ballot, J., Lignières, F., Prat, V., Reese, D. R., & Rieutord, M. 2012, in *Astronomical Society of the Pacific Conference Series*, Vol. 462, *Progress in Solar/Stellar Physics with Helio- and Asteroseismology*, ed. H. Shibahashi, M. Takata, & A. E. Lynas-Gray, 389
Ballot, J., Lignières, F., Reese, D. R., & Rieutord, M. 2010, *A&A*, 518, A30
Balona, L. A., Pigulski, A., Cat, P. D., et al. 2011, *MNRAS*, 413, 2403
Blackwell, D. E., Shallis, M. J., & Selby, M. J. 1979, *MNRAS*, 188, 847
Briquet, M., Lefever, K., Uytterhoeven, K., & Aerts, C. 2005, *MNRAS*, 362, 619
Briquet, M., Morel, T., Thoul, A., et al. 2007, *MNRAS*, 381, 1482
Briquet, M., Uytterhoeven, K., Morel, T., et al. 2009, *A&A*, 506, 269
Butler, K. & Giddings, J. R. 1985, in *Newsletter of Analysis of Astronomical Spectra*, Vol. 9
Cameron, C., Saio, H., Kuschnig, R., et al. 2008, *ApJ*, 685, 489
Castelli, F. & Kurucz, R. L. 2003, in *IAU Symposium*, Vol. 210, *Modelling of Stellar Atmospheres*, ed. N. Piskunov, W. W. Weiss, & D. F. Gray, 20P
Christensen-Dalsgaard, J. 2002, *Reviews of Modern Physics*, 74, 1073
Claret, A. 1995, *A&AS*, 109, 441
Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *2MASS All Sky Catalog of point sources*, (NASA/IPAC Infrared Science Archive)
Cutri, R. M., Wright, E. L., Conrow, T., et al. 2011, *Explanatory Supplement to the WISE Preliminary Data Release Products*, Tech. rep.
De Ridder, J., Teltng, J. H., Balona, L. A., et al. 2004, *MNRAS*, 351, 324
Degroote, P., Acke, B., Samadi, R., et al. 2011, *A&A*, 536, A82
Degroote, P., Aerts, C., Baglin, A., et al. 2010, *Nature*, 464, 259
Degroote, P., Briquet, M., Catala, C., et al. 2009, *A&A*, 506, 111
Desmet, M., Briquet, M., Thoul, A., et al. 2009, *MNRAS*, 396, 1460
Diago, P. D., Gutiérrez-Soto, J., Auvergne, M., et al. 2009, *A&A*, 506, 125
Dupret, M.-A., Thoul, A., Scuflaire, R., et al. 2004, *A&A*, 415, 251
Dziembowski, W. A. & Pamyatnykh, A. A. 2008, *MNRAS*, 385, 2061
Giddings, J. R. 1981, *PhD thesis*, University of London
Grevesse, N. & Sauval, A. J. 1998, *Space Sci. Rev.*, 85, 161
Gutiérrez-Soto, J., Floquet, M., Samadi, R., et al. 2009, *A&A*, 506, 133
Handler, G., Jerzykiewicz, M., Rodríguez, E., et al. 2006, *MNRAS*, 365, 327
Handler, G., Shobbrook, R. R., Jerzykiewicz, M., et al. 2004, *MNRAS*, 347, 454
Handler, G., Shobbrook, R. R., & Mokgwetsi, T. 2005, *MNRAS*, 362, 612
Høg, E., Fabricius, C., Makarov, V. V., et al. 2000, *A&A*, 355, L27
Huat, A.-L., Hubert, A.-M., Baudin, F., et al. 2009, *A&A*, 506, 95
Ishihara, D., Onaka, T., Kataza, H., et al. 2010, *A&A*, 514, A1
Jerzykiewicz, M., Handler, G., Shobbrook, R. R., et al. 2005, *MNRAS*, 360, 619
Koen, C. & Eyer, L. 2002, *MNRAS*, 331, 45
Lanz, T. & Hubeny, I. 2007, *ApJS*, 169, 83
Lenz, P. & Breger, M. 2005, *Communications in Asteroseismology*, 146, 53
Lyubimkov, L. S., Rachkovskaya, T. M., Rostopchin, S. I., & Lambert, D. L. 2002, *MNRAS*, 333, 9
Mermilliod, J., Mermilliod, M., & Hauck, B. 1997, *A&AS*, 124, 349
Mermilliod, J. C. 1986, *Bulletin d'Information du Centre de Données Stellaires*, 31, 185
Michel, E., Baglin, A., Auvergne, M., et al. 2006, in *ESA Special Publication*, Vol. 1306, *ESA Special Publication*, ed. M. Fridlund, A. Baglin, J. Lochard, & L. Conroy, 39
Morel, T. & Butler, K. 2008, *A&A*, 487, 307
Morel, T., Butler, K., Aerts, C., Neiner, C., & Briquet, M. 2006, *A&A*, 457, 651
Morel, T., Hubrig, S., & Briquet, M. 2008, *A&A*, 481, 453
Neiner, C., Gutiérrez-Soto, J., Baudin, F., et al. 2009, *A&A*, 506, 143
Ofek, E. O. 2008, *PASP*, Chicago, 120, 1128
Pamyatnykh, A. A. 1999, *Acta Astron.*, 49, 119
Pamyatnykh, A. A., Handler, G., & Dziembowski, W. A. 2004, *MNRAS*, 350, 1022
Pápics, P. I., Briquet, M., Auvergne, M., et al. 2011, *A&A*, 528, A123

- Pápics, P. I., Briquet, M., Baglin, A., et al. 2012, *A&A*, 542, A55
- Raskin, G., van Winckel, H., Hensberge, H., et al. 2011, *A&A*, 526, A69
- Rufener, F. 1988, Catalogue of stars measured in the Geneva Observatory photometric system : 4 : 1988
- Rufener, F. & Nicolet, B. 1988, *A&A*, 206, 357
- Saio, H., Cameron, C., Kuschnig, R., et al. 2007, *ApJ*, 654, 544
- Scuflaire, R., Théado, S., Montalbán, J., et al. 2008, *Ap&SS*, 316, 83
- Simón-Díaz, S. 2010, *A&A*, 510, A22+
- Simón-Díaz, S. & Herrero, A. 2007, *A&A*, 468, 1063
- Thompson, G. I., Nandy, K., Jamar, C., et al. 1978, Catalogue of stellar ultraviolet fluxes. A compilation of absolute stellar fluxes measured by the Sky Survey Telescope (S2/68) aboard the ESRO satellite TD-1, (NASA/STI)
- Wesselius, P. R., van Duinen, R. J., Aalders, J. W. G., & Kester, D. 1980, *A&A*, 85, 221
- Wesselius, P. R., van Duinen, R. J., de Jonge, A. R. W., et al. 1982, *A&AS*, 49, 427
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868