

**MEUSE-RHINE GEOLOGISTS MEETING
MONT-RIGI (HAUTES FAGNES)
MAY 18-19, 1990**

**Organized by M. STREEL and M.J.M. BLESS
with the cooperation of the «Société géologique de Belgique»**

**Location : Station scientifique des Hautes Fagnes
(Dir. : Prof. R. Schumacker), 4950 - Robertville-Waimes (Belgium)**

**BIOSTRATIGRAPHICAL CORRELATION
BETWEEN THE HANGENBERG
SCHIEFER (NORTHERN
«RHEINISCHES SCHIEFERGEBIRGE»)
AND GLACIGENIC DEPOSITS
IN BRAZIL**

M. STREEL¹ & S. LOBOZIAK²

ABSTRACT.- Clastic sedimentary sequences with glacigenic characteristics of Late Devonian age are described in several basins of Brazil. They were dated as Famennian *sensu lato*. Samples from these sequences in the Amazonas and Parnaiba basins contain well preserved miospores which allow a accurate correlation with the uppermost Famennian (Middle *praesulcata* Zone

133

equivalent) *R. lepidophyta* - *H. explanatus* and *R. lepidophyta* - *V. nitidus* Zones (LE and LN Zones). The same miospore-zones characterize the Hangenberg Schiefer and Sandstein in the Northern «Rheinisches Schiefergebirge» which are considered to correspond to a sudden drop of sea level. Such an accurate time relationship between glacigenic sediments in Brazil and drop of sea-level in Germany (and around the world !) suggests that they might have the same climatic origin.

¹. Paléontologie, Université de Liège, 7, place du Vingt-Août, B-4000 LIEGE, Belgium.

². Paléobotanique, Université des Sciences et Techniques de Lille, URA 1365, F-59655 VILLENEUVE D'ASCO, France.