

ABOUT SOME SKELETAL PARTICULARITIES OF THE FIRST VERTEBRAE RELATED TO THE MODE OF PREY CAPTURE IN *URANOSCOPIUS SCABER* (URANOSCOPIDAE)

by

Laurence HUET, Véronique GOOSSE, Eric PARMENTIER & Pierre VANDEWALLE* (1)

ABSTRACT. - Feeding in *Uranoscopus scaber* is characterised by bending of the body just behind the head. Bending is related to the organisation of the first five vertebrae: vertebral centra of lesser height fitting one into the next; short, less tilted neural spines, presence of many intervertebral ligaments. Bending of the vertebral column causes the head to rotate upward and the mouth to open just beneath the prey; it also brings the pharyngeal jaws, shifted with respect to each other at rest, to face each other.

RÉSUMÉ. - Particularités squelettiques des premières vertèbres en relation avec le mode de prise de nourriture chez *Uranoscopus scaber* (Uranoscopidae).

La prise de nourriture chez *Uranoscopus scaber* est caractérisée par une pliure du corps, située à l'arrière de la tête. Cette pliure est en rapport avec l'organisation des cinq premières vertèbres: corps vertébraux peu élevés et emboîtés les uns dans les autres, neurépinnes courtes et redressées, présence de nombreux ligaments intervertébraux. Cette pliure entraîne une rotation de la tête qui amène la bouche à s'ouvrir vers le haut immédiatement sous la proie et place les mâchoires pharyngiennes, décalées au repos, les unes en face des autres, les rendant aptes à saisir la proie.

Key-words. - Uranoscopidae, *Uranoscopus scaber*, Feeding, Vertebral column.

The stargazer, *Uranoscopus scaber* Linné, 1758, is a Mediterranean fish that feeds by lying in wait for its prey. Its head is flat and wide, corresponding with a benthic life. Buried in the sand almost to its eyes, it attracts its prey by means of a protractile appendix attached to the mandible and waved like a lure (Bauchot and Pras, 1980; Pietsch, 1989). At the right moment, it lunges quickly out of the substrate and catches the prey, or tries to. The lunge is caused by bending between the head and trunk, by 60° and more (Goosse *et al.*, 1995). The aim of the present paper is to examine whether there are special adaptations of the anterior vertebral column that make this bending possible, and to assess changes in the disposition of cephalic skeletal structures as bending occurs. The cephalic skeleton has been described by Pietsch (1989) and will thus not be described again here.

(1) Université de Liège, Institut de Zoologie, Laboratoire de Morphologie fonctionnelle et évolutive, 22 Quai Van Beneden, B-4020 Liège, Belgique. [P.Vandewalle@ulg.ac.be]

* Author for correspondance.

MATERIAL AND METHODS

Twelve dead *Uranoscopus scaber* specimens (standard length between 16 and 25.5 cm) were stored frozen or in alcohol. One was cleared with trypsin and stained with alizarin red for observation of the bony skeleton, the others were dissected and examined with a Wild M10 stereobinocular microscope equipped with a camera lucida and a photographic tube.

Three live specimens (standard length: 17.5, 19 and 22 cm) were acclimated to living and feeding on live prey in an aquarium. They were filmed at 400 fps with a Photosonics IPL camera (Eastman Kodak High Speed film, 400 ASA, 3200K°).

All specimens studied were from Calvi Bay (STARESO station) and Lion Gulf (Grau du Roy aquarium), France.

Abbreviations used in the figures

BOC: basioccipital	NA: neural arch	POZ: postzygapophysis
E: eye	NS: neural spine	PZ: prezygapophysis
HY: hyobranchial apparatus	OP: opercle	SPNF: spinal nerve foramen
LI 1,2,3,4,5: ligament 1 to 5	PF: pectoral fin	VC: vertebral centrum
MAX: maxilla	PJ: pharyngeal jaw	V1,2,3,4,5,6: vertebra 1 to 6
MD: mandible	PAP: parapophysis	

RESULTS

Movements

Bending of the vertebral column in *Uranoscopus scaber* occurs when a live prey passes above its head (Fig. 1). Bending causes a slight elevation of the trunk and a major lifting of the neurocranium. It always takes less than 3/100 s, after which the fish resumes its initial position at a quite variable speed (32/100 s to 55/100 s).

Vertebrae

The first five abdominal vertebrae display several original features. They bear broader neural spines than the following vertebrae. The first neural spine is shorter than the neurocranium height. The length of the neural spines increases from the first to the sixth vertebra and these first neural spines are not tilted as far backward as the following ones. Viewed from the side (Fig. 3), the first four vertebral centra are lower than those of the following vertebrae; they display no prominent parapophyses (Fig. 2).

The first two vertebrae bear no ribs. The first vertebra is longer than the following ones and is attached to the occipital region by short fibres, from the top of the short neural spine to the base of the vertebral centrum; this makes it almost completely immobile. The next three vertebrae are short and their vertebral centra are wider than high. The anterior end of each one is narrower than the posterior end of the preceding one, so that the first four vertebrae interlock. The second, third, and fourth neural arches are situated quite forward on the vertebral centra. The front part of each vertebral centrum is narrower than the rear part of the preceding one. They bear prezygapophyses with extended, outward-turned articulation surfaces. Each of these surfaces is coapted to that of the inward-turned postzygapophysis of the preceding vertebra (Fig. 3).




Fig. 1. - *Uranoscopus scaber*. Scheme and below, the fish with raised head. The prey has already been sucked into the buccal cavity at 400 frames per second.



Fig. 2. - *Uranoscopus scaber*. A: Vertebral centra of the LI.5 ligaments. B: Lateral view of the vertebrae where the postzygapophysis articulates.

The third vertebra bears no ribs. The fourth is flanked by ribs, articulating with the fifth.

DS

Standard length between 16 and 22 cm were acclimated to light with trypsin and stained with alizarin red. They were dissected and examined under a camera lucida and a photo-

microscope (22 cm) were acclimated to light with trypsin and stained with alizarin red. They were dissected and examined under a camera lucida and a photo-

microscope (22 cm) were acclimated to light with trypsin and stained with alizarin red. They were dissected and examined under a camera lucida and a photo-

- POZ: postzygapophysis
- VC: vertebral centrum
- SPNF: spinal nerve foramen
- NA: neural arch
- NS: neural spine
- Li5: ligament 5
- PAP: postzygapophysis
- VC1-6: vertebra 1 to 6




Fig. 1. - *Uranoscopus scaber*. Schematic lateral views showing above, the fish lying in wait for its prey, and below, the fish with raised head, when the mouth is wide open and bending is maximal. The prey has already been sucked into the buccal cavity. The drawings were made from frames of a film made at 400 frames per second.

Uranoscopus scaber occurs when a live prey is captured. It causes a flexion of the trunk and a major loss of equilibrium, after which the fish resumes its normal position (55/100 s).

Original features. They bear a broad neural spine which is shorter than the dorsal spine from the first to the sixth vertebrae as the following ones. The neural spines are lower than those of the other species (Fig. 2).

The first neural arch is longer than the following ones, from the top of the short neural spine it almost completely immo- bilizes the centra. The neural arches are wider than high. The width of the preceding one, so that the fourth neural arches are situated above the fifth vertebral centrum is narrower than the preceding ones with extended, the postzygapophysis is coapted to that of the intervertebral centrum (Fig. 3).




Fig. 2. - *Uranoscopus scaber*. A: Ventral view of the first vertebrae: the interrupted lines show the limits of the Li5 ligaments. B: Lateral view of the second vertebra: the interrupted lines situate the zone where the postzygapophysis articulates with the prezygapophysis of the next vertebra.

The third vertebra bears a pair of ribs at the upper extremity of the neural arch. The fourth is flanked by ribs, articulated on the lateral walls of the neural arch. Not until the




Fig. 3. - *Uranoscopus scaber*. A: Lateral view of the first vertebrae and associated ligaments; B: Lateral view of the positions of these vertebrae when bending exceeds 45°. Asterisks indicate where the ribs articulate with the vertebrae.

fifth vertebra do the ribs appear as usual on the parapophyses. The fifth vertebra looks partially "normal". Although the front of its vertebral centrum is still narrow and flattened, its posterior surface has the same shape as the following vertebra. The parapophyses protrude ventro-laterally at the front of the vertebral centrum. The neural arch is located almost completely above the vertebral centrum and is extended by a more tapered neural spine.

All of the vertebrae are attached to each other by ligaments. A large dorsal septum (LI.1) encloses the neural spines from the rear to the fifth vertebra; then it thickens and shortens as the height of the neural spines decreases. It attaches to the rear of the neurocranium where it spreads out. A part of the epaxial musculature (wide and voluminous in the back of the head) inserts onto this septum.

Thick ligaments (LI.2) link each neural arch with the dorsal side of the prezygapophysis of the next vertebra. The first four ligaments are a little more tilted than the following ones, which are wider and thinner. In between the first five vertebrae there is a ligament LI.3, almost horizontal and partly masked by LI.2 (completely masked between the next vertebrae). Another thick ligament (LI.4) links the inner surface of a prezygapophysis with the postzygapophysis and the vertebral centrum of the preceding vertebra. Lastly, each vertebral centrum is linked to the next by a ligament LI.5 consisting of short fibres. The LI.5 fibres connecting the first four vertebrae appear longer than those of the following ligaments.

Skeletal particularities

Buccal parts and jaws

One original feature of this fish is lying in wait, the two halves of the jaws for prey capture, the neurocranial region is practically the same vertical position as the jaws, almost horizontal (Fig. 1B). This is different from the hyoid depression in comparison with other species (elevation).

Observed on dead animals, the jaws in extended position (Fig. 1A) appear opposite (Fig. 1B).

Teleosts have different systems of the cranium. One possible system is the elevation of the neurocranium (Vandewalle, 1978; Liem, 1982; Sibbing, 1982; Muller, 1987; Muller, 1987; Howes, 1979), 40° for *Lucioperca* (Howes, 1979), 40° for *Lucioperca* (Howes, 1979), 40° for *Lucioperca* (Howes, 1979), 40° for *Lucioperca* (Howes, 1979). Lesiuk (1981) have evidenced the specific modification of extreme markedly bending of the vertebral column. In the characiform fish, the head simply by rotating around the vertical axis (Howes, 1979). In the characiform fish, the head and occurs posterior to the first vertebra and the Weberian apparatus. The fifth vertebra is taller than the following ones and changes in the vertebral column. A large bending angle can be exceptional. The epaxial musculature, fixed on a strong septum (Li, 1989) contracts, vertebrae two to three times as tall as possible by the interlocking of the postzygapophyses, and the difference in the neural arches, enabling them to counteract this rotation are probably the result of the bending of the vertebrae, because of their standard shape to curve differently from what is observed in other species (Li, 1982; Vandewalle *et al.*, 1995). The difference in the vertebral column should be minimal slight elevation of ligaments LI.1, LI.2, LI.3.

In acanthopterygians, during feeding, the mouth, pointed towards the prey, opens first, after which the buccal cavity, then the opercular cavity dilate, causing the prey to be sucked in. Finally the pharyngeal jaws, facing each other, separate to grasp the prey (Lauder, 1980, 1981; Westneat and Wainwright, 1989; Vandewalle *et al.*, 1992, 1995). In *Uranoscopus scaber*, the mouth is turned upward. Opening of the mouth without rotation of the neurocranium would protrude the mouth opening, whereas the prey is located above the predator. Rotation of the neurocranium makes it possible to open the mouth upward, (practically) without modifying the orientation of the mandible. The distance between the mandibular and the premaxillaries increases, nevertheless, as upon any opening of the mouth (Alexander, 1967; Lauder, 1980, 1981; Vandewalle, 1980). This requires participation of the ventral musculature of the head (protractor hyoideus and sternohyoideus muscles) to maintain the mandible at its position. Contraction of this musculature is attested by depression of the buccal cavity bottom (ventral part of the hyobranchial system) with respect to the skull. If this did not occur, the mandible would follow at least partially, like the rest of the head, the rotatory motion of the neurocranium. The originality of mouth opening in *U. scaber* thus does not reside in any special motion of the buccal parts with respect to one another, but solely in the possibility of bending the vertebral column to a larger degree than other teleosts. In the stonefish *Synanceia verrucosa*, which also lies in wait like *U. scaber*, bending causes only raising of the head without motion of the body. In the stonefish the mandible is also practically vertical at rest and remains so during opening of the mouth (Grobecker, 1983).

Bending of the vertebral column most probably has a second beneficial consequence. At rest, the toothed plates of the pharyngeal jaws of *Uranoscopus scaber* do not face each other as in most teleosts (Liem, 1970; Liem and Osse, 1975; Aerts *et al.*, 1986; Gobalet, 1989; Vandewalle *et al.*, 1992); the lower ones are markedly in front of the upper ones. In this position, the pharyngeal jaws would be unable to grasp a prey. Major bending of the front of the vertebral column, causing rotation of the neurocranium, may cause an upward rotation of the upper pharyngeal jaws while the lower ones are retained by the depression of the mouth bottom; this would place the toothed plates one above the other, position enabling them to grasp a prey.

Acknowledgements. - We thank Dr D. Bay (Stareso, Calvi, France) for providing us with specimens of *Uranoscopus scaber*, and Mrs K. Broman for translating the text in English.

REFERENCES

- AERTS P., DE VREE F. & P. VANDEWALLE, 1986. - Pharyngeal jaw movements in *Oreochromis niloticus* (Teleostei: Cichlidae): Preliminary results of a cineradiographic analysis. *Ann. Soc. r. Zool. Belg.*, 116: 75-82.
- ALEXANDER R. McN., 1967. - The functions and mechanism of the protrusible jaws of some Acanthopterygian fishes. *J. Zool., Lond.*, 151: 43-64.
- BAUCHOT M.L. & A. PRAS, 1980. - Guide des Poissons marins d'Europe. 427 p. Lausanne - Paris: Delachaux & Niestlé Edit.
- BERGERT B.A. & P.C. WAINWRIGHT, 1997. - Morphology and kinematics of the prey capture in the syngnathid fishes *Hippocampus erectus* and *Syngnathus floridae*. *Mar. Biol.*, 127: 563-570.
- GOBALET K.W., 1989. - Morphology of the parrotfish pharyngeal jaw apparatus. *Amer. Zool.*, 29: 319-331.
- GOOSSE V., HUET L. & P. VANDEWALLE, 1998. - Les particularités squelettiques chez *Uranoscopus scaber* (Perciformes, Uranoscopusidae). *Env. Biol. Fish.*, 8(3/4): 191-200.
- GROBECKER D.B., 1983. - The "upward" mouth of the stonefish *Synanceia verrucosa*. *Env. Biol. Fish.*, 8(3/4): 191-200.
- HOWES G.J., 1979. - Notes on the feeding mechanism of the stonefish *Synanceia verrucosa* with comments on the Cultridae. *J. Zool., Lond.*, 187: 543-570.
- LAUDER G.V., 1979. - Feeding mechanism of the stonefish *Synanceia verrucosa*. *J. Zool., Lond.*, 187: 543-570.
- LAUDER G.V., 1980. - The suction feeding mechanism of the stonefish *Synanceia verrucosa*. *J. Exp. Biol.*, 88: 43-54.
- LAUDER G.V., 1981. - Intraspecific variation in the feeding mechanism of the stonefish *Synanceia verrucosa*. *J. Zool., Lond.*, 191: 1-16.
- LAUDER G.V. & K.F. LIEM, 1981. - The mechanism of jaw protrusion in teleost fishes. *J. Zool., Lond.*, 191: 1-16.
- LESIAUK T.P. & C.C. LINDSEY, 1981. - The feeding mechanism of the stonefish *Synanceia verrucosa*. *J. Zool., Lond.*, 191: 1-16.
- LIEM K.F., 1970. - Comparative functional anatomy of the mouth of teleost fishes. *J. Zool., Lond.*, 165: 1-166.
- LIEM K.F., 1979. - Modulatory mechanisms of the mouth of teleost fishes. *J. Zool., Lond.*, 187: 1-166.
- LIEM K.F. & J.W.M. OSSE, 1975. - The mouth of African cichlid fishes. *Amer. Zool.*, 15: 1-166.
- MULLER M., 1987. - Optimization of the mouth bottom depression in teleost fishes. *J. Zool., Lond.*, 201: 1-166.
- PIESTSCH T.W., 1989. - Phylogenetic relationships of the stonefish *Synanceia verrucosa*. *Copeia*, 1989: 253-303.
- SIBBING F.A., 1982. - Pharyngeal jaw structure in teleost fishes. *J. Zool., Lond.*, 191: 1-166.
- VANDEWALLE P., 1978. - Analyse cineradiographique et électrocinématique de la dépression buccale chez le *Gobio gobio* (L.) (Poissons, Cyprinidae). *Cybius*, 1: 1-166.
- VANDEWALLE P., 1980. - Etude cineradiographique de la dépression buccale chez le *Gobio gobio* (L.) (Poissons, Cyprinidae). *Cybius*, 3: 1-166.
- VANDEWALLE P., HAVARD M., 1981. - Les particularités squelettiques de la pharyngienne pendant la dépression buccale chez le *Gobio gobio* (L.) (Poissons, Serranidae). *Can. J. Zool.*, 59: 1-166.
- VANDEWALLE P., SAINTIN P. & M. HAVARD, 1982. - Les particularités squelettiques de la pharyngienne pendant la dépression buccale chez le *Gobio gobio* (L.) (Poissons, Serranidae). *Can. J. Zool.*, 60: 1-166.
- VANDEWALLE P., SAINTIN P. & M. HAVARD, 1983. - Les particularités squelettiques de la pharyngienne pendant la dépression buccale chez le *Gobio gobio* (L.) (Poissons, Serranidae). *Can. J. Zool.*, 61: 1-166.
- WESTNEAT M.W., 1994. - Transitional forms of the pharyngeal jaws in related teleost fishes (Teleostei, Perciformes). *J. Zool., Lond.*, 234: 1-166.
- WESTNEAT M.W. & P. WAINWRIGHT, 1989. - Evolution of the pharyngeal jaw apparatus in teleost fishes (Labridae; Teleostei): Evolutionary trends in the structure and function of the pharyngeal jaw apparatus. *J. Zool., Lond.*, 223: 1-166.

Reçu le 16.12.1998.

Accepté pour publication le 14.04.1999.

inted towards the prey, opens
dilates, causing the prey to be
r. separate to grasp the prey
dewalle *et al.*, 1992, 1995). In
of the mouth without rotation
reas the prey is located above
le to open the mouth upward,
ble. The distance between the
as upon any opening of the
(1980). This requires participa-
deus and sternohyoideus mus-
of this musculature is attested
he hyobranchial system) with
d follow at least partially, like
um. The originality of mouth
otion of the buccal parts with
ling the vertebral column to a
verrucosa, which also lies in
l without motion of the body.
at rest and remains so during

as a second beneficial conse-
of *Uranoscopus scaber* do not
osse, 1975; Aerts *et al.*, 1986;
markedly in front of the upper
e to grasp a prey. Major ben-
of the neurocranium, may cause
lower ones are retained by the
ed plates one above the other.

for providing us with specimens of
nglish.

l jaw movements in *Oreochromis*
radiographic analysis. *Ann. Soc. r.*

of the protrusible jaws of some

l'Europe. 427 p. Lausanne - Paris:


cinematics of the prey capture in the
idae. Mar. Biol., 127: 563-570.

l jaw apparatus. *Amer. Zool.*, 29:

- GOOSSE V., HUET L. & P. VANDEWALLE, 1995. - Introduction à l'étude de la prise de nourriture chez *Uranoscopus scaber* L. (Pisces, Perciformes). *Rapp. Comm. int. Mer Médit.*, 34: 244.
- GROBECKER D.B., 1983. - The "lie-in-wait" feeding mode of a criptic teleost, *Synanceia verrucosa*. *Env. Biol. Fish.*, 8(3/4): 191-202.
- HOWES G.J., 1979. - Notes on the anatomy of *Macrochirichthys macrochirus* (Valenciennes), 1844, with comments on the Cultrinae (Pisces, Cyprinidae). *Bull. Brit. Mus. Nat. Hist.*, 36: 147-200.
- LAUDER G.V., 1979. - Feeding mechanic in primitive teleosts and in the halecomorph fish *Amia calva*. *J. Zool., Lond.*, 187: 543-578.
- LAUDER G.V., 1980. - The suction feeding mechanism in sunfishes (*Lepomis*): An experimental analysis. *J. Exp. Biol.*, 88: 49-72.
- LAUDER G.V., 1981. - Intraspecific functional repertoires in the feeding mechanism of the characoid fishes *Lebiasina*, *Hoplias* and *Chalceus*. *Copeia*, 1981: 154-168.
- LAUDER G.V. & K.F. LIEM, 1981. - Prey capture by *Luciocephalus pulcher*: Implications for models of jaw protrusion in teleost fishes. *Env. Biol. Fish.*, 6: 257-268.
- LESIUK T.P. & C.C. LINDSEY, 1978. - Morphological peculiarities in the neck-bending amazonian characoid fish *Rhaphiodon vulpinus*. *Can. J. Zool.*, 56: 991-997.
- LIEM K.F., 1970. - Comparative functional anatomy of the Nandidae (Pisces: Teleostei). *Field. Zool.*, 56: 1-166.
- LIEM K.F., 1979. - Modulatory multiplicity in the feeding mechanism in cichlid fishes, as exemplified by the invertebrate pickers of Lake Tanganyika. *J. Zool., Lond.*, 189: 93-125.
- LIEM K.F. & J.W.M. OSSE, 1975. - Biology, versatility, evolution and food resource partitioning in african cichlid fishes. *Amer. Zool.*, 15: 427-454.
- MULLER M., 1987. - Optimization principles applied to the mechanism of neurocranium levation and mouth bottom depression in bony fishes (Halecostomi). *J. Theor. Biol.*, 126: 343-368.
- PIESTSCH T.W., 1989. - Phylogenetic relationships of trachinoid fishes of the family Uranoscopidae. *Copeia*, 1989: 253-303.
- SIBBING F.A., 1982. - Pharyngeal mastication and food transport in the carp (*Cyprinus carpio* L.): A cineradiographic and electromyographic study. *J. Morphol.*, 172: 223-258.
- VANDEWALLE P., 1978. - Analyse des mouvements potentiels de la région céphalique du goujon *Gobio gobio* (L.) (Poissons, Cyprinidae). *Cybium*, 3: 15-33.
- VANDEWALLE P., 1980. - Etude cinématographique et électromyographique de la prise de nourriture et du crachement chez le goujon, *Gobio gobio* (L.) et la vandoise, *Leuciscus leuciscus* (L.) (Pisces, Cyprinidae). *Cybium*, 5: 3-14.
- VANDEWALLE P., HAVARD M., CLAES G. & F. DE VREE, 1992. - Mouvements des mâchoires pharyngiennes pendant la prise de nourriture chez *Serranus scriba* (Linné, 1758) (Pisces: Serranidae). *Can. J. Zool.*, 70: 145-160.
- VANDEWALLE P., SAINTIN P. & M. CHARDON, 1995. - Structures and movements of the buccal and pharyngeal jaws in relation to feeding in *Diplodus sargus*. *J. Fish Biol.*, 46: 623-656.
- WESTNEAT M.W., 1994. - Transmission of force and velocity in the feeding mechanisms of labrid fishes (Teleostei, Perciformes). *Zoomorphology*, 114: 103-118.
- WESTNEAT M.W. & P. WAINWRIGHT, 1989. - Feeding mechanism of *Epibulus insidiator* (Labridae; Teleostei): Evolution of a novel functional system. *J. Morphol.*, 202: 129-150.

Reçu le 16.12.1998.

Accepté pour publication le 14.04.1999.



Buccal parts and pharyngeal jaws

One original feature of the buccal and pharyngeal jaws is their position. While the fish is lying in wait, the two buccal jaws are almost vertical (Fig. 1A). Upon bending and prey capture, the neurocranial elevation raises the buccal jaws but the mandible retains practically the same vertical position whereas the maxillaries and premaxillaries become almost horizontal (Fig. 1B). The expansion of the buccal cavity is confirmed by the hyoid depression in comparison with the neurocranium top, but not with the bottom (elevation).

Observed on dead animals, the lower pharyngeal jaws are located forward the upper jaws in extended position (Fig. 1A). In curved position, the upper and lower pharyngeal jaws appear opposite (Fig. 1B).

DISCUSSION

Teleosts have different systems for opening their mouths. Elevation of the neurocranium is one possible system; it is always caused by contraction of the epaxial musculature (Vandewalle, 1978; Liem, 1979; Sibbing, 1982; Westneat and Wainwright, 1989). Elevation of the neurocranium by about 10° during feeding has been described in many fish (Alexander, 1967; Liem, 1979; Lauder, 1979, 1980, 1981; Lauder and Liem, 1981; Sibbing, 1982; Muller, 1987; Westneat, 1994; Vandewalle *et al.*, 1995). This lifting does not seem to require special anatomical features. Some teleosts are capable of a greater elevation of the cephalic region: 35° for *Macrochirichthys macrochirus* (Valenciennes) (Howes, 1979), 40° for *Luciocephalus pulcher* (Gray) (Lauder and Liem, 1981), 50° for *Synanceia verrucosa* Bloch & Schneider (Grobeck, 1983), 40° for Syngnathidae (Bergert and Wainwright, 1997). Lesiuk and Lindsey (1978), Howes (1979) and Lauder and Liem (1981) have evidenced the specialised morphological features of certain teleosts capable of extreme markedly bending of the front of the vertebral column. In all cases the anterior vertebrae are modified, but in different manners. *Macrochirichthys macrochirus* raises its head simply by rotating around the first, reduced vertebra, fitting into the second vertebra (Howes, 1979). In the characoid *Raphiodon vulpinus* (Agassiz), bending can reach 45° and occurs posterior to the first four vertebrae, probably because of the presence of the Weberian apparatus. The fifth vertebra is wide, not very high, and bears a shorter neural spine than the following ones (Lesiuk and Lindsey, 1978). In *Uranoscopus scaber*, the changes in the vertebral column concern the first five vertebrae, which explains why the bending angle can be exceptionally great in this species (Goosse *et al.*, 1995). When the epaxial musculature, fixed on a small occipital crest and a wide occipital region (Piestch, 1989) contracts, vertebrae two to five, being not high, are lifted. This rotation is made possible by the interlocking of the first vertebral centra, the incline of the pre- and postzygapophyses, and the difference in width between the front and rear of the first neural arches, enabling them to cover each other to a certain degree. The structures limiting this rotation are probably the LI.5 ligaments and the neural spines. The subsequent vertebrae, because of their standard morphology, should not enable the vertebral column to curve differently from what is usually observed in teleosts (Lauder, 1979; Sibbing, 1982; Vandewalle *et al.*, 1995). Downward bending of the anterior region of the vertebral column should be minimal slight to impossible in *Uranoscopus scaber*, due to the presence of ligaments LI.1, LI.2, LI.3, and LI.4.



and associated ligaments; B: Lateral
Asterisks indicate where the ribs

ses. The fifth vertebra looks
um is still narrow and flatte-
: vertebra. The parapophyses
m. The neural arch is located
ded by a more tapered neural

ments. A large dorsal septum
vertebra; then it thickens and
ches to the rear of the neuro-
ture (wide and voluminous in

dorsal side of the prezygapo-
le more tilted than the follo-
ive vertebrae there is a liga-
mpletely masked between the
r surface of a prezygapophy-
t of the preceding vertebra.
ment LI.5 consisting of short
ear longer than those of the