


## The NeQuick Ionosphere Model

- **Ionosphere affecting** radio propagation and hence **GNSS**
  - Positioning errors exceeding 100 m in extreme cases
  - **Total Electron Content (TEC)**, integral of the electron density Ne = main driver
- Importance of **TEC modelling**
  - Crucial especially for **single frequency receivers**, the most common ones constituting the mass market
  - By means of a 3D method using the **NeQuick model for GALILEO** (Orus et al., 2007a)
- NeQuick = **empirical model** of the electron density Ne
  - “**Profiler**” = several mathematical functions fitted on anchor points corresponding to the maxima of the layers of the ionosphere (Radicella et Leitingner, 2001)
  - Peaks and profile characteristics calculated on the basis of **monthly median measurements**
  - **New version (NeQuick 2)**: main modification regarding the description of the **higher part of the ionosphere** (“topside”) → two formulas for shape parameter k (each for six months of the year) replaced by a single one (Nava et al., 2008)




## Tools and Method

- Investigation of **NeQuick profile formulation**
  - Monthly median measurements replaced by actual ones → **model constrained by means of ionosonde data** (Bidaine et Warnant, 2007)
  - Obtained vertical TEC compared to GPS TEC data → **collocated ionosonde and GPS receiver needed**
- Data types
  - **Manually validated digisonde data**
  - **Slant TEC data levelled using Global Ionospheric Maps** (Orus et al., 2007b) and mapped to vertical + elevation filter and average to obtain vertical TEC
- Tests for **mid-latitudes and high solar activity**
  - Year 2002
  - Three European locations with (nearly) collocated digisonde and IGS/EUREF station



## Yearly Statistics



- Influence of **latitude**: vertical TEC mean decreasing northwards (TEC in TECu =  $10^{16} e^{-m^{-2}}$ )
- TEC **underestimated** on average (potential bias in GPS TEC data to take into account)
- **Bigger** (around 20%) **underestimation** with NeQuick 2
- **Lower** (around 20%) **standard deviation** for NeQuick 2 → better behaviour


$$dTEC = TEC_{meas} - TEC_{mod} \quad Bias = \langle dTEC \rangle$$

$$Standard\ deviation = \sqrt{\langle (dTEC - Bias)^2 \rangle}$$

$$Relative\ parameter = \frac{Parameter}{\langle TEC_{meas} \rangle}$$

$$Evolution = \frac{Parameter_{NeQuick\ 2} - Parameter_{NeQuick\ 1}}{Parameter_{NeQuick\ 1}}$$


## Influence of k Unification



- **Major modification between model versions**: 2 formulas for k (April to September and October to March) in NeQuick 1 replaced by a single one in NeQuick 2
- Apparently **different behaviour for periods corresponding to both k formulas** in NeQuick 1
- April to September: lower (20%) bias and slightly higher standard deviation in NeQuick 2
- **October to March**:
  - In NeQuick 1, lower bias and higher standard deviation than first period
  - In NeQuick 2, much higher (200%) bias and lower (15%) standard deviation → **more homogenous with first period**


## TEC Splitting

- **Integrate bottomside Ne profile** from digisonde to obtain bottomside TEC
- **Subtract bottomside TEC to GPS TEC** to obtain estimate of topside TEC (caution with interpretation about topside because resulting TEC value containing whole GPS TEC uncertainty)
- **Big proportion of TEC within topside** (3/4, 1/4)
- Bottomside: low bias (at least in absolute value) and relatively high standard deviation, no big evolution between NeQuick versions
- **Topside**: higher relative bias and standard deviation than bottomside, **bias/standard deviation evolution between NeQuick versions corresponding to global statistics**



## Conclusion and Perspectives

- **Investigation of NeQuick profile formulation** for mid-latitudes and high solar activity
  - Constraining NeQuick with ionosonde measurements and comparing resulting vertical TEC with GPS TEC
  - **Standard deviation decreasing by 20% to reach less than 20% with NeQuick 2**, bias increasing by 20% up to 25% but caution with GPS TEC data
  - Homogenisation thanks to topside modification
  - **Major role of the topside**
- **Further research**
  - **Ingestion**: adapt NeQuick TEC to GPS TEC by means of effective parameters
  - Investigate **GALILEO Single Frequency Ionospheric Correction Algorithm**



## Acknowledgements

- Research undertaken under F.R.S.-FNRS fellowship
- Sandro Radicella, Pierdavide Coisson and Bruno Nava from ICTP in Trieste for providing NeQuick latest version and comments about it
- Iñigo Blanco Alegre from INTA (El Arenosillo), David Altadill from Observatori de l'Ebre (Roquetes), Elise Van Malderen and Luc Lejeune from RMI (Dourbes) for providing ionosonde data and comments about them
- Grigori Khmyrov and Bodo Reinisch from UMLCAR in Boston for providing access to the DIDBase (digisonde database)
- Roberto Prieto Cerdeira and Raul Orus from ESA/ESTEC for providing TEC data and comments about them

## References

- BIDAINE, B. and R. WARNANT (2007): Assessment of the NeQuick Model at Mid-latitudes Using GPS TEC and Ionosonde Data, *Proc. 1st Colloquium Scientific and Fundamental Aspects of the Galileo Programme*.
- NAVA, B., P. COISSON and S.M. RADICELLA (2008): A new version of the NeQuick ionosphere electron density model, *J. Atmos. and Sol.-Terr. Phys.*, in press, corrected proof. doi:10.1016/j.jastp.2008.01.015
- ORUS, R., B. ARBESSER-RASTBURG, R. PRIETO-CERDEIRA, M. HERNANDEZ-PAJARES, J. M. JUAN and J. SANZ (2007a): Performance of Different Ionospheric Models for Single Frequency Navigation Receivers, *Proc. Beacon Satellite Symposium 2007*.
- ORUS, R., Lj. R. CANDER and M. HERNANDEZ-PAJARES (2007b): Testing regional vTEC maps over Europe during the 17-21 January 2005 sudden space weather event, *Radio Sc.*, Vol. 42, RS3004. doi:10.1029/2006RS003515
- RADICELLA, S. M., and R. LEITINGER (2001): The evolution of the DGR approach to model electron density profiles, *Adv. Space Res.*, Vol. 27, No. 1, p. 35-40. doi:10.1016/S0273-1177(00)00138-1