

Narrow-band photometry of the eclipsing WN7+O binary WR 22

Pierre Royer¹, Gregor Rauw¹, Jean Manfroid¹, Eric Gosset¹, and
Jean-Marie Vreux

*Institut d'Astrophysique et de Géophysique, Université de Liège,
5, Avenue de Cointe, B-4000 Liège, Belgium*

Abstract. In the present paper, we discuss photometric observations of the February 1996 eclipse of the very massive WR+O binary WR 22. Our data were obtained with a set of narrow-band filters, specially designed for the study of WR stars.

1. Introduction

The WN7+abs+O binary WR 22 ($P_{\text{orb}} = 80.3$ d) displays a shallow eclipse around its periastron passage, when the WR star is in front of its companion (Balona *et al.* 1989; Gosset *et al.* 1991). Such an atmospheric eclipse results from the phase-dependent opacity along our sightline towards the O star and is a common feature among WR+O binaries with orbital periods shorter than 30 days (Lamontagne *et al.* 1996).

In the case of WR 22, as a result of the combination of the inclination of the orbit, of its eccentricity ($e \simeq 0.56$), and of the argument of the periastron ($\omega \simeq 272^\circ$), only the secondary eclipse near periastron is present, whereas no occultation is observed near apastron (Gosset *et al.* 1991; Rauw *et al.* 1996).

2. Narrow-band photometry of WR 22

Photometric observations, through specially designed narrow-band filters, of the secondary eclipse of WR 22 were acquired during five nights in February 1996 (JD 2450125–129) at the Bochum 0.6 m telescope (La Silla, Chile) equipped with the DLR CCD camera using a Tektronix 1024×1024 chip (pixel size $23 \mu\text{m}$). For a detailed description of the characteristics and performances of our set of narrow-band filters, see Royer *et al.* (1998). The data were reduced using the algorithm described by Manfroid (1993).

Given the incomplete phase-coverage of the eclipse by the narrow-band observations, we have fitted the data with the mean light-curve, as derived from a more extensive data set obtained in the Strömgren y filter, keeping the depth of the eclipse as the only free parameter (see Fig. 1). The Strömgren data set will be discussed and analysed in a forthcoming paper (Rauw *et al.* in preparation).

¹At the Fonds National de la Recherche Scientifique, Belgium




Figure 1. Light-curve of WR 22 as observed with a filter centered on 6051 Å. Differential magnitudes are shown as a function of $\theta = \psi - \frac{3\pi}{2} + \omega$ where ψ is the true anomaly and ω the argument of the periastron (orbital elements from Rauw *et al.* 1996).

As expected, the eclipse as observed with the filter centered on the He II λ 4686 emission line is shallower than with the continuum filters (centered on λ 5057 and λ 6051). The fitted depths are $\Delta m(\lambda 4686) = 0.062$, $\Delta m(\lambda 5057) = 0.112$ and $\Delta m(\lambda 6051) = 0.102$. The depths of the eclipse in the continuum filters provide upper limits on the luminosity ratio L_{WR}/L_O of 9.2 (λ 5057) and 10.1 (λ 6051). We have performed numerical simulations to quantify the contribution of the He II λ 4686 line to the integrated flux in the narrow filter as a function of the line's orbital Doppler shift. On the average, the emission line contributes for some 42% of the total flux and hence accounts for the depth difference of the eclipse as observed with the He II λ 4686 filter compared to the continuum filters. During the phase interval $\phi = 0.97 \rightarrow 0.03$, the radial velocity of the He II λ 4686 line increases from ~ 0 to $\sim +120 \text{ km s}^{-1}$. Due to the Doppler shift of the line within the narrow filter passband, our simulations, accordingly, yield an apparent faintening by 0.007 magnitude during this phase interval, *i.e.*, too small an effect to be detected in our data given their accuracy and the intrinsic variability of the WN7+abs star. In contrast to most of the light-curves of eclipsing WR+O systems reported by Lamontagne *et al.* (1996), the light-curve of WR 22 displays a rather *narrow shape without prominent extended wings*, pointing towards a rather low mass-loss rate ($\log \dot{M} \leq -4.8$, Rauw 1997).

References

Balona, L.A., Egan, J., Marang, F. 1989, MNRAS 240, 103
 Gosset, E., Remy, M., Manfroid, J., *et al.* 1991, IBVS 3571
 Lamontagne, R., Moffat, A.F.J., Drissen, L., Robert, C., Matthews, J.M. 1996, AJ 112, 2227
 Manfroid, J. 1993, A&A 271, 714
 Rauw, G. 1997, PhD Thesis, Université de Liège
 Rauw, G., Vreux, J.-M., Gosset, E., *et al.* 1996, A&A 306, 771
 Royer, P., Vreux, J.-M., Manfroid, J. 1998, A&AS 130, 407