

Reintroduction of the invasive mosquito species *Aedes albopictus* in Belgium in July 2013

Slimane Boukraa^{1,*}, Fara N. Raharimalala^{1,2}, Jean-Yves Zimmer¹, Francis Schaffner³, Thomas Bawin¹, Eric Haubruge¹, and Frédéric Francis¹

¹ Gembloux Agro-Bio Tech, University of Liège, 5030 Gembloux, Belgium

² Medical Entomology Unit, Pasteur Institute, Antananarivo, Madagascar

³ National Centre for Vector Entomology, Institute of Parasitology, University of Zurich, Switzerland

Received 12 November 2013, Accepted 3 December 2013, Published online 12 December 2013

Abstract – Since its first report in 2000, the invasive mosquito *Aedes albopictus* was not found any more during the different entomological inspections performed at its place of introduction in Belgium between 2001 and 2012. In July 2013, one adult male was captured at the same site (a platform of imported used tires located in Vrasene, Oost-Vlaanderen Province), during a monitoring using CO₂-baited trap. This finding suggests the reintroduction of the species in Belgium via the used tire trade.

Key words: *Aedes albopictus*, Belgium, Used tire.

Résumé – Réintroduction du moustique invasif *Aedes albopictus* en Belgique en Juillet 2013. Depuis sa première observation en 2000, le moustique invasif *Aedes albopictus* n'a plus été trouvé au cours des différentes inspections entomologiques réalisées en Belgique sur le site d'introduction entre 2001 et 2012. En Juillet 2013, un adulte mâle a été capturé sur ce même site (une plateforme de pneus usagés importés située à Vrasene, Province de Flandre-Orientale), durant une surveillance ayant recours à un piège à CO₂. Cette redécouverte suggère la réintroduction de cette espèce en Belgique via le commerce de pneus usagés.

Introduction

Emerging arbovirogenesis occurrences are related to a change in pathogens and/or arthropod vector distributions [10]. Environment and climate change, as well as globalization of international trade can affect these distributions [6]. Since the late 1970s, the Tiger mosquito *Aedes (Stegomyia) albopictus* (Skuse, 1894) [14] shows an explosive worldwide spreading, being currently the most invasive mosquito in the world. In Europe, it has been reported from 20 countries and is nowadays well established in the Mediterranean region [8]. *Aedes albopictus* belongs to the most important arbovirus vectors, in particular for chikungunya and dengue viruses [8]. The risk for emergence and spread of these arboviruses to no-epidemic regions has increased especially in regions where *Ae. albopictus* has established, as demonstrated by recent local transmissions of chikungunya and dengue in Croatia and France [8]. In Belgium, *Ae. albopictus* was reported for the first time in 2000 from Vrasene (Oost-Vlaanderen Province), on an used tire storage of a recycling company that imports from the USA and

Japan, among other countries [11]. Several other inspections were performed after that report but no additional specimens have been found in Belgium, although the site where it has been reported from has been monitored [15, F. Schaffner unpublished data]. Thus, the species is considered as not established in Belgium and, as no control measures have been applied, it is assumed that the introduced population was naturally eliminated. Besides, there is no evidence of any additional introduction at other points of entries [15]. During this last decade, two other Asian mosquitoes were recorded in Belgium. *Ochlerotatus japonicus japonicus* (Theobald, 1901) was introduced and has established in southern Belgium (Hamois, Namur province) and *Oc. koreicus* (Edwards, 1917) was collected in Eastern Belgium (Maasmechelen, Limburg province), where it successfully established as well [15].

Material and methods

In the frame of a study of bacterial endosymbionts in Belgian mosquitoes, adult mosquitoes were regularly trapped at several places throughout the country by CO₂-baited traps Mosquito Magnet Liberty Plus® (MMLP) and collected by sweep

*Corresponding author: entomologie.gembloux@ulg.ac.be

Table 1. Adult and larval mosquitoes collected in Oost-Vlaanderen province, Belgium (02-07 VII 2013).

Site	Collection method / place	<i>Anopheles plumbeus</i>	<i>Aedes albopictus</i>	<i>Culex pipiens s.l.</i>	<i>Culex torrentium</i> *	<i>Coquillettidia richiardii</i>	<i>Ochlerotatus geniculatus</i>	<i>Ochlerotatus dorsalis</i>
Vrasene	MMLP / TSOG	458 F	1 M	10 F	0	1 F	61 F, 20 M	0
	Sweep net / TSOG	28 F	0	0	0	0	6 F	0
	Dipper / TSS	0	0	503 L	5 L	0	36 L	0
	Dipper / TSOG	428 L	0	638 L	1925 L	0	0	0
Lochristi	MMLP / TSI	0	0	65 F	0	0	0	1 M
	Dipper / TSS	75 L	0	1424 L	0	0	0	0

L: larva; M: male; F: female; MMLP: Mosquito Magnet Liberty Plus; TSS: Tires stored in shelters; TSI: Tires stored inside; TSOG: Tires stored outside, bordering grassland; *Identification of this species based on male genitalia after emerging in laboratory.

netting, and immatures were collected by the dipping method. For the present study, the surveys were conducted in two storage center of used tires recycling companies located in Oost-Vlaanderen Province. Study site A was at Vrasene (51°12'49" N, 4°11'37" E; 5 m above sea level) and at less of 10 km from port of Antwerp. This platform of imported used tires was regularly inspected following the find of *Ae. albopictus* in 2000 by Schaffner et al. [11]. Study site B is a company for recycling tires of local origin, located in Lochristi (51°06'18" N, 3°52'12" E; 5 m above sea level) and approximately 20 km southwest of the first study site. Identification of *Ae. albopictus* was first performed by morphology and then confirmed by molecular tools. The mitochondrial cytochrome oxidase subunit I (COI) was amplified [9] using the primers CI-J-1632 and CI-N-2191 [7] and sequenced. The nucleotide sequence is deposited in GenBank under accession number [KF657725](https://www.ncbi.nlm.nih.gov/nuclot/KF657725). Blast analysis was used to compare the obtained COI sequence with data available in NCBI (www.ncbi.nlm.nih.gov).

Results and discussion

Early July 2013, one adult male of *Ae. albopictus* was trapped (MMLP) in Vrasene at the same used tire recycling company where it was observed in 2000. The obtained sequence compared with data available in NCBI showed 99% of similarity with those of *Ae. albopictus* from the United States and Germany. Other species collected together with *Ae. albopictus* were *Anopheles plumbeus* (Stephens, 1828), *Culex pipiens* (Linnaeus, 1758) s.l., *Cx. torrentium* (Martini, 1925), *Coquillettidia richiardii* (Ficalbi, 1899), and *Oc. geniculatus* (Olivier, 1791) (Table 1).

This rediscovery of *Ae. albopictus* together with the absence of any finding during previous years (2001–2012) suggests its reintroduction into Belgium via the used tire trade. In addition, given the information about the origin of used tires recently imported by the company, the source of the species reintroduction is possibly the United States. *Aedes albopictus* is a confirmed efficient vector of Dengue and Chikungunya viruses [8], and Belgium regularly registers imported chikungunya cases [2]. Thus, if established in the country, *Ae. albopictus* may become a substantial threat to public health. A study of the survival and dispersal of this mosquito in Belgium, as well as of its bioecology in neighboring countries might provide important insights to further elucidate its invasiveness, and identify

high-risk areas for mosquito proliferation and pathogen transmission. A rapid proactive response is critical for vector management because of the possibility of its establishment, according to several models [3, 4]. This includes rapid implementation of control measures, before elimination is impossible [12]. Several countries in Western Europe recently confirmed repeated introductions of *Ae. albopictus* [1, 5, 13], and thus we suggest broader and more thorough entomological surveys at the European scale to survey introduction pathways and prevent establishment, and subsequently to reduce the risks of future arbovirus transmission.

Acknowledgements. We thank both tire company managers for allowing unrestricted access to their sites. We also thank Sophie Vandermotten (Gembloux Agro-Bio Tech, ULg) for help with the molecular study. This work was supported by the University of Liège (ULg) and Wallonie-Bruxelles International (WBI).

References

1. Becker N, Geier M, Balczun C, Bradersen U, Huber K, Kiel E, Krüger A, Lühken R, Orendt C, Plenge-Bönig A, Rose A, Schaub GA, Tannich E. 2013. Repeated introduction of *Aedes albopictus* into Germany, July to October 2012. *Parasitology Research*, 112, 1787–1790.
2. Bottieau E, Van Esbroeck M, Cnops L, Clerinx J, Van Gompel A. 2009. Chikungunya infection confirmed in a Belgian traveller returning from Phuket (Thailand). *Euro Surveillance*, 14, pii = 19248. Available online: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19248> (accessed 11 August 2013).
3. Caminade C, Medlock JM, Ducheyne E, McIntire KM, Leach S, Baylis M, Morse AP. 2012. Suitability of European climate for the Asian tiger mosquito *Aedes albopictus*: recent trends and future scenarios. *Journal of the Royal Society Interface*, 9, 2708–2717.
4. European Centre for Disease Prevention and Control (ECDC). 2009. Development of *Aedes albopictus* risk maps. ECDC Technical Report. Stockholm: ECDC. Available online: http://ecdc.europa.eu/en/publications/Publications/0905_TER_Development_of_Aedes_Alboipictus_Risk_Maps.pdf (accessed 10 August 2013).
5. Expatica, DutchNews.nl. *Aedes albopictus* – Netherlands. ProMed2013 Aug 17, <http://www.promedmail.org/direct.php?id=20130817.1885955> (accessed 28 August 2013).
6. Gould EA, Higgs S. 2009. Impact of climate change and other factors on emerging arbovirus diseases. *Transaction of the Royal Society of Tropical Medicine and Hygiene*, 103, 109–121.

7. Kambhampati S, Smith PT. 1995. PCR primers for the amplification of four insect mitochondrial gene fragments. *Insect Molecular Biology*, 4, 233–236.
8. Medlock JM, Hansford KM, Schaffner F, Versteirt V, Hendrickx G, Zeller H, Van Bortel W. 2012. A review of the invasive mosquitoes in Europe: Ecology, public health risks, and control options. *Vector Borne Zoonotic Diseases*, 12, 435–447.
9. Mousson L, Dauga C, Garrigues T, Schaffner F, Vazeille M, Failloux AB. 2005. Phylogeography of *Aedes (Stegomyia) aegypti* (L.) and *Aedes (Stegomyia) albopictus* (Skuse) (Diptera: Culicidae) based on mitochondrial DNA variations. *Genetics Research*, 86, 1–11.
10. Randolph SE, Rogers DJ. 2010. The arrival, establishment and spread of exotic diseases: patterns and predictions. *Nature Reviews Microbiology*, 8, 361–371.
11. Schaffner F, Van Bortel W, Coosemans M. 2004. First record of *Aedes (Stegomyia) albopictus* in Belgium. *Journal of the American Mosquito Control Association*, 20, 201–203.
12. Schaffner F, Bellini R, Petrić D, Scholte E-J, Zeller H, Marrama Rakotoarivony L. 2013. Development of guidelines for the surveillance of invasive mosquitoes in Europe. *Parasites and Vectors*, 6, 209.
13. Scholte EJ, Dijkstra E, Blok H, De Vries A, Takken W, Hofhuis A, Koopmans M, De Boer A, Reusken CB. 2008. Accidental importation of the mosquito *Aedes albopictus* into the Netherlands: a survey of mosquito distribution and the presence of dengue virus. *Medical and Veterinary Entomology*, 22, 352–358.
14. Skuse F. 1894. The banded mosquito of Bengal, *Indian Museum Notes*, 3, 20.
15. Versteirt V, Boyer S, Damiens D, De Clercq EM, Dekoninck W, Ducheyne E, Grootaert P, Garros C, Hance T, Hendrickx G, Coosemans M, Van Bortel W. 2012. Nationwide inventory of mosquito biodiversity (Diptera: Culicidae) in Belgium, Europe. *Bulletin of Entomological Research*, 103, 193–203.

Cite this article as: Boukraa S, Raharimalala FN, Zimmer J-Y, Schaffner F, Bawin T, Haubruge E & Francis F: Reintroduction of the invasive mosquito species *Aedes albopictus* in Belgium in July 2013. *Parasite*, 2013, 20, 54.



An international open-access, peer-reviewed, online journal publishing high quality papers on all aspects of human and animal parasitology

Reviews, articles and short notes may be submitted. Fields include, but are not limited to: general, medical and veterinary parasitology; morphology, including ultrastructure; parasite systematics, including entomology, acarology, helminthology and protistology, and molecular analyses; molecular biology and biochemistry; immunology of parasitic diseases; host-parasite relationships; ecology and life history of parasites; epidemiology; therapeutics; new diagnostic tools.

All papers in Parasite are published in English. Manuscripts should have a broad interest and must not have been published or submitted elsewhere. No limit is imposed on the length of manuscripts.

Parasite (open-access) continues **Parasite** (print and online editions, 1994-2012) and **Annales de Parasitologie Humaine et Comparée** (1923-1993) and is the official journal of the Société Française de Parasitologie.

Editor-in-Chief:
Jean-Lou Justine, Paris

Submit your manuscript at
<http://parasite.edmgr.com/>