

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

Photoferrotrophy and Fe-cycling in a freshwater column

LLIRÓS M^{1,2}; CROWE SA³; GARCÍA-ARMISEN T⁴;
DARCHAMBEAU F⁵; MORANA C⁶; BORREGO CM^{7,8};
TRIADÓ-MARGARIT X⁹; BOUILLON S⁶; BORGES AV⁵;
SERVAIS P⁴; CANFIELD D³; AND DESCY JP¹

¹ Laboratory of Freshwater Ecology – URBE, Department of Biology, University of Namur, B-5000 Namur, Belgium. marc.lliros@uab.cat; jpdescy@fundp.ac.be

² Department of Genetics and Microbiology, Universitat Autònoma de Barcelona, 08913 Bellaterra, Spain. marc.lliros@uab.cat

³ Institute of Biology, Nordic Center for Earth Evolution, University of Southern Denmark, 5230 Odense, Denmark. sacrowe1@gmail.com; dec@biology.sdu.dk

⁴ Ecologie des Systèmes Aquatiques – ESA, Université Libre de Bruxelles, B-1050 Brussels, Belgium. tgarciaa@ulb.ac.be; pservais@ulb.ac.be

⁵ Chemical Oceanography Unit, Université de Liège, B-4000 Liège, Belgium. Francois.Darchambeau@ulg.ac.be; alberto.borges@ulg.ac.be

⁶ Department of Earth & Environmental Sciences, Katholieke Universiteit Leuven, B-3001 Leuven, Belgium. Cedric.Morana@ees.kuleuven.be; Steven.Bouillon@ees.kuleuven.be

⁷ Group of Molecular Microbial Ecology, Institute of Aquatic Ecology, University of Girona, E-17071 Girona, Spain. carles.borrego@udg.edu

⁸ Water Quality and Microbial Diversity, Catalan Institute for Water Research (ICRA), E-17003, Girona, Spain. carles.borrego@udg.edu

⁹ Limnological Observatory of the Pyrenees (LOOP) – Biogeodynamics & Biodiversity Group, Department of Continental Ecology, Centre d'Estudis Avançats de Blanes, CEAB–CSIC, Blanes, Spain. xtriado@ceab.csic.es

Emerging insight shows that ferruginous (anoxic and iron-rich) conditions dominated ocean chemistry throughout the first 3.5 billion years of Earth evolution. Modern ferruginous water masses are rare, but detailed examination of these oddities, especially of photoferrotrophs and their Fe-reducing respiratory counterparts, could yield important insights into the early evolution of life on Earth and its impact on global element cycles.

Here, we describe an abundant microbial community of pelagic photoferrotrophic and Fe-reducing microbes in a freshwater basin in East Africa (Kabuno Bay, DR Congo), recently discovered to be ferruginous. The application of

This abstract is too long to be accepted for publication in Mineralogical Magazine. Please revise it so that it fits into the column on one page.

culture-dependent and -independent techniques allowed the identification of active photoferrotrophs genetically similar to laboratory cultures of *Chlorobium ferrooxidans*, the only member of the *Chlorobi* previously known to conduct photoferrotrophy. The Kabuno Bay photoferrotrophic Green Sulfur Bacteria (GSB) community ranged between the 16.2% and 38.3% of the entire bacterial community retrieved by 454 pyrotag analyses of the euphotic Fe-rich chemocline of Kabuno bay. The GSB maximum coincided with high rates of ferrous Fe oxidation and bacterial production. Up to 60% of total depth integrated BP is carried out at the depths where photoferrotrophic GSB dominate. Main bacterial groups involved in methanogenesis were also recovered from the upper oxic-anoxic transition zone, but in contrast to the GSB community, other microorganisms typically implicated in the sulfur cycle were only present at low relative abundances. DAPI counts revealed a high microbial biomass in the water column of Kabuno bay, which we speculate supports large populations of pelagic iron-reducers. Reactive iron oxides, however, are exhausted within the chemocline and excess organic matter is thus channelled through mineralization by methanogenesis directly in the water column.

Our study documents for the first time that photoferrotrophs are suited to pelagic lifestyles and putatively possess a low-light adapted light harvesting apparatus. That pelagic carbon mineralization is channelled through methanogenesis even under ferruginous conditions implying that gaseous reduced equivalents can escape to the atmosphere, allowing excess ferric iron to be exported to sediments such as Banded Iron Formations. In this way, photoferrotrophs could contribute to the overall oxidation of the Earth's surface prior to the oxygenation of the atmosphere.