

New cycles found in air temperature data and proxy series using the Morlet wavelet method

S. Nicolay¹, G. Mabille², X. Fettweis², M. Erpicum²

¹Department of Mathematics, University of Liège,

²Department of Physical Geography, University of Liège.

January 12, 2009

The Continuous Wavelet transform

The wavelet analysis provides a two-dimensional unfolding of a one-dimensional signal by decomposing it into scale–time coefficients.

The continuous wavelet transform turns a signal s into a function W

$$W[s](t, a) = \int s(x) \bar{\psi}\left(\frac{x-t}{a}\right) \frac{dx}{a},$$

where $\bar{\psi}$ denotes the complex conjugate of the function ψ , a the scale and t the time.

Conditions on ψ

The function ψ must be integrable, square integrable and satisfy some admissibility condition. Such a function is called a wavelet. The Morlet wavelet is particularly well conditioned for frequency-based study. It satisfies the following equality

$$\hat{\psi}(\omega) = \exp\left(-\frac{(\omega - \Omega)^2}{2}\right) - \exp\left(-\frac{\omega^2 + \Omega^2}{2}\right),$$

where $\Omega > 5$ is called the central frequency.



Some Properties of the Wavelet Transform

- the wavelet transform is linear:
$$W[c(u + v)] = cW[s] + cW[v],$$

Some Properties of the Wavelet Transform

- the wavelet transform is linear:
$$W[c(u + v)] = cW[u] + cW[v],$$
- the wavelet transform allows to handle noisy data,

Some Properties of the Wavelet Transform

- the wavelet transform is linear:
$$W[c(u + v)] = cW[s] + cW[v],$$
- the wavelet transform allows to handle noisy data,
- the wavelet transform is blind to polynomial behaviors (up to a degree n , depending on ψ): $W[s + P] = W[s]$, where P is a polynomial of degree $\leq n$.

Consequently, non-zero mean and linear tendencies do not affect the wavelet transform.

The Scale Spectrum

For wavelets such as the Morlet wavelet, we have

$$W[\cos(\omega_0 t)](t, a) = \exp(i\omega_0 t) \hat{\psi}(a\omega_0),$$

so that the frequency ω_0 is given by the maximum of $\hat{\psi}(a\omega_0)$:
 $a_\omega = \Omega/\omega_0$. Consequently, the unknown frequency ω_0 can be obtained through the maximum of $|W[\cos(\omega_0 t)]|$.

The Scale Spectrum

For wavelets such as the Morlet wavelet, we have

$$W[\cos(\omega_0 t)](t, a) = \exp(i\omega_0 t) \hat{\psi}(a\omega_0),$$

so that the frequency ω_0 is given by the maximum of $\hat{\psi}(a\omega_0)$:
 $a_\omega = \Omega/\omega_0$. Consequently, the unknown frequency ω_0 can be obtained through the maximum of $|W[\cos(\omega_0 t)]|$.

Definition

The scale spectrum of a signal s is defined by

$$\Lambda(a) = E|W[s](t, a)|,$$

where E denotes the mean over the time t .


For What Purpose?

The scale spectrum should be useful for signals which are not stationary but whose characteristics do not evolve too quickly: the scale spectrum allows to recover frequencies even if they “kindly depend on the time”, i.e. if $\omega_0 = \omega_0(t)$ with $\frac{d}{dt}\omega_0 \ll 1$, one should be able to recover $E\omega_0$.




Example

A Visual Example




Example

A Visual Example




Example

A Visual Example



Example


Known long cycles are observed in the spectrum



EPICA dome

Example


Known long cycles are observed in the spectrum



EPICA dome

Example


Known long cycles are observed in the spectrum



EPICA dome

Known long cycles are observed in the spectrum

Milankovitch cycles are detected, as expected



EPICA dome


A weather station

The time series we are interested in...



Global temperature data

What About the Global Temperature Data?




HadCRUT3, NH

1950–2007 monthly-sampled data

Global temperature data

What About the Global Temperature Data?




HadCRUT3, NH

1950–2007 monthly-sampled data

Global temperature data

What About the Global Temperature Data?




HadCRUT3, NH

1950–2007 monthly-sampled data


Global temperature data

What About the Global Temperature Data?



HadCRUT3, NH

1950-2007 monthly-sampled data




HadCRUT3, SH

1950-2007 monthly-sampled data


Global temperature data

What About the Global Temperature Data?



HadCRUT3, NH

1950–2007 monthly-sampled data




HadCRUT3, SH

1950–2007 monthly-sampled data


Global temperature data

What About the Global Temperature Data?



HadCRUT3, NH

1950-2007 monthly-sampled data




HadCRUT3, SH

1950-2007 monthly-sampled data

Global temperature data

What About the Global Temperature Data?




HaDSST2

1950–2007 monthly-sampled data

Global temperature data

What About the Global Temperature Data?




HaDSST2

1950–2007 monthly-sampled data


Global temperature data

What About the Global Temperature Data?



HaDSST2

1950–2007 monthly-sampled data




CRUTEM3gl

1950–2007 monthly-sampled data


Global temperature data

What About the Global Temperature Data?



HaDSST2

1950–2007 monthly-sampled data




CRUTEM3gl

1950–2007 monthly-sampled data


Global temperature data

What About the Global Temperature Data?



HaDSST2

1950–2007 monthly-sampled data




CRUTEM3gl

1950–2007 monthly-sampled data


Global temperature data

What About the Global Temperature Data?



HaDSST2

1950–2007 monthly-sampled data




CRUTEM3gl

1950–2007 monthly-sampled data

Global temperature data


What About the Local Temperature Data?



Bierset-aero, Belgium
1950–2007 hourly-sampled data

Global temperature data


What About the Local Temperature Data?



Bierset-aero, Belgium
1950–2007 hourly-sampled data

Global temperature data


What About the Local Temperature Data?



Bierset-aero, Belgium
1950–2007 hourly-sampled data

Global temperature data


What About the Local Temperature Data?




Bierset-aero, Belgium
1950–2007 hourly-sampled data

Global temperature data

What About the Local Temperature Data?




Bierset-aero, Belgium
1950–2007 hourly-sampled data




The Pas, Canada
1950–2007 monthly-sampled data

Global temperature data

What About the Local Temperature Data?




Bierset-aero, Belgium
1950–2007 hourly-sampled data




The Pas, Canada
1950–2007 monthly-sampled data

Global temperature data

What About the Local Temperature Data?



Bierset-aero, Belgium
1950–2007 hourly-sampled data



The Pas, Canada
1950–2007 monthly-sampled data

A Random Process to Compare the Data With

One of the methods to check if “This cycle which is present in this time series did not just occur by pure chance” consists in comparing the scale spectra to spectra obtained from a random process “looking like” the data.



A Random Process to Compare the Data With


One of the methods to check if “This cycle which is present in this time series did not just occur by pure chance” consists in comparing the scale spectra to spectra obtained from a random process “looking like” the data.

A good model seems to be the AR(1):

$$x_{n+1} = \alpha x_n + \sigma \eta_{n+1},$$


where , as usual, η_n is i.i.d., with zero mean and unit variance (e.g. Gaussian white noise $N(0,1)$).

A Visual Illustration



Wien, Austria


A Visual Illustration




Wien, Austria / A simulation

A Model for the Data



The scale spectra of the Wien Temperature Records vs. the Mean Spectra of the Model




30 Months Cycle




NCEP/NCAR Reanalysis Monthly Means




42 Months Cycle



NCEP/NCAR Reanalysis Monthly Means




Scale Spectrum of Some Climatic Indices



NAO (CPC)

1950–2007 monthly-sampled data


Scale Spectrum of Some Climatic Indices



NAO (CPC)


1950–2007 monthly-sampled data

Scale Spectrum of Some Climatic Indices



NAO (CPC)


1950–2007 monthly-sampled data



ENSO (SST)


1950–2007 monthly-sampled data

Scale Spectrum of Some Climatic Indices



NAO (CPC)


1950–2007 monthly-sampled data




ENSO (SST)

1950–2007 monthly-sampled data

Millennial Temperature Reconstructions of Jones et al.





Years
Northern hemisphere





Years
Southern hemisphere



Millennial Temperature Reconstructions of Jones et al.




Millennial Temperature Reconstructions of Jones et al.



Millennial Temperature Reconstructions of Jones et al.




Scale spectrum of the Sunspot Number




1950–2007 monthly-sampled data

Scale spectrum of the Sunspot Number



1950–2007 monthly-sampled data


Scale spectrum of the Sunspot Number



1950–2007 monthly-sampled data


What About the Sun

A Relation with the Sun?



What About the Sun?

A Relation with the Sun?



For Further Reading



S. Nicolay, G. Mabilie, X. Fettweis and M. Erpicum,
30 and 43 months period cycles found in air temperature time series using the
Morlet wavelet method,
Climate Dynamics, doi: [10.1007/s00382-008-0484-5](https://doi.org/10.1007/s00382-008-0484-5) (2008)



G. Mabilie and S. Nicolay,
New cycles found in air temperature data and proxy series,
submitted



http://reflexions.ulg.ac.be/cms/c_5030/mathematiques