



Transport fluxes and emission of greenhouse gases of the Middle Niger River (west Africa): disproportionate importance of the recent red floods in the Niamey region

François Darchambeau (1), Steven Bouillon (2), Bassirou Alhou (3), Thibault Lambert (1), and Alberto V. Borges (1)

(1) University of Liège, Chemical Oceanography Unit, Belgium (francois.darchambeau@ulg.ac.be), (2) Katholieke Universiteit Leuven, Department of Earth and Environmental Sciences, Leuven, Belgium, (3) Université Abdou Moumouni, Ecole Normale Supérieure, Département des Sciences de la Vie et de la Terre, Niamey, Niger

The Niger River is Africa's third longest river and drains an area of $\sim 2,120,000$ km². It encompasses six hydrographic regions and crosses almost all possible ecosystem zones in West Africa. Since few decades, the Middle Niger River presents a two flood hydrograph, the local flood, or red flood, occurring during the rainy season being the more pronounced one. Here, we report initial results of a monitoring campaign whereby 2-weekly samples were collected at Niamey (Niger) [2.01°E 13.57°N] between April 2011 and March 2013 for a suite of physico-chemical and biogeochemical characteristics, including total suspended matter (TSM) concentrations, concentration and stable isotope composition of particulate organic carbon (POC and $\delta^{13}\text{C}$ -POC) and particulate nitrogen (PN and $\delta^{15}\text{N}$ -PN), chromophoric dissolved organic matter (CDOM), dissolved organic carbon (DOC and $\delta^{13}\text{C}$ -DOC), dissolved inorganic carbon (DIC and $\delta^{13}\text{C}$ -DIC), concentration of greenhouse gases (GHGs) (CO₂, CH₄ and N₂O), as well as major elements, total alkalinity, and oxygen isotope signatures of water ($\delta^{18}\text{O}$ -H₂O). This dataset allows us to construct seasonal budgets for particulate and dissolved carbon fluxes, nutrient exports, as well as a first seasonally resolved characterisation of the GHGs emitted to the atmosphere by the Middle Niger River. The red flood, concentrated on 2 months (August-September), contributed to more than 80% of the annual transport fluxes of TSM and POC and to approximately 30% of the annual transport fluxes of DIC and DOC.