

## Neuronal and Psychological Underpinnings of Pathological Gambling

Bryan Frederick Singer, Patrick Anselme, Mike James Ferrar Robinson and Paul Vezina

Journal Name:	Frontiers in Behavioral Neuroscience
ISSN:	1662-5153
Article type:	General Commentary Article
Received on:	29 May 2014
Accepted on:	06 Jun 2014
Provisional PDF published on:	06 Jun 2014
www.frontiersin.org:	<a href="http://www.frontiersin.org">www.frontiersin.org</a>
Citation:	Singer BF, Anselme P, Robinson MJ and Vezina P(2014) Neuronal and Psychological Underpinnings of Pathological Gambling. <i>Front. Behav. Neurosci.</i> 8:230. doi:10.3389/fnbeh.2014.00230
/Journal/FullText.aspx?s=99&name=behavioral%20neuroscience&ART_DOI=10.3389/fnbeh.2014.00230:	<a href="http://Journal/FullText.aspx?s=99&amp;name=behavioral%20neuroscience&amp;ART_DOI=10.3389/fnbeh.2014.00230">/Journal/FullText.aspx?s=99&amp;name=behavioral%20neuroscience&amp;ART_DOI=10.3389/fnbeh.2014.00230</a>
	(If clicking on the link doesn't work, try copying and pasting it into your browser.)
Copyright statement:	© 2014 Singer, Anselme, Robinson and Vezina. This is an open-access article distributed under the terms of the <a href="http://creativecommons.org/licenses/by/4.0/">Creative Commons Attribution License (CC BY)</a> . The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after rigorous peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

## **Neuronal and Psychological Underpinnings of Pathological Gambling**

Bryan F. Singer<sup>1\*</sup>, Patrick Anselme<sup>2</sup>, Mike J. F. Robinson<sup>3</sup>, Paul Vezina<sup>4</sup>

<sup>1</sup>Department of Psychology, The University of Michigan, Ann Arbor, Michigan, USA.

<sup>2</sup>Department of Psychology, The University of Liège, Liège, Belgium.

<sup>3</sup>Psychology Department, Wesleyan University, Middletown, CT, USA.

<sup>4</sup>Department of Psychiatry and Behavioral Neuroscience, The University of Chicago, Chicago, IL, USA.

**This is a General Commentary serving as the Introduction to the Frontiers in Behavioral Neuroscience Research Topic “Neuronal and Psychological Underpinnings of Pathological Gambling”**

Words: 997

Keywords: Dopamine, Gambling, Reward, Uncertainty, Addiction, Stress, Cortisol, Incentive Salience, Conditioning (Psychology), Reward Prediction Error, Ventral Striatum

\*, Correspondence:

Bryan F. Singer

Department of Psychology, The University of Michigan

530 Church Street

Ann Arbor, MI, USA, 48109

E-MAIL: [bfsinger@umich.edu](mailto:bfsinger@umich.edu)

Although pathological gambling (PG) is a prevalent disease, its neurobiological and psychological underpinnings are not well characterized. As legal gambling increases in prominence in a growing number of casinos as well as on the internet, the potential for a rise in PG diagnoses warrants investigation of the disorder. The recent reclassification of PG as a behavioral addiction in the DSM-5 raises the possibility that similar cognitive and motivational phenotypes may underlie both gambling and substance use disorders. Indeed, in this Research Topic, **Zack et al.** tested the hypothesis that exposure to reward unpredictability can recruit brain dopamine (DA) systems in a similar way to chronic exposure to drugs of abuse (see also Singer et al., 2012). Over the years a variety of models have proposed that alterations in DA signaling may mediate the transition from drug use to dependence; similarly, the hypothesis that aberrant DA responses may influence the transition from recreational, to problematic, and finally PG has only recently begun to be tested. The collection of articles in this Research Topic highlights the complexity of PG and posits several theories of how dopaminergic signaling may contribute to behavioral maladaptations that contribute to PG.

In this Research Topic, **Pagliari et al.** report a growing incidence of PG with a lack of effective treatments. As described by **Goudriaan et al.** (this Research Topic), PG is thought to result from “diminished cognitive control over the urge to engage in addictive behaviors” that manifests in the inability to control desire to gamble despite negative consequences. PG is characterized by several cognitive dysfunctions, including increased impulsivity and cognitive interference. Similar to drug addictions, gambling behavior is powerfully modulated by exposure to gambling-related conditioned stimuli. In this Research Topic both **Anselme & Robinson**, as well as **Linnet**, describe the supporting role of gambling-related cues in this behavioral addiction. **Anselme & Robinson** present a series of findings suggesting that surprising nonrewards enhance incentive salience attribution to conditioned cues in

conditioning procedures as well as during gambling episodes. They discuss a possible evolutionary origin of this counterintuitive process. **Linnet** reviews the contribution of DA signaling to incentive salience and reward prediction. Noting the research demonstrating brain activation during gambling tasks despite the possibility of a loss, he suggests a role for DA dysfunction in reward “wanting” and anticipation.

Ventral striatal activation is thought to be critical for the attribution of incentive salience to reward-related cues. In this Research Topic, **Lawrence & Brooks** found that healthy individuals who are more likely to display disinhibitory personality traits, such as financial extravagance and irresponsibility, show increased capacity for ventral striatal DA synthesis. Thus it is possible that individual variation in DA signaling due to genetics or environmental factors may influence PG. **Porchet et al.** (this Research Topic) also investigated whether physiological and cognitive responses observed during the performance of gambling tasks could be altered in recreational gamblers with pharmacological manipulations. As the commentary from **Zack** suggests, the **Porchet et al.** results may reflect important differences in neurobiological function between recreational and pathological gamblers. This hypothesis, along with the results of **Lawrence & Brooks** demonstrating increased DA capacity in individuals thought to be more prone to gambling, illustrates the complexity of PG as a disease and the need to sample different populations with different techniques and behavioral tasks.

Two papers in this Research Topic suggest a role for cortisol in modulating incentive motivation in the ventral striatum. **Li et al.** demonstrate an imbalanced sensitivity to monetary vs. non-monetary incentives in the ventral striatum of pathological gamblers. They show that cortisol levels in PG positively correlate with ventral striatal responses to monetary cues. **van den Bos et al.** provide further evidence for the importance of cortisol by highlighting the strong positive correlation observed in men between salivary cortisol levels and risk-taking measures.

This was a significant contrast to the weak negative correlation seen in women. Their findings highlight important gender differences in how stress hormones affect risky-decision making, and by extension, the role of stress in gambling.

In this Research Topic, **Clark & Dagher** provide a review of the literature investigating the relationship between DA agonists and impulse control disorders in Parkinson's patients, and how this relates to potential gains and losses within a decision-making framework. They provide the beginnings of a hypothetical model of how DA agonist treatments affect value and risk assessments. While a variety of research suggests that dopaminergic treatments for Parkinson's disease may affect PG, few have probed whether individuals with Huntington's disease (HD) display gambling-related phenotypes. **Kalkhoven et al.** (this Research Topic) show that HD patients exhibit symptoms of behavioral disinhibition similar to those observed in PG. However, HD patients do not typically develop problem gambling. Based on neurobehavioral evidence, these authors suggest why HD patients are unlikely to start gambling but have a higher chance of developing PG if they encounter a situation that promotes such behavior.

The investigation of neural mechanisms underlying PG is currently at an early stage. As emphasized by **Potenza** in this Research Topic, while previous research and the present findings suggest that DA may underlie gambling-related behaviors, other neurotransmitters and signaling pathways may also play vital roles in the emergence of the disease. Individual variation in PG populations (e.g., differing levels of impulsivity, compulsivity, decision making, and DA pathology) has produced discrepancies in the PG literature, warranting a systematic approach to investigating the disease in the future. **Paglieri et al.** also suggest the need for greater methodological integration of animal studies (rodents and primates) to better understand the mechanisms underlying PG. In particular, **Tedford et al.** note in this Research

Topic that gambling activity involves costs/benefits decision-making and that intracranial self-stimulation provides experimental advantages over traditional reinforcement methods used to model PG in animals. Finally, **Pagliari et al.** suggest that computational modeling, already used to account for other psychiatric diseases, might be applied to PG as well. Taken together, this collection of articles suggests novel avenues for future research of PG to improve treatment options for the disease.

## **Bibliography**

- Anselme, P., and Robinson, M. J. F. (2013). What motivates gambling behavior? Insight into dopamine's role. *Front. Behav. Neurosci.* 7, 182. doi:10.3389/fnbeh.2013.00182.
- Clark, C., and Dagher, A. (2014). The Role Of Dopamine In Risk Taking: A Specific Look At Parkinson's Disease And Gambling. *Front. Behav. Neurosci. In Press*. doi:10.3389/fnbeh.2014.00196.
- Goudriaan, A. E., Holst, R. J. Van, and Yücel, M. (2014). Getting a grip on problem gambling: What can neuroscience tell us ? *Front. Behav. Neurosci. In Press*. doi:10.3389/fnbeh.2014.00141.
- Kalkhoven, C., Sennef, C., Peeters, A., and van den Bos, R. (2014). Risk-taking and pathological gambling behavior in Huntington's disease. *Front. Behav. Neurosci.* 8, 103. doi:10.3389/fnbeh.2014.00103.
- Lawrence, A. D., and Brooks, D. J. (2014). Ventral striatal dopamine synthesis capacity is associated with individual differences in behavioral disinhibition. *Front. Behav. Neurosci.* 8, 86. doi:10.3389/fnbeh.2014.00086.
- Li, Y., Sescousse, G., and Dreher, J.-C. (2014). Endogenous cortisol levels are associated with an imbalanced striatal sensitivity to monetary versus non-monetary cues in pathological gamblers. *Front. Behav. Neurosci.* 8, 83. doi:10.3389/fnbeh.2014.00083.
- Linnet, J. (2014). Neurobiological underpinnings of reward anticipation and outcome evaluation in gambling disorder. *Front. Behav. Neurosci.* 8, 100. doi:10.3389/fnbeh.2014.00100.

- Paglieri, F., Addessi, E., De Petrillo, F., Laviola, G., Mirolli, M., Parisi, D., Petrosino, G., Ventricelli, M., Zoratto, F., and Adriani, W. (2014). Nonhuman gamblers: lessons from rodents, primates, and robots. *Front. Behav. Neurosci.* 8, 33. doi:10.3389/fnbeh.2014.00033.
- Porchet, R. I., Boekhoudt, L., Studer, B., Gandamaneni, P. K., Rani, N., Binnamangala, S., Müller, U., and Clark, L. (2013). Opioidergic and dopaminergic manipulation of gambling tendencies: a preliminary study in male recreational gamblers. *Front. Behav. Neurosci.* 7, 138. doi:10.3389/fnbeh.2013.00138.
- Potenza, M. N. (2013). How central is dopamine to pathological gambling or gambling disorder? *Front. Behav. Neurosci.* 7, 206. doi:10.3389/fnbeh.2013.00206.
- Singer, B. F., Scott-Railton, J., and Vezina, P. (2012). Unpredictable saccharin reinforcement enhances locomotor responding to amphetamine. *Behav. Brain Res.* 226, 340–4. doi:10.1016/j.bbr.2011.09.003.
- Tedford, S. E., Holtz, N. A., Persons, A. L., and Napier, T. C. (2014). A new approach to assess gambling-like behavior in laboratory rats: using intracranial self-stimulation as a positive reinforcer Stephanie. *Front. Behav. Neurosci. In Press*. doi: 10.3389/fnbeh.2014.00215.
- van den Bos, R., Taris, R., Scheppink, B., de Haan, L., and Verster, J. C. (2013). Salivary cortisol and alpha-amylase levels during an assessment procedure correlate differently with risk-taking measures in male and female police recruits. *Front. Behav. Neurosci.* 7, 219. doi:10.3389/fnbeh.2013.00219.
- Zack, M. (2013). Opioid and dopamine mediation of gambling responses in recreational gamblers. *Front. Behav. Neurosci.* 7, 147. doi:10.3389/fnbeh.2013.00147.
- Zack, M., Featherstone, R. E., Mathewson, S., and Fletcher, P. J. (2014). Chronic exposure to a gambling-like schedule of reward predictive stimuli can promote sensitization to amphetamine in rats. *Front. Behav. Neurosci.* 8, 36. doi:10.3389/fnbeh.2014.00036.