

Analysis of the Technical/Economic Performance of Four Cropping Systems Involving *Jatropha curcas* L. in the Kinshasa Region (Democratic Republic of the Congo)

J.D. Minengu¹, P. Mobambo¹ & G. Mergeai²

Keywords: *Jatropha*- Crop combinations- Yield- Family farm income- Kinshasa- DR Congo

Summary

In order to assess the sustainability of cultivating *Jatropha curcas* L. in rural areas in the Kinshasa region, four cropping systems were compared: cultivation of *J. curcas* as a sole crop with and without fertilisers, a combination of *J. curcas* with subsistence crops (maize - *Zea mays* L., the common bean - *Phaseolus vulgaris* L.) with and without fertilisers. The major attacks by pests (mainly *Apthona* sp.) suffered by *J. curcas* plants in the region make it vital to conduct at least two insecticide treatments per year. Dry seed yields of *J. curcas* obtained in the 4th year of cultivation amounted to 753 kg ha⁻¹ when *J. curcas* was cultivated as a sole crop without fertilisers, 797 kg ha⁻¹ for intercropping without fertilisers, 1158 kg ha⁻¹ when *J. curcas* was cultivated as a sole crop with fertilisers and 1173 kg ha⁻¹ for intercropping with fertilisers. Yields from the two annual crops were not improved by the application of mineral fertilisers on the *J. curcas* plants. They amounted to an average of 815 kg ha⁻¹ for maize and 676 kg ha⁻¹ for the beans. It is more profitable to cultivate *J. curcas* with maize and beans than to cultivate it as a sole crop. By combining crops in this way, a one-hectare farm can earn 1102 USD ha⁻¹ without fertilisers and 1049 USD ha⁻¹ with fertilisers. Sustainable cultivation of *J. curcas* under the test conditions requires the development of efficient weed/pest control methods and improved soil fertility management, in order to minimise the use of mineral fertilisers.

Résumé

Analyse des performances technico-économiques de quatre systèmes de culture basés sur la production de *Jatropha curcas* L. dans la région de Kinshasa (RDC)

Afin d'évaluer la durabilité de la production de *Jatropha curcas* L. dans la partie rurale de la région de Kinshasa, quatre systèmes de culture de la plante ont été comparés: la culture pure de *J. curcas* avec et sans engrais, l'association de *J. curcas* avec des cultures vivrières (maïs *Zea mays* L., haricot commun *Phaseolus vulgaris* L.) avec et sans engrais. Les fortes attaques de ravageurs (principalement *Apthona* sp.) subies par les plantes de *J. curcas* dans la région rendent indispensable la réalisation d'au moins deux traitements insecticides par an. Les rendements en graines sèches de *J. curcas* obtenus en 4^{ème} année de culture s'élevaient respectivement à 753 kg ha⁻¹ en culture pure de *J. curcas* sans engrais, 797 kg ha⁻¹ en cultures associées sans engrais, 1158 kg ha⁻¹ en culture pure de *J. curcas* avec engrais et 1173 kg ha⁻¹ en cultures associées avec engrais. Les rendements des cultures vivrières n'ont pas été améliorés par l'application d'engrais. Ils s'élevaient en moyenne à 815 kg ha⁻¹ pour le maïs et 676 kg ha⁻¹ pour le beans commun. La rentabilité de l'association de *J. curcas* avec le maïs et le beans commun est meilleure que la culture pure de *J. curcas*. Le revenu agricole d'un ha de cette association s'élève respectivement à 1102 USD ha⁻¹ sans engrais et à 1049 USD ha⁻¹ avec engrais. L'exploitation durable de *J. curcas* dans les conditions d'étude nécessite la mise au point de méthodes efficaces de contrôle des adventices et des ravageurs et d'amélioration de la fertilité du sol minimisant l'emploi d'engrais minéraux.

¹University of Kinshasa, Faculty of Agricultural Sciences, Kinshasa, Democratic Republic of the Congo.

²University of Liege, Gembloux Agro-Bio Tech, Laboratory of Tropical Crop Husbandry and Horticulture, Gembloux, Belgium.

*Corresponding author: E-mail: jminengum@yahoo.fr

Received on 11.03.13 and accepted for publication on 17.04.13.

Introduction

Jatropha curcas L. is a plant from the Euphorbiaceae family and comes from Central America. It was introduced to the Cap Verde islands by Portuguese sailors in the 16th century, followed by Guinea-Bissau. It later spread across Africa and Asia. It is cultivated in the tropical regions of the world (4). With its seeds rich in oil (35% on average) with a high fuel value, *J. curcas* has attractive characteristics for agrofuel production. However, the development of *J. curcas* plants and yields are low in most regions where it is grown (15). Some authors state that *J. curcas* can be cultivated on marginal land without the use of inputs (16, 19).

The Batéké Plateau near Kinshasa is an ideal place for testing the possibilities opened up by the cultivation of *J. curcas*, in terms of making full use of largely infertile land in a humid tropical climate. Using the soil for cultivation in this area actually leads to a rapid fall in its fertility, due to the mineralisation of the humus accumulated during fallow time. Different *J. curcas* cropping systems co-exist in this area, which are characterised by cultivation of this crop alone or in intercropping with subsistence crops (mainly the common bean *Phaseolus vulgaris* L. and maize *Zea mays* L.) and the quantity of mineral fertilisers used. This article aims to assess the technical/economic performance of the four cropping systems used for *J. curcas* cultivation (sole cropping with and without fertilisers; intercropping with maize, followed by beans with and without fertilisers) on a 4-year old plantation established in poor quality soil in rural areas in the province of Kinshasa.

Materials and methods

Characteristics of the study site

The study site is located on the Batéké Plateau (4°47' latitude south, 16°12' longitude east) at an altitude of 684 m. The plantation was set up after a short fallow period (5 years) on an area covering 3 ha in December 2007 in very poor soil, the arable horizon of which consisted of 94.2% sand (48.7% coarse sand 0.2–2 mm, 45.5% fine sand 0.05–0.2 mm), 1.4% silt (1.1% coarse silt 0.02–0.05 mm,

0.3% fine silt 0.002–0.02 mm), 3.3% clay (<0.002 mm), 5.3 g/kg organic carbon and 1.1% humus, with a pH of 5.3. The climate belongs to the AW4 type, according to Köppen's classification system. This humid tropical climate is characterised by a rainy season that lasts from mid-September until mid-May and is interrupted by a brief dry season between mid-January - mid-February. The first part of the rainy season occurring before the short dry season is called "season A" and the second part of the rainy season occurring after the short dry season is called "season B". The long dry season lasts four months – from mid-May until mid-September. The average annual temperature is 25 °C and rainfall fluctuates at around 1500 mm/year (18). Plant formations on the Batéké Plateau consist mainly of shrub savannahs, which alternate with grassy savannahs (23).

Composition and arrangement of test plots

The study was conducted on a plantation established on completely flat terrain. The planting density was 2500 plants per ha⁻¹ (2 m x 2 m). The planting materials consisted of plants grown from seedlings from seeds collected from a Mwabo ecotype (Bandundu Province in the DRC), which were grown for 3 months in a nursery. In order to increase the number of shoots, the *J. curcas* plants were pruned twice using the method advocated by Henning (12).

No inputs were used on the plants, which were not combined with other crops in the first two years after planting. Dry seed yields of *J. curcas* obtained during the first three years, after the plantation was established, rose to 97 kg ha⁻¹ in the first year, 182 kg.ha⁻¹ in the second year and 372 kg ha⁻¹ in the third year. In April 2011, the total area of the plantation was divided into four 0.5 ha plots (100 m x 50 m) on which the treatments indicated in Table 1 were applied in a random fashion.

Mineral fertilisers (50 kg urea.ha⁻¹ and 50 kg NPK 17-17-17 ha⁻¹) were applied on 5 April 2011 within 30 cm-radius circles at the foot of each *J. curcas* plant. As the canopy of *J. curcas* was still far from covering all the soil, 4 years after the plantation was established, the space left free made it possible to plant intercrops (Figure 1).

Table 1
J. curcas cropping systems compared in the test.

System without fertilisers (1 ha)	System with fertilisers (1 ha)
<i>Jatropha</i> crops alone 0.5 ha	<i>Jatropha</i> crops alone 0.5 ha
<i>Jatropha</i> combined with beans, followed by maize 0.5 ha	<i>Jatropha</i> combined with beans, followed by maize 0.5 ha



A

B

C

Figure 1: *Jatropha* cultivated on its own (A), intercropping of *Jatropha*-maize (B) and *Jatropha*-beans (C) in the 4th year of cultivating *Jatropha*.

Local varieties of the common bean (semi-erect dwarf variety from the Lower Congo, 10 April 2011) and maize (local population from the Batéké Plateau) were sown as intercrops on 3 October 2011, with 5 lines of beans (83,500 plants.ha⁻¹ – 0.3 m x 0.3 m) between two lines of *J. curcas* and 4 lines of maize (40,200 plants ha⁻¹ – 0.5 m x 0.4 m) between two lines of *J. curcas*.

Morphological observations and yields

The root collar diameter, plant height (main stem length) and total number of branches were measured on 3 April 2011 (first measurement) and 15 February 2012 (second measurement) on 125 *J. curcas* plants from each experimental unit. The dry seed yields of *J. curcas*, beans and maize were measured, after harvesting all the plants from each plot.

Calculation of technical/economic performance

This evaluation of the technical/economic performance of the different systems used to cultivate *J. curcas* is based on Dufumier's method (7). The amortisation period for *J. curcas* plantation

has been estimated at 30 years (4). For agricultural tools and drying racks, the amortisation period has been estimated at 5 and 2 years, respectively.

Validation of results

The study site is representative of the most unfavourable cultivation conditions on the plateau, in terms of soil fertility and pest pressure (*Apthona* sp. and *Stomphastis thraustica* Meyrick). In order to compare the results measured during the test to those obtained for *J. curcas* crops at other locations in the region, the other *J. curcas* plantations established on the Batéké Plateau (Menkao, Domaine des Sources in Mongata and N'sele) were visited from early 2008.

Results and discussion

Morphological characteristics of *J. curcas* plants

The use of fertilisers promoted the plant development for all the parameters analysed (Table 2). Vegetative growth achieved by *J. curcas* after four years of cultivation is low compared to that observed in other regions of the world, such as Brazil (Aw climate and oxisol soil type) and other locations in the Kinshasa region, where cultivation

Table 2
Comparison of influence of treatments on vegetative growth of *J. curcas* plants.

Cultivation methods	Root collar diameter (cm)			Plant height (m)			Number of shoots		
	On 3 April 2011	On 15 February 2012	Growth (cm)	On 3 April 2011	On 15 February 2012	Growth (m)	On 3 April 2011	On 15 February 2012	Increase
<i>Jatropha</i> as a sole crop without fertilisers	10.5±1.2	15.1±2.2	4.6	1.6±0.1	1.9±0.1	0.3	10±2.1	14±3.3	4
<i>Jatropha</i> as a sole crop with fertilisers	10.4±2.1	16.6±3.1	6.2	1.7±0.2	2.1±0.2	0.4	9±1.8	16±2.4	7
<i>Jatropha</i> as an intercrop without fertilisers	11.1±2.3	14.8±1.7	3.7	1.6±0.1	1.8±0.1	0.2	11±3.0	15±2.8	4
<i>Jatropha</i> as an intercrop with fertilisers	10.8±1.9	17.1±2.3	6.2	1.8±0.1	2.0±0.1	0.1	12±2.3	16±1.9	4

The averages and standard deviations were calculated using 125 plants selected at random for each treatment.

Table 3
Crop yield.

Cultivation methods	Yield (kg/ha/year) System without fertilisers			Yield (kg/ha/year) System with fertilisers*		
	Season B 2011	Season A 2011	Total	Season B 2011	Season A 2011	Total
<i>Jatropha</i> as a sole crop	271	482	753	376	782	1158
<i>Jatropha</i> as an intercrop	287	510	797	353	820	1173
Average yield: <i>Jatropha</i>	279	496	775	365	801	1166
Beans cultivated with <i>Jatropha</i>	667	-	667	684	-	684
Maize cultivated with <i>Jatropha</i>	-	783	783	-	846	846

Season B: from March until May. Season A: from September until December. * 50 kg of urea ha⁻¹ and 50 kg NPK 17-17-17 ha⁻¹.

conditions are more favourable. A study of *J. curcas* in Brazil indicates that average plant heights at 12, 24 and 36 months were 1.3±0.4 m, 2.3±0.3 m and 2.5±0.3 m, respectively (17). In a market garden located in N'sele, in the suburbs of Kinshasa, where the plants are irrigated and regularly fertilised, *J. curcas* grows to a height of over 2 metres after three years and the number of shoots per plant is well over 20. If, at the same time, we consider plants cultivated as sole crops and intercrops, the use of mineral fertilisers results in increased root collar diameter, height and number of branches. If we consider plants cultivated with and without the use of fertilisers, those that were cultivated as sole crops show the most growth in terms of the observed parameters.

Crop yields

The productivity of *J. curcas* depends on the rainfall, which determines the number of fructifications and therefore annual harvests (16). In the Kinshasa region, *J. curcas* can be harvested twice per year (Table 3). The first production is harvested between April and July (this period extends from Season B until the start of the dry season) and the second production is harvested between September and December (Season A). The better yields obtained in Season A can be explained by the higher and better distribution of rainfall during this period. Season B is shorter and rainfall is more irregular. Due to the very poor soil water economy, approx. 70% of the seeds contained in the capsules produced during the short rainy season fail to reach maturity, as the last stage of their development

occurs after the start of the long dry season (July). Tests involving the application of mulch on soil on *J. curcas* plots close to our crops have shown that this technique makes it possible to significantly improve the maturation of seeds produced during the short rainy season. The yield obtained by cultivating *J. curcas* is subject to the same order of magnitude, whether it is a sole crop or intercrop. If fertilisers are not used, it amounts on average to 774 kg ha⁻¹ (753 kg ha⁻¹ as a sole crop and 797 kg ha⁻¹ as an intercrop). If fertilisers are used, it reaches an average of 1166 kg ha⁻¹ (1158 kg ha⁻¹ as a sole crop and 1173 kg ha⁻¹ as an intercrop).

The use of mineral fertilisers only makes it possible to obtain an average gain of 396 kg ha⁻¹, which is equivalent to 4 kg of *J. curcas* seeds kg⁻¹ of fertiliser used. Until now, no research has been conducted, in order to determine the most suitable formula and dosage of mineral fertilisers for promoting *J. curcas* growth in the soil of the Batéké Plateau. It is possible that higher yields could be obtained with a different dosage of mineral fertilisers than that used during our test.

The average yields obtained for annual intercrops, with or without the use of fertilisers, were approx. 815 kg ha⁻¹ for maize (783 kg ha⁻¹ without fertilisers and 846 kg ha⁻¹ with fertilisers) and 675 kg ha⁻¹ for the common bean (667 kg ha⁻¹ without fertilisers and 684 kg ha⁻¹ with fertilisers). The absence of any effect caused by intercrops on the *J. curcas* yield can be attributed to the fact that *J. curcas* plants grown by seed propagation develop taproots, which hardly compete with the subsistence crops that are cultivated with them (6). This is true, as long as the perennial plant does not grow too much, in relation to the density of the two intercrops.

J. curcas is a plant, whose growth depends greatly on soil fertility and responds positively to the use of mineral fertilisers (20). The yields obtained after four years of cultivation are lower than those indicated for crops of the same age in Nicaragua (2500 kg ha⁻¹) with 650 mm rainfall year⁻¹ (8), in Sadivayal (India) (4000 kg ha⁻¹) with 2000 mm rainfall per year⁻¹ (11) and in Allahabad (India) (2000 kg ha⁻¹) with 1000 mm rainfall year⁻¹ (1). The very low soil fertility and major pest pressure suffered by the plants in the test area may explain

the low yields obtained. According to different authors (14, 19, 21), *J. curcas* produces its full potential yield after 3-5 years. However, recent studies (3, 5) have shown that it could take longer than 5 years for the shrubs to produce their maximum yield. It is likely that the *J. curcas* plants on the plots observed in Mbankana will continue to grow and therefore produce a higher yield during the next few years. It is difficult to predict with accuracy the maximum yield that they should produce, but it is unlikely that the latter will exceed the estimated yields suggested by Trabucco et al. (22) (2500 kg ha⁻¹) for the region, once they have reached their full potential.

Technical/economic performance of *J. curcas* cultivation systems

Gross income

As there are still no structured markets for *J. curcas* seeds in the DRC, the price of 0.125 USD kg⁻¹, based on the average selling price in other African countries where a market already exists (10), has been used to calculate the gross income generated. The gross income generated by cultivating *J. curcas* seeds as a sole crop and without using fertilisers (94 USD ha⁻¹) is lower than that generated by the corresponding intercrops (1236 USD ha⁻¹) (Table 4). The use of fertilisers on *J. curcas* only makes it possible to obtain an additional gain of approx. 100 US dollars ha⁻¹, if it is cultivated as an intercrop with subsistence crops, and about 50 US dollars ha⁻¹ if it is cultivated as a sole crop.

Inputs

In cropping systems without mineral fertilisers, the most costly inputs are the insecticides (40 USD ha⁻¹ year⁻¹) (Table 5). The use of pesticides is vital in the study area, if *J. curcas* is to be cultivated. If they are not used, the leaves and buds are completely destroyed by a beetle of the *Aphthona* genus (18). In cropping systems with mineral fertilisers, the latter represent by far the most costly input (150 USD ha⁻¹ year⁻¹).

Table 4
Gross income ha⁻¹ year⁻¹.

Cropping systems	<i>Jatropha</i> (USD)			Maize			Beans			Total GI (USD)
	Harvest (kg.ha ⁻¹)	Selling price (USD kg ⁻¹)	GI (USD)	Harvest (kg.ha ⁻¹)	Selling price (USD kg ⁻¹)	GI (USD)	Harvest (kg.ha ⁻¹)	Selling price (USD kg ⁻¹)	GI (USD)	
<i>Jatropha</i> as a sole crop without fertilisers	753	125	94	-	-	-	-	-	-	94
<i>Jatropha</i> cultivated as a sole crop with fertilisers	1158	125	145	-	-	-	-	-	-	145
<i>Jatropha</i> cultivated as an intercrop without fertilisers	797	125	100	783	0.6	470	667	1	667	1236
<i>Jatropha</i> cultivated as an intercrop with fertilisers	1173	125	147	846	0.6	508	684	1	684	1338

Key: GI (gross income)

Table 5
Inputs.

Expenses	Production costs USD ha ⁻¹ year ⁻¹			
	<i>Jatropha</i> as a sole crop without fertilisers	Intercropping (<i>Jatropha</i> +beans+maize) without fertilisers	<i>Jatropha</i> as a sole crop with fertilisers	Intercropping (<i>Jatropha</i> +beans+maize) with fertilisers
Fertilisers (Urea+NPK)	0	0	150	150
Insecticides	40	40	40	40
Bean seeds	0	15	0	15
Maize seeds	0	10	0	10
<i>Jatropha</i> harvest bags	10	11	15	16
Maize harvest bags	0	5	0	5
Bean harvest bags	0	3	0	3
Total	50	84	205	239

Table 6
Cost of planting one hectare of *J. curcas* shrubs (2,500 plants).

Expenses	Cost in USD/ha
<i>Jatropha</i> seeds	18
Ploughing and harrowing (tractor)	160
Preparing and maintaining plants in the nursery for 3 months (5 man-days)	50
Stumping (10 man-days)	20
Staking out and digging (15 man-days)	30
Planting <i>Jatropha</i> plantlets (5 man-days)	10
Replanting <i>Jatropha</i> (2 man-days)	4
Total	292

Table 7
Required labour (man-days ha⁻¹ year⁻¹).

Activities/tasks	<i>Jatropha</i> as a sole crop without fertilisers	Intercropping (<i>Jatropha</i> +beans+maize) without fertilisers	<i>Jatropha</i> as a sole crop with fertilisers	Intercropping (<i>Jatropha</i> +beans+maize) with fertilisers
Maize seeding	0	10	0	10
Bean seeding	0	15	0	15
Weeding	80	80	80	80
<i>Jatropha</i> plant pruning	5	5	5	5
Mineral fertilisation	0	0	10	10
Phytosanitary treatment	10	10	10	10
<i>Jatropha</i> harvesting and shelling	47	50	72	73
Maize harvesting	0	10	0	10
Bean harvesting and shelling	0	15	0	15
Maize shelling	0	15	0	15
Total	142	210	177	243

Table 8
Family farm income.

Cropping systems	GI ha ⁻¹ year ⁻¹ (USD)	I.C. ha ⁻¹ year ⁻¹ (USD)	Amortisation ha ⁻¹ year ⁻¹ (USD)	Land rent ha ⁻¹ year ⁻¹ (USD)	FI ha ⁻¹ year ⁻¹ (USD)	No. man-days ha ⁻¹ year ⁻¹	F.I. man-days ⁻¹ (USD)
<i>Jatropha</i> as a sole crop without fertilisers	94	50	46	4	-6	142	0
Intercropping without fertilisers	1236	84	46	4	1102	210	5,2
<i>Jatropha</i> as a sole crop with fertilisers	145	205	46	4	-144	177	-0,8
Intercropping with fertilisers	1338	239	46	4	1049	243	4,3

Key: GI (Gross income), IC (Input cost), FI (Family farm income).

Amortisation

In this context, amortisation refers to the initial cost of establishing the plantation and purchasing small agricultural equipment. The cost of setting up a one-hectare plantation is estimated at 292 USD (Table 6) and the life span of this investment is estimated at 30 years. The small equipment used for the cultivation of *J. curcas* consists of small agricultural tools (hoes, spades, rakes and backpack pressure sprayer), which cost a total of 80 USD and pay for themselves over 5 years (16 USD year⁻¹). The last fixed asset consists of trays used for drying *J. curcas* seeds, two of which are needed per ha. They cost 20 USD each and pay for themselves over a 2-year period. The total value of annual amortisations is therefore 46 USD.

Labour

Weeding, harvesting and shelling *J. curcas* fruit are the most labour-intensive tasks (Table 7). Eighty man-days ha⁻¹ year⁻¹ are needed for weeding and 50 man-days ha⁻¹ year⁻¹ for harvesting and shelling fruit, for an average yield of 800 kg ha⁻¹ year⁻¹ of dry seeds, based on 16 kg of dry seeds collected per person and per day (harvesting and shelling of the capsules).

As the longer the plantation is cultivated the more the yield increases, it is certain that the labour required to complete seed collecting tasks will also increase. The high number of man-days devoted to weeding is linked to major pressure from weeds (*Cynodon dactylon* L., *Digitaria* sp., *Imperata cylindrica*, etc.) on cultivated plots on the Batéké Plateau. The quantity of seeds obtained per man-

day under the conditions of our test appears low compared to figures published for other parts of the world (9). It is, however, consistent with the figures put forward by Henning (13) based on observations in Mali. The number of capsules harvested per working hour depends on a range of factors: the height and width of the shrub, method used to collect fruits, planting density and productivity of each shrub (the higher the yield, the more efficient the collecting process becomes) (5).

Land rent

The land rent (4 USD year⁻¹) was calculated by dividing the purchase price of the land, on which the plantation was established (120 USD ha⁻¹), by the life span of the plantation (30 years).

Family farm income

As a sole crop, if the selling price is 0.125 USD kg⁻¹ for dry seeds, the cultivation of *J. curcas* results in a loss of family farm income amounting to 6 USD ha⁻¹ without fertilisers and 144 USD ha⁻¹ with fertilisers (Table 8). Based on the price of 0.125 USD kg⁻¹, the dry seed yields that need to be produced, in order to cover the costs associated with *J. curcas* cultivation as a sole crop, amount to 872 kg.ha⁻¹ without fertilisers and 2112 kg.ha⁻¹ with fertilisers. In order to make 2 USD man-day⁻¹, depending on conditions when the plantation was established, the selling price of *J. curcas* seeds should be 0.52 USD kg⁻¹ without fertilisers and 0.54 USD kg⁻¹ with fertilisers. The profitability of combining *J. curcas* cultivation with maize and beans is better than that of cultivating *J. curcas* as a sole crop, in terms of income per ha and income per man-day. Due to the selling price of *J. curcas* seeds, the use of mineral fertilisers is not at all viable. One kg of fertilisers costing on average 1.5 USD only makes it possible to increase production by 4 kg dry seeds or 0.5 USD (for a seed selling price of 0.125 USD). The application of at least two insecticide treatments is vital for the growth of *J. curcas* plants. As has been observed in other parts of the world, it is likely that pest pressure increases as the cultivated areas are extended (2, 6). The very high cost of inputs (fertilisers and insecticides), the amount of work that it takes to weed the plots and

the small quantities of seeds harvested per man-day are the main causes of the very small amounts paid for family labour, as calculated for the cultivation of *J. curcas* as a sole crop.

A major reduction in production costs for *J. curcas* as a sole crop could be obtained, by planting a cover crop that requires little or no maintenance. However, with the productivity observed during the test in terms of collecting seeds, even an infinite yield increase would not make it possible to pay family workers an amount that would correspond to the labour opportunity cost (2 USD day⁻¹). In fact, the amount paid for one working day spent harvesting, shelling pods and drying seeds corresponds to its opportunity cost (16 kg of dry seeds day⁻¹ x 0.125 USD kg⁻¹ = 2 USD day⁻¹). This means that, based on the selling price of 0.125 USD kg⁻¹, the cultivation of *J. curcas* as a sole crop will always result in a loss, as long as no solution is found, in order to increase the productivity of work devoted to harvesting and post-harvesting.

In other regions of the world, the increased yield ha⁻¹ and use of mechanical shellers (6, 9) make it possible to achieve yields of 40 kg seeds collected per man-day. With this kind of daily yield, based on a seed selling price of 0.125 USD kg⁻¹, the yield to be achieved per ha, in order to cover all expenses, if fertilisers are used (including the labour at its opportunity cost), amounts to 6129 kg ha⁻¹ with manual weeding and 3549 kg.ha⁻¹ if a cover crop is planted. If we allow for a seed collection yield of 40 kg man-day⁻¹ and current production costs (with the use of fertilisers and labour costs estimated at their opportunity cost), the selling price of seeds, in order to cover running costs, if the maximum yield reaches 2500 kg ha⁻¹, amounts to 0.215 USD kg⁻¹ with manual weeding and 0.143 USD kg⁻¹ if a maintenance-free cover crop is planted.

Conclusion

The quantification of the technical and economic performance of *J. curcas* sole cropping, with and without the use of fertilisers, on a 4-year old plantation in the Batéké Plateau region, has made it possible to highlight the absence of financial profitability for the production of this crop, using the cultivation methods compared in this study. Based

on a seed selling price similar to that given in other African countries (0.125 USD kg⁻¹), losses in family farm income per ha amount to 6 USD if no fertilisers are used and 144 USD if fertilisers are applied. The low yields obtained, the very high cost of inputs (fertilisers and insecticides) and weeding, combined with the small quantities of seeds harvested per man-day, are the main causes of the low income generated by cultivating of *J. curcas* as a sole crop. The profitability of combined cultivation of *J. curcas* shrubs with maize and beans is greater than for the

cultivation of *J. curcas* as a sole crop. Family farm income from one ha, if crops are combined in this way, amounts to 1102 USD ha⁻¹ without fertilisers and 1049 USD ha⁻¹ with fertilisers. Cultivating *J. curcas* as an intercrop with subsistence crops therefore appears the best solution for establishing rural plantations in the region. Sustainable cultivation of these plantations will require, however, the development of efficient methods for controlling weeds and pests, as well as improving soil fertility, while minimising the use of mineral fertilisers and pesticides.

Literature

1. Achten W.M.J., Verchot L., Franken Y.J., Mathijs E., diesel production and use, *Biomass Bioenergy*, 32, 1063-1084.
2. Achten W.M.J., Maes W.H., Aerts R., Verchot L., Trabucco A., Mathijs E., Singh V.P. & Muys B., 2010, *Jatropha: from global hype to local opportunity*, *J. Arid Environ.*, 74, 1, 164-165.
3. Anonyme, 2011, Evaluation et optimisation du potentiel de développement d'une culture oléagineuse à hautes performances énergétique et environnementale pour la production de biodiesel, le *Jatropha curcas*. Rapport final Cirad, Enerbio, Fondation Tuck. <http://www.fondation-tuck.fr/resultats/projets/2008/documents-projet-01/2008-P01-rapport-final.pdf>, (29/02/2012).
4. Brittain R. & Litaladio N., 2010, *Jatropha: a smallholder bioenergy crop. The potential for pro-poor development*. Integrated Crop Management, vol 8. Food and Agriculture Organization of The United Nations, Rome.
5. De Jongh J. & Nielsen F., 2011, Lessons Learned: *Jatropha* for local development. Repport for FACT Foundation.
6. Domergue M. & Pirot R., 2008, *Jatropha curcas* L. Rapport de synthèse bibliographique, CIRAD, France, 118.
7. Dufumier M., 1996, Les projets de développement agricoles, manuel d'expertise. Karthala, Paris, France.
8. Foidl N., Foidl G., Sanchez M., Mittelbach M. & Hackel S., 1996, *Jatropha curcas* L. as a source for the production of bio-fuel in Nicaragua, *Bioresour. Technol.*, 58, 77-82.
9. Franken Y.J. & Nielsen F., 2010, Plantation establishment and management. In: *The Jatropha Handbook: From Cultivation to Application*, Eds: Jan de Jongh. Fact Foundation, Netherlands, 8-27
10. GEXSI, 2008, Global Market Study on *Jatropha*. Final Report. Prepared for the World Wide Fund for Nature (WWF). London/Berlin: Global Exchange for Social Investment.
11. Gunaseelan V.N., 2009, Biomass estimates, characteristics, biochemical methane potential, kinetics and energy flow from *Jatropha curcas* on dry lands, *Biomass Bioenergy*, 33, 589-596.
12. Henning R., 2007, Identification, selection and multiplication of high yielding *Jatropha curcas* L. plants and economic key points for viable *Jatropha* oil production costs. (Henning paper).
13. Henning R., 2009, The *Jatropha* System- An integrated approach of rural development. The *Jatropha* Book.
14. Jongschaap R.E.E., Corre W.J., Bindraban P.S. & Brandenburg W.A., 2007, Claims and facts on *Jatropha curcas* L.: Global *Jatropha curcas* evaluation, breeding and propagation programme. Report 158 (Plant Research International BV, Wageningen, The Netherlands and Stichting Het Groene Woudt, Laren, The Netherlands).
15. Les Amis de la terre, 2010, Afrique: terre(s) de toutes les convoitises, Ampleur et conséquences de l'accaparement des terres pour produire des agrocarburants. Rapport, 36.

-
16. Maes W.H., Trabucco A., Achten W.M.J. & Muys B., 2009, Climatic growing conditions of *Jatropha curcas* L, *Biomass Bioenergy*, 33, 1481-1485.
17. Moreira Miquelino E.T.C., António Gonçalves Jacovine L., De Paula Toledo D., Pedro Boechat Soares C., Cerruto Ribeiro S. & Cristina Martins M., 2011, Biomass and Carbon stock in *Jatropha curcas* L. *Cerne, Lavras*, 17, 3, 353-359.
18. Nyst J., 2010, Contribution à l'étude des possibilités de culture de *Jatropha curcas* L. sur le Plateau des Batéké RDC. Travail de fin d'études présenté en vue de l'obtention de diplôme de master bioingénieur en sciences agronomiques, orientation agronomie tropicale, Université de Liège-Gembloux Agro-Bio Tech, Belgique, 81.
19. Openshaw K., 2000, A review of *Jatropha curcas*: an oil plant of unfulfilled promise, *Biomass Bioenergy*, 9, 1-15.
20. Patolia, J., Ghosh A., Chikara J., Chaudhary D.R., Parmar D.R. & Bhuvra H.M., 2007, Response of *Jatropha curcas* grown on wasteland to N and P fertilization. Expert Seminar on *Jatropha curcas* L. Agronomy and Genetics. Wageningen
21. Rao G.R., Korwar G.R., Shanker A.K. & Ramakrishna Y.S., 2008, Genetic associations, variability and diversity in seed characters, growth, reproductive phenology and yield in *Jatropha curcas* (L.) accessions, *Trees- Structure Function*, 22, 5, 697-709.
22. Trabucco A., Achten W.M.J., Bowe C., Aerts R., Van Orshoven J., Norgrove L. & Muys B., 2010, Global mapping of *Jatropha curcas* yield based on response of fitness to present and future climate. *Global Change Biol, Bioenergy*, 2, 3, 139-151.
23. Vermeulen C. & Lanata F., 2006, Le domaine de chasse de Bombo-Lumene: un espace naturel en péril aux frontières de Kinshasa. *Parcs et réserves*, 61, 2, 4-8.
-

J.D. Minengu, Congolese, PhD Student, Assistant, University of Kinshasa, Faculty of Agricultural Sciences, Kinshasa, Democratic Republic of the Congo.

P. Mobambo, Congolese, PhD, Professor, University of Kinshasa, Faculty of Agricultural Sciences, Kinshasa, Democratic Republic of the Congo.

G. Mergeai, Belgian, PhD, Professor, University of Liege, Gembloux Agro-Bio Tech, Laboratory of Tropical Crop Husbandry and Horticulture, Gembloux, Belgium.