

Organic and inorganic carbon fluxes in a tropical river system (Tana River, Kenya) during contrasting wet seasons

Naomi Geeraert ^{(1,*),} Fred O. Omengo ^{(1,2),} Steven Bouillon ^{(1),} Alberto V. Borges ^{(3),} and Gerard Govers ⁽¹⁾

⁽¹⁾ KU Leuven, Department of Earth and Environmental Sciences, Celestijnenlaan 200E, 3001 Leuven, Belgium

⁽²⁾ Kenya Wildlife Service, P.O. Box 40241-00100, Nairobi, Kenya

⁽³⁾ Université de Liège, Chemical Oceanography Unit, Belgium

* naomi.geeraert@ees.kuleuven.be



Introduction

➤ The majority of discharge in tropical river systems occurs in relatively short time periods. E.g. in Tana River, ~60% of the annual discharge in 4 months.



➤ This seasonality has effect on the carbon (C) transport as C processes, such as respiration and primary production vary with discharge.


➤ Furthermore, connection with the floodplain during flooding significantly alters the biogeochemical functioning.

Research question: What is the role of floodplains on delivery or retention of C to the main channel?

Study area and methods

➤ 2 sampling sites: Garissa and Garsen, situated 385 km apart in the lower Tana River, Kenya

➤ Daily samples to measure the concentration of particulate and dissolved organic carbon (POC and DOC) and dissolved inorganic carbon (DIC) *.



➤ 3 sampling campaigns: Oct-Nov 2012, May-June 2013 and April-May 2014.


➤ In 2013, considerable overbank flooding took place between the sampling sites. No significant flooding occurred in the other wet seasons.

➤ Carbon loads were calculated by multiplying water discharge and carbon concentration. Missing values were interpolated based on the flux-discharge relationship. Seasonal fluxes are the sum of the loads.

* **POC:** 25 ml of water filtered on 25 mm Whatman GF/F filters. Concentration measured on EA-IRMS (ThermoFinnigan Flash HT). **DOC:** 40 ml of water filtered on 0.2 µm Acrodisc syringe filters, preserved with H₃PO₄. Analysed on a wet oxidation TOC-analyzer (IO Analytical Aurora 1030W) coupled with an IRMS (ThermoFinnigan DeltaV Advantage). **DIC:** calculated from TA (automated electro-titration) and pCO₂ (headspace technique on LI-820 gas analyzer) with thermodynamic constants from Millero.

Results


Carbon loads




- Highest DOC and DIC loads during peaks with flooding
- Highest POC loads during peaks without flooding
- At low discharge, daily loads increase in the downstream direction for all C pools (end of 2014)

Conclusion

Related to changes in the dominant carbon pool, the floodplains between both sites are a source of C towards the river during the flooded season and a sink of riverine C during the non-flooded seasons.



Seasonal fluxes



Water flux
POC
DIC
DOC

GSA: Garissa
GSN: Garsen

Numbers in the bars are percentage share of the carbon pool per site and per season

- Downstream decrease in water and total C flux during non-flooded seasons; Downstream increase in water and total C flux during the flooded season
- POC flux decreased downstream (33%, 8% and 30% in 2012, 2014 and 2013)
- DIC flux increased downstream (9%, 6% and 62% in 2012, 2014 and 2013)
- DOC flux decreased during non-flooded seasons (38% and 10% in 2012 and 2014), and increased by 163% in the flooded season (2013)
- POC was the dominant C pool during the non-flooded season; DIC was the dominant pool during the flooded season

Notes

➤ The calculated loss/gain of C are net fluxes between the two sites. Fluxes between the different pools are not yet taken into account.

➤ The fluxes are only valid for the observation periods and can not be directly extrapolated to the whole season or year.

Acknowledgements

Funding for this work was provided by the Research Foundation Flanders (FWO-Vlaanderen, project G024012N), KU Leuven BOF, and ERC-StG 240002 (AFRIVAL, <http://ees.kuleuven.be/project/afriaval/>). We are grateful to the Kenya Wildlife Service for the field assistance.