

Freikofel Formation

MONICA PONDRELLI, DAMIEN PAS, CLAUDIA SPALLETTA, HANS-PETER SCHÖNLAUB, ENZO FARABEGOLI, CARLO CORRADINI,
THOMAS J. SUTTNER, MARIA G. CORRIGA, MARIA CRISTINA PERRI, ANNE-CHRISTINE DA SILVA, SUSANNE M.L. POHLER,
LUCA SIMONETTO, CLAUDIA DOJEN, ANGELO MOSSONI, ERIKA KIDO & HEIKO HÜNEKE

Österreichische Karte 1:50.000

Blatt BMN 197 Kötschach

Blatt UTM 3109 Oberdrauburg

Blatt UTM 3110 Kötschach-Mauthen

Blatt UTM 3116 Sonnenalpe Naßfeld

Carta Topografica d'Italia 1:50.000

Foglio 018 Passo di Monte Croce Carnico

Foglio 031 Ampezzo

Foglio 032 Tolmezzo

Definition

Prevalent floatstone/rudstone interbedded with lithoclastic packstone/grainstone and pelagic mudstone/wackestone (BANDEL, 1972; SPALLETTA et al., 1982; SPALLETTA & VAI, 1984; SCHÖNLAUB, 1985; KREUTZER, 1990, 1992a, b; SCHÖNLAUB et al., 2004; SCHNELLBÄCHER, 2010; PAS et al., 2014).

Description

The Freikofel Formation consists of three well-bedded facies listed in order of decreasing abundance.


Facies A: medium dark gray, medium to thick bedded, moderately sorted lithoclastic rudstone (subordinately floatstone) sometimes showing fining upward grading; matrix consists of grainstone (subordinately wacke-/packstone);

Facies B: medium dark gray, thin to medium bedded grainstone and subordinate packstone locally showing fining upward grading; planar and subordinate cross lamination is present;



Facies C: very thin to thin bedded, moderate pink to gray mud-/wackestone.

Fossil content

Brachiopods, calcispheres, cephalopods, conodonts, crinoids, foraminifers, trilobites, ostracods, rugose corals, stromatoporoids and stylolinids have been documented (KREUTZER, 1992b; PAS et al., 2014).



Areas of outcrop of the Freikofel Formation with indication of the stratotype (asterisk).



The Freikofel Section. a) log of the Freikofel Formation (SCHNELLBÄCHER, 2010); b) view of the section taken from the west (photo M. PONDRELLI).

Depositional environment

The Freikofel Formation was formed at the slope of a carbonate apron (SCHNELLBÄCHER, 2010; PAS et al., 2014). Deposits of hyperconcentrated and concentrated density flows (Facies A) and turbidity flows (Facies B) predominate. Thin interbeds of pelagic sediments (Facies C and D) are rarely preserved.

Stratotype

Freikofel Section, located on the western slope of Mt. Freikofel (BANDEL, 1972; SCHNELLBÄCHER, 2010; PAS et al., 2014), between coordinates N 46°36'05.1", E 12°58'33.9" (base of the section) and N 46°36'08.0", E 12°58'35.0" (top of the section).

Reference sections -

Type area

Central Carnic Alps.

Main outcrop areas


The Freikofel Formation crops out in the Rauchkofel area, in the south side of the Valentintal and between the Cellon/Creta di Collinetta and the Creta di Timau.

Thickness

About 65 m (Freikofel) to about 100 m (Kellerwand).

Boundaries

Underlying units – Cellon Formation (conformable sharp contact).



Views of the Freikofel Formation in the field. a) breccia bed: top of Mt. Freikofel (photo M. PONDRELLI); b) part of a breccia bed showing a crude fining upward trend: Freikofel Section (photo M. PONDRELLI); c) rudstone: PR327 section (photo C. SPALLETTA); d) particular of a rudstone bed: PR327 section (photo C. SPALLETTA).

Overlying units – Pal Grande Formation (conformable interfingering contact).

Lateral units – Kellergrat Formation, Creta di Collina Formation (proximal part); Hoher Trieb Formation, Pal Grande Formation (distal part).

Derivation of name

After Mount Freikofel.

Synonymy

'Lithoklastkalk' [partim]: BANDEL (1974).

Freikofel Kalk: SCHÖNLAUB (1985).

Freikofel limestone: KREUTZER (1992b).

Calcareniti di transizione prossimali [partim]: VENTURINI (2006).

Freikofel Rudstone [partim]: BRIME et al. (2008).

Calciruditi del Freikofel [partim]: SPALLETTA & PONDRELLI (2009).

Chronostratigraphic age

Devonian: Givetian to Frasnian (KREUTZER, 1990, 1992a, b; SCHNELLBÄCHER, 2010; PAS et al., 2014).

Biostratigraphy

Conodonts. – The Freikofel Formation spans to Frasnian Zone 13b (*linguiformis* Zone) (Freikofel T section)-Frasnian Zone 13a (middle-upper part of the Upper *rhenana* Zone) (Pramosio A section).

Complementary references -

Remarks -

References

BANDEL, K. (1972): Palökologie und Paläogeographie im Devon und Unterkarbon der zentralen Karnischen Alpen. – *Palaeontographica Abteilung A*, **141**/1–4, 1–117, Stuttgart.

BANDEL, K. (1974): Deep-water limestones from the Devonian-Carboniferous of the Carnic Alps, Austria. – Special Publications of the International Association of Sedimentologists, **1974**/1, 93–115, Oxford.

BRIME, C., PERRI, M.C., PONDRELLI, M., SPALLETTA, C. & VENTURINI, C. (2008): Polyphase metamorphism in the eastern Carnic Alps (N Italy-S Austria): Clay minerals and conodont Colour Alteration Index evidence. – *International Journal of Earth Sciences*, **97**/6, 1213–1229, Berlin-Heidelberg.

KREUTZER, L.H. (1990): Mikrofazies, Stratigraphie und Paläogeographie des Zentralkarnischen Hauptkammes zwischen Seewarte und Cellon. – *Jahrbuch der Geologischen Bundesanstalt*, **133**/2, 275–343, Wien.

KREUTZER, L.H. (1992a): Palinspastische Entzerrung und Neugliederung des Devons in den Zentralkarnischen Alpen aufgrund von neuen Untersuchungen. – *Jahrbuch der Geologischen Bundesanstalt*, **135**/1, 261–272, Wien.

KREUTZER, L.H. (1992b): Photoatlas zu den variszischen Karbonat-Gesteinen der Karnischen Alpen (Österreich/Italien). – *Abhandlungen der Geologischen Bundesanstalt*, **47**, 1–129, Wien.

PAS, D., DA SILVA, A.-C., SUTTNER, T., KIDO, E., BULTYNCK, P., PONDRELLI, M., CORRADINI, C., DE VLEESCHOUWER, D., DOIEN, C. & BOULVAIN, F. (2014): Insight into the development of a carbonate platform through a multi-disciplinary approach: a case study from the Upper Devonian slope deposits of Mount Freikofel (Carnic Alps, Austria/Italy). – *International Journal of Earth Sciences*, **103**, 519–538, Berlin-Heidelberg.

SCHNELLBÄCHER, P. (2010): Stratigraphy, transport mechanisms and provenience of Devonian carbonate slope deposits (Freikofel, Carnic Alps). – Unpublished master thesis, Universität Greifswald. – 124 p., Greifswald.

SCHÖNLAUB, H.P. (1985): Das Paläozoikum der Karnischen Alpen. – In: SCHÖNLAUB, H.P. (ed.): *Arbeitstagung der Geologischen Bundesanstalt 1985 Kötschach-Mauthen, Gailtal – Geologische Bundesanstalt*, 34–52, Wien.

SCHÖNLAUB, H.P., HISTON, K. & POHLER, S. (2004): The Palaeozoic of the Carnic Alps. – In: SCHÖNLAUB, H.P. (ed.): *Field Trip Carnic Alps Guidebook. June 23–24, 2004, Carinthia, Austria*. – Geologische Bundesanstalt, 2–32, Wien.

SPALLETTA, C. & PONDRELLI, M. (2009): Calciruditi del Freikofel. – In: VENTURINI, C. (ed.): *Note Illustrative del Foglio 031 Ampezzo. – Carta Geologica d'Italia alla scala 1:50000, Istituto Superiore per la Protezione e la Ricerca Ambientale (ex-Agenzia per la Protezione dell'Ambiente e per i Servizi Tecnici, Servizio Geologico d'Italia)*, 43, Stampa A.T.I. – S.E.L.C.A. srl. – L.A.C. srl. – System Cart srl, Firenze.

SPALLETTA, C. & VAI, G.B. (1984): Upper Devonian intraclast para-breccias interpreted as seismites. – *Marine Geology*, **55**/1–2, 133–144, Amsterdam.

SPALLETTA, C., VAI, G.B. & VENTURINI, C. (1982): La Catena Paleocarnica. – In: CASTELLARIN, A. & VAI, G.B. (eds.): *Guida alla geologia del Sudalpino centro-orientale. – Guide Geologiche Regionali, Società Geologica Italiana*, 281–292, Bologna.

VENTURINI, C. (2006): Evoluzione geologica delle Alpi Carniche. – *Edizioni del Museo Friulano di Storia Naturale*, **48**, 208 p., Udine.