

itobiase y est évidente

Gmelin, dont le régime  
ons pas trouvé trace de  
ans les extraits aqueux  
mais faible élaboration  
muqueuses gastrique et

re démonstration d'une  
de chitinase chez un

siol. Bioch., 66, 408.

rch. internat. Physiol. Bioch.,

(Reprinted from *Nature*, Vol. 192, No. 4798, pp. 135-136,  
October 14, 1961)

## CHITINASE: AN ADDITION TO THE LIST OF HYDROLASES IN THE DIGESTIVE TRACT OF VERTEBRATES

By DR. CHARLES JEUNIAUX

Department of Biochemistry, University of Liège

IT was thought for a long time that, in animals, chitinases were present in snails only. The distribution of these enzymes among invertebrates has appeared to be much more extensive. They have been found by Tracey in protozoa<sup>1</sup>, in nematodes<sup>2</sup>, and in earthworms<sup>3</sup>, and by me in arthropods<sup>4-6</sup>, coelenterates<sup>7</sup>, polychaete worms<sup>7</sup>, etc. The presence of chitinolytic enzymes among vertebrates has, however, never been suspected. The enzymic lysis of chitin is known to be actually performed by two different enzymes, acting consecutively<sup>8-9</sup>: a chitinase (mucopolysaccharidase) and a chitobiase ( $\beta$ -acetylglucosaminidase). This 'chitinolytic system' and these two enzymes have been searched for in the digestive juices and in aqueous extracts of digestive glands and digestive epithelia of some fishes, lizards, birds and mammals.

In order to detect the chitinolytic system (chitinase + chitobiase), the enzymic extracts were incubated with a chitin suspension (from *Sepia pen*) at 37° C. and pH 5.2 for 3 hr.; the acetylglucosamine liberated was measured by a method described by Reissig *et al.*<sup>10</sup>. Controls without enzyme or without substrate were run simultaneously.

In order to detect chitinase in enzymic extracts containing little or no chitobiase, the activity of these extracts on chitin has also been investigated after addition of an excess of chitobiase in the enzymic test. A commercial preparation of  $\beta$ -glucosidase (Nutritional Biochemicals Corporation) provided a convenient source of chitobiase, free from chitinase. In these cases, the presence of chitinase was also confirmed by the nephelometric method<sup>11</sup>, which does not depend on the presence or absence of chitobiase<sup>9</sup>. In order to measure the chitobiase, the liberation of acetylglucosamine has been followed, using a preparation of 'depolymerized chitin' as substrate<sup>12</sup> (a preparation which contains chitobiose and other small polymers of acetylglucosamine).

64.

63.

Table 1. CHITINOLYTIC ENZYMES IN DIGESTIVE JUICES OF SOME VERTEBRATES

Species	Digestive juices or contents	Activity of chitinolytic system: acetylglucosamine liberated, $\mu\text{gm./hr./ml.}$
<i>Carassius auratus</i> L.	Anterior intestine	118
	Posterior intestine	814
<i>Gasterosteus aculeatus</i> L.	Whole digestive contents	872
<i>Lacerta viridis</i> Laur.	Gastric juice	1,200
<i>Testudo hermanni</i> J. F. Gmelin	Gastric juice	0
	Intestinal contents	0
<i>Passer domesticus</i> L.	Gastric juice	180
<i>Turdus merula</i> L.	Gastric juice	1,560
<i>Oryzolagus cuniculus</i> L.	Gastric contents	0
	Intestinal contents	2.5*
	Caecal contents	0

\* Non-significant result.

Table 2. DISTRIBUTION AND LOCALIZATION OF CHITINASE AND CHITOBIASE IN DIGESTIVE GLANDS AND TISSUES OF SOME VERTEBRATES

Species	Organs	Activity = acetylglucosamine liberated, $\mu\text{gm./hr./gm. fresh tissues}$		
		Chitinolytic system*	Chitinase†	Chitobiase‡
<i>Carassius auratus</i> L.	Mucosa anterior intestine	43-88	490	440
	Mucosa posterior intestine	288	—	72
	Pancreas (+ Liver)	—	67	62
<i>Lacerta viridis</i> Laur.	Gastric mucosa	1437-1570	8,064	260-740
	Intestine	0-3§	—	45
	Pancreas	810-1350	13,120	0-160
<i>Testudo hermanni</i> J. M. Gmelin	Gastric mucosa	0	0	45
	Intestinal mucosa	0	0	57
<i>Turdus merula</i> L.	Mucosa of glandular stomach	340	3,114	89
<i>Rhinolophus ferrum-equinum</i> Schreb.	Intestine (first third)	26	144	56
	Intestine (third)	13§	13§	0
	Pancreas	347	5,280	71
	Gastric mucosa	0	12§	23
<i>Oryzolagus cuniculus</i> L.	Intestine	0	0	29
	Pancreas	0	0	0
	Gastric mucosa	0	0	70
	Intestine	0	7.8§	72
	Pancreas	0	—	—

\* Liberation of acetylglucosamine from chitin, by the enzyme extract.

† Liberation of acetylglucosamine from chitin, by the enzymic extract after addition of an excess of chitobiase.

‡ Liberation of acetylglucosamine from 'depolymerized chitin', a preparation containing chitobiase and other small polymers of acetylglucosamine.

§ Non-significant results.

The results are summarized in Tables 1 and 2. Chitinolytic enzymes have been found in the digestive tract of six species of vertebrates, among the eight species so far investigated, the digestive juices and tissues of turtle and rabbit being entirely devoid of chitinases. In all these eight species, muscles, kidney and liver extracts were found to be free from chitinases, the presence of chitobiase only being evident in the livers of *Lacerta* and *Testudo*.

The following tentative conclusions may be drawn from these results:

(1) The chitinolytic enzymes have been found not only in the digestive contents, but also in glands and

in care  
Such lo  
of a ba  
(2) A  
are ins  
least o  
rabbit,  
secretio  
observa  
repres  
of a co  
among  
(3) T  
an evol  
in the g  
of the v  
faculty  
pancre  
has thi  
In c  
often v  
even in  
(4) ?  
gastric  
first k  
thesize  
very s  
1 Tracey  
2 Tracey  
3 Tracey  
4 Jenni  
5 Jenni  
6 Jenni  
7 Jenni  
8 Zech  
9 Jenni  
10 Reiss  
217  
11 Jenni  
12 Jenni  
68,

Printed

IN DIGESTIVE JUICES OF  
RATES

Species or Contents	Activity of chitinolytic system: acetylglucosamine liberated, $\mu\text{gm./hr./ml.}$
113	
814	
872	
1,200	
0	
0	
180	
1,550	
0	
2.5*	
0	

LOCALIZATION OF CHITINASE AND  
TISSUES OF SOME VERTEBRATES

Activity = acetylglucosamine liberated, $\mu\text{gm./hr./gm. fresh tissues}$	Chitinase†	Chitobiase‡
43-88	490	440
288	—	72
—	87	62
437-1570	8,064	250-740
0-3.8§	—	45
110-1350	13,120	0-180
0	0	45
0	0	57
340	3,114	89
26	144	50
13§	13§	0
347	5,280	71
0	12§	23
0	0	29
0	0	0
0	0	70
0	7.8§	72

chitin, by the enzymic extract.  
chitin, by the enzymic extract  
chitobiase.  
from 'depolymerized chitin', a  
and other small polymers of

ed in Tables I and 2.  
n found in the digestive  
rates, among the eight  
the digestive juices and  
being entirely devoid of  
eight species, muscles,  
e found to be free from  
chitobiase only being  
a and *Testudo*.  
conclusions may be drawn

es have been found not  
, but also in glands and

in carefully washed gastric or intestinal mucosa. Such localization does not support the hypothesis of a bacterial origin.

(2) All the species where chitinase has been found are insectivorous, or eat chitin-covered preys, at least occasionally. In contrast, the turtle and the rabbit, which do not possess any trace of chitinase secretion, are typical phytophagous animals. This observation, if confirmed for other species, would represent the only clear-cut case, with that of lactase, of a correlation between diet and digestive enzymes among vertebrates.

(3) The site of chitinase secretion seems to show an evolution towards a higher specialization of tissues: in the goldfish, the secretion of chitinase is a property of the whole epithelium and of pancreatic islets. This faculty is restricted to the gastric mucosa and the pancreas in *Lacerta*: in bats, the gastric mucosa alone has this property.

In contrast, the secretion of chitobiase, although often very weak, occurs in the majority of tissues, even in the turtle and in the rabbit.

(4) The cases of the pancreas of *Lacerta* and of the gastric mucosa of *Turdus* and of *Rhinolophus* are the first known examples of tissues being able to synthesize large amounts of chitinase, but secreting only very small amounts of chitobiase.

1 Tracey, M. V., *Nature*, 175, 815 (1955).  
 2 Tracey, M. V., *Second Intern. Cong. Biochem.*, Paris, 242 (1952).  
 3 Tracey, M. V., *Nature*, 187, 776 (1961).  
 4 Jeuniaux, Ch., *Mém. Soc. Roy. Entom. Belg.*, 27, 312 (1956).  
 5 Jeuniaux, Ch., *Arch. Intern. Physiol. Biochem.*, 64, 583 (1956).  
 6 Jeuniaux, Ch., *Arch. Intern. Physiol. Biochem.*, 68, 684 (1960).  
 7 Jeuniaux, Ch., *Cahiers Biol. Marine* (in the press).  
 8 Zechmeister, L., and Toth, G., *Enzymologia*, 7, 165 (1939).  
 9 Jeuniaux, Ch., *Arch. Intern. Physiol. Biochem.*, 67, 115 (1959).  
 10 Reissig, J. L., Strominger, J. L., and Leloir, L. F., *J. Biol. Chem.*, 217, 959 (1955).  
 11 Jeuniaux, Ch., *Arch. Intern. Physiol. Biochem.*, 66, 408 (1958).  
 12 Jeuniaux, Ch., and Devigne, J., *Arch. Intern. Physiol. Biochem.*, 68, 411 (1960).