

Biotic homogenization can decrease landscape-scale forest multifunctionality

Fons van der Plas^{a,b,1}, Pete Manning^{a,b}, Santiago Soliveres^a, Eric Allan^a, Michael Scherer-Lorenzen^c, Kris Verheyen^d, Christian Wirth^{e,f}, Miguel A. Zavala^g, Evy Ampoorter^d, Lander Baeten^{d,h}, Luc Barbaro^{i,j}, Jürgen Bauhus^k, Raquel Benavides^c, Adam Benneter^k, Damien Bonal^l, Olivier Bouriaud^m, Helge Bruelheide^{f,n}, Filippo Bussotti^{o,2}, Monique Carnol^p, Bastien Castagnyrol^{i,j}, Yohan Charbonnier^{i,j}, David Anthony Coomes^q, Andrea Coppi^o, Cristina C. Bestias^r, Seid Muhie Dawud^s, Hans De Wandeler^t, Timo Domisch^u, Leena Finér^u, Arthur Gessler^v, André Granier^l, Charlotte Grossiord^w, Virginie Guyot^{i,j,x}, Stephan Hättenschwiler^y, Hervé Jactel^{i,j}, Bogdan Jaroszewicz^z, François-xavier Joly^y, Tommaso Jucker^q, Julia Koricheva^{aa}, Harriet Milligan^{aa}, Sandra Mueller^c, Bart Muys^t, Diem Nguyen^{bb}, Martina Pollastrini^o, Sophia Ratcliffe^e, Karsten Raulund-Rasmussen^s, Federico Selvi^o, Jan Stenlid^{bb}, Fernando Valladares^{r,cc}, Lars Vesterdal^s, Dawid Zielinski^z, and Markus Fischer^{a,b,dd}

^aPlant Ecology Group, Institute of Plant Sciences, University of Bern, 3013 Bern, Switzerland; ^bSenckenberg Gesellschaft für Naturforschung, Biodiversity and Climate Research Centre, 60325 Frankfurt, Germany; ^cFaculty of Biology/Geobotany, University of Freiburg, 79104 Freiburg, Germany; ^dForest & Nature Lab, Department of Forest and Water Management, Ghent University, 9000 Ghent, Belgium; ^eSystematic Botany and Functional Biodiversity Study Group, University of Leipzig, 04103 Leipzig, Germany; ^fGerman Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig, Germany; ^gForest Ecology and Restoration Group, Department of Life Sciences, University de Alcalá, 28805 Alcalá de Henares, Madrid, Spain; ^hTerrestrial Ecology Unit, Department of Biology, Ghent University, 9000 Ghent, Belgium; ⁱInstitut National de la Recherche Agronomique (INRA), UMR 1202, Biodiversité Gènes et Communautés (BIOGECO), F-33610 Cestas, France; ^jUniversity of Bordeaux, UMR 1202, Biodiversité Gènes et Communautés, F-33600 Pessac, France; ^kFaculty of Environment and Natural Resources, University of Freiburg, 79085 Freiburg, Germany; ^lInstitut National de la Recherche Agronomique, UMR Ecologie et Écophysiologie Forestières, 54280 Champenoux, France; ^mFaculty of Forestry, Stefan cel Mare University of Suceava, Suceava 720229, Romania; ⁿInstitute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, 06108 Halle, Germany; ^oLaboratory of Applied and Environmental Botany, Department of Agri-Food Production and Environmental Science, University of Florence, 50144 Firenze, Italy; ^pLaboratory of Plant and Microbial Ecology, Department of Biology, Ecology, Evolution, University of Liège, 4000 Liège, Belgium; ^qForest Ecology and Conservation, Department of Plant Sciences, University of Cambridge, CB2 3EA Cambridge, United Kingdom; ^rDepartment of Biogeography and Global Change, National Museum of Natural Sciences (MNCS), Consejo Superior de Investigaciones Científicas, 28006 Madrid, Spain; ^sDepartment of Geosciences and Natural Resource Management, University of Copenhagen, 1958 Frederiksberg C, Denmark; ^tDivision of Forest, Nature, and Landscape, Department of Earth and Environmental Sciences, KU Leuven, BE-3001 Leuven, Belgium; ^uNatural Resources Institute Finland, FI-80101 Joensuu, Finland; ^vSwiss Federal Institute for Forest, Snow, and Landscape Research, CH-8903 Birmensdorf, Switzerland; ^wEarth and Environmental Sciences Division, Los Alamos National Laboratory, Los Alamos, NM 87545; ^xInstitut National de la Recherche Agronomique, UMR 1201, Dynamics and Ecology of Forest Landscapes, F-31326 Castanet-Tolosan, France; ^yCentre of Evolutionary and Functional Ecology UMR 5175–University of Montpellier–University Paul-Valéry Montpellier–École Pratique des Hautes Études, 34293 Montpellier Cedex 5, France; ^zBiałowieża Geobotanical Station, Faculty of Biology, University of Warsaw, 17-230 Białowieża, Poland; ^{aa}Ecology, Evolution and Behaviour, School of Biological Sciences, Royal Holloway University of London, TW20 0EX Egham, Surrey, United Kingdom; ^{bb}Department of Forest Mycology and Plant Pathology, Swedish University of Agricultural Sciences, SE-750 07, Uppsala, Sweden; ^{cc}Departamento de Biología y Geología, Escuela Superior de Ciencias Experimentales y Tecnología, Universidad Rey Juan Carlos, 28933 Móstoles, Spain; and ^{dd}Oeschger Centre for Climate Change Research, University of Bern, CH-3012 Bern, Switzerland

Edited by William H. Schlesinger, Cary Institute of Ecosystem Studies, Millbrook, NY, and approved January 29, 2016 (received for review September 8, 2015)

Many experiments have shown that local biodiversity loss impairs the ability of ecosystems to maintain multiple ecosystem functions at high levels (multifunctionality). In contrast, the role of biodiversity in driving ecosystem multifunctionality at landscape scales remains unresolved. We used a comprehensive pan-European dataset, including 16 ecosystem functions measured in 209 forest plots across six European countries, and performed simulations to investigate how local plot-scale richness of tree species (α -diversity) and their turnover between plots (β -diversity) are related to landscape-scale multifunctionality. After accounting for variation in environmental conditions, we found that relationships between α -diversity and landscape-scale multifunctionality varied from positive to negative depending on the multifunctionality metric used. In contrast, when significant, relationships between β -diversity and landscape-scale multifunctionality were always positive, because a high spatial turnover in species composition was closely related to a high spatial turnover in functions that were supported at high levels. Our findings have major implications for forest management and indicate that biotic homogenization can have previously unrecognized and negative consequences for large-scale ecosystem multifunctionality.

β -diversity | biodiversity | ecosystem functioning | FunDivEUROPE | spatial scale

It is widely established that high local-scale biodiversity increases levels of individual ecosystem functions in experimental ecosystems (1–4), and that biodiversity is even more important for the simultaneous maintenance of multiple functions at high levels (i.e., ecosystem multifunctionality) (5–8). Because the capacity of natural ecosystems to maintain multiple functions and services is crucial for human well-being (9), the positive

Significance

Numerous studies have demonstrated the importance of biodiversity in maintaining multiple ecosystem functions and services (multifunctionality) at local spatial scales, but it is unknown whether similar relationships are found at larger spatial scales in real-world landscapes. Here, we show, for the first time to our knowledge, that biodiversity can also be important for multifunctionality at larger spatial scales in European forest landscapes. Both high local (α -) diversity and a high turnover in species composition between locations (high β -diversity) were found to be potentially important drivers of ecosystem multifunctionality. Our study provides evidence that it is important to conserve the landscape-scale biodiversity that is being eroded by biotic homogenization if ecosystem multifunctionality is to be maintained.

Author contributions: F.v.d.P., P.M., S.S., E. Allan, M.S.-L., K.V., C.W., M.A.Z., and M.F. designed research; F.v.d.P., E. Ampoorter, L. Baeten, L. Barbaro, J.B., R.B., A.B., D.B., O.B., H.B., F.B., M.C., B.C., Y.C., D.A.C., A.C., C.C.B., S.M.D., H.D.W., T.D., L.F., A. Gessler, A. Granier, C.G., V.G., S.H., H.J., B.J., F.-x.J., T.J., J.K., H.M., S.M., B.M., D.N., M.P., S.R., K.R.-R., F.S., J.S., F.V., L.V., and D.Z. performed research; F.v.d.P. and S.S. analyzed data; F.v.d.P. and P.M. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: Data on forest plot locations, environmental variables, tree diversity data, and values for all ecosystem functions are available at: https://figshare.com/articles/PNAS_data_on_ecosystem_functions_tree_communities_and_multifunctionality/3082180. German National forest inventory data is available at: forestportal.efi.int/view.php?id=201&pl=01.20.

¹To whom correspondence should be addressed. Email: fonsvanderplas@gmail.com.

²Present address: Department of Biology, University of Florence, 50121 Florence, Italy.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1517903113/-DCSupplemental.

diversity–multifunctionality relationship is often used as an argument to promote biodiversity conservation (6, 10). However, although society seeks to maximize the delivery of potentially conflicting ecosystem services, such as food production, bioenergy generation, and carbon storage at the landscape scale (11–13), research into the relationship between biodiversity and ecosystem multifunctionality has been largely limited to local-scale studies, where diversity is manipulated in experimental plant communities. Although some studies have focused on more natural communities distributed over larger spatial extents (e.g., 14–16), they examined relationships between local-scale biodiversity and local-scale multifunctionality. The only previous study to investigate multifunctionality at larger scales (17) simulated artificial landscapes using data from experimental grassland communities. It showed that although different aspects of biodiversity affected multifunctionality, local-scale (α -) diversity was a much stronger driver than the turnover of species between sites (β -diversity). However, whether those findings can be extrapolated to real-world (i.e., natural, seminatural) ecosystems, such as forests, is unknown. As a result, we have a poor understanding of how multifunctionality relates to biodiversity at the larger spatial scales that are most relevant to ecosystem managers. This question is of particular concern, given recent findings suggesting that human-driven homogenization of communities [loss of β -diversity (18–21)] may be just as widespread as α -diversity declines (22, 23).


Multifunctionality can be measured by a variety of methods, and the most appropriate means of doing so remains unresolved (24–27), particularly at larger scales, where the desired distribution of ecosystem function across the landscape has not been quantified. At local scales, one can quantify ecosystem multifunctionality as the number of ecosystem functions that exceed a given threshold value, where the threshold equals a certain percentage of the maximum observed value of each function (10, 24) (hereafter “threshold-based multifunctionality”; Fig. 1*B*). This threshold reflects the minimum value of ecosystem functioning that is deemed satisfactory. Because trade-offs between ecosystem functions or services are commonplace (5, 7, 28, 29), it is often impossible to maximize all of the desired functions in a local community (6). However, when different species provide different functions (5, 7) at larger spatial scales, a high spatial turnover in community composition (i.e., a high β -diversity) across the landscape can cause different parts of the landscape to provide different functions at high levels (defined as high threshold-based β -multifunctionality; Fig. 1*B*). Therefore, high β -diversity might cause all desired ecosystem functions to be provided at high levels in at least one patch within a landscape [and hence promote threshold-based landscape-scale or γ -multifunctionality (30)] (Fig. 1*B*), but only if

(i) species differ in the functions they support and (ii) there is no “superspecies” that supports the majority of functions. This threshold-based γ -multifunctionality may be relevant for cases where forest landscapes are managed for many different services (e.g., timber production, limitation of nutrient runoff, ecotourism), but where each of these services only needs to be provided at high levels in a part of the landscape, not everywhere (31). Alternatively, a manager may seek to promote the total delivery of many summed individual ecosystem functions across a landscape. We define this total delivery as sum-based γ -multifunctionality (Fig. 1*B*). This metric may be a more appropriate measure of multifunctionality in cases where the benefits of ecosystem services are manifested at large scales, such as carbon sequestration or water purification (32). In this case, β -diversity might only promote sum-based γ -multifunctionality if nonadditive diversity effects, such as resource partitioning, species–environment matching, or spillover effects, operate at relatively large spatial scales (33, 34). It is therefore likely that the importance of β -diversity for γ -multifunctionality varies depending on the desired pattern of ecosystem service provision.

Forests provide many ecosystem services, including wood production, regulation of water quality and climate, and recreation (35, 36). Most present-day European forests and almost all forest plantations worldwide are dominated by only one or a few tree species (15, 37), although their diversity could be promoted relatively easily by planting more species or by encouraging natural regeneration. This fact makes the understanding of diversity–multifunctionality relationships in these ecosystems highly relevant for forest management.

We therefore assessed the importance of α - and β -diversity of tree species in driving γ -multifunctionality in mature European forests. To do so, we used data taken from a pan-European forest dataset consisting of 209 forest plots, specifically selected to investigate relationships between tree diversity and ecosystem functioning by maximizing variation in dominant “target” species richness and minimizing (i) variation in other potential drivers of ecosystem function (e.g., soil and climatic conditions) and (ii) covariation between tree α -diversity, species composition, and environmental variables as much as possible (38). Our plot selection therefore aimed to mimic biodiversity experiments to investigate relationships between biodiversity and ecosystem functioning in mature forests, which are difficult to undertake with manipulative approaches due to the longevity of tree species. The plots were widely distributed across six European countries, spanning boreal to Mediterranean zones and representing six major European forest types (38). In each plot, 16 ecosystem processes, functions, or properties (termed “functions” hereafter) were measured. These functions represented a wide range of

Fig. 1. Quantifying biodiversity and multifunctionality across spatial scales. The light yellow areas represent hypothetical landscapes, consisting of (white) local communities. In these communities, some species are present (colored icons in *A* and *C*), whereas others are absent (gray icons). Similarly, some functions are performing above a hypothetical threshold of 0.5 (colored icons in *B*), whereas others are not (gray icons). Diversity and threshold-based multifunctionality are quantified at (i) the local plot (α -) scale as the number of species present (two and three in *A*) or functions performing above a given threshold (two and three in *B*), (ii) the β -scale as the turnover in species composition [$=1 - \log((a + b + 2c)/(a + b + c)) = 1 - \log((1 + 2 + 2)/(1 + 2 + 1)) = 0.90$ in *A* (49)] or functions [$=1 - \log((a + b + 2c)/(a + b + c)) = 1 - \log((1 + 2 + 2)/(1 + 2 + 1)) = 0.90$ in *B* (49)] across plots, and (iii) the landscape (γ -) scale as the number of functions (four in *B*) present in at least one plot. Sum-based γ -multifunctionality is defined as the sum of all standardized ecosystem values in a landscape ($= 0.8 + 0.2 + 0.7 + 0.4 + 0.9 + 1.0 + 0.1 + 0.6 = 4.7$). In contrast to threshold-based multifunctionality, sum-based multifunctionality is not analogous to biodiversity (where species are either present or absent), and can therefore not be partitioned into α - or β -components. (*C*) This framework allows investigation of whether γ -multifunctionality is promoted by α - and/or β -diversity.






Fig. 2. Scale-dependent effects of biodiversity on forest ecosystem multifunctionality. Bars represent the standardized regression coefficients of α -diversity (light gray) and β -diversity (dark gray) in generalized LMMs explaining α - (A), β - (B), or γ - (C) multifunctionality. Multifunctionality was quantified at different scales using a threshold approach, with thresholds of 40%, 50%, 70%, and 90%. In addition, sum-based γ -multifunctionality was calculated as the sum of scaled (between 0 and 1) individual function values. Diversity measures were calculated based on individuals of target tree species. * $P < 0.05$; *** $P < 0.001$; **** $P < 0.0001$. NS, not significant.

supporting, provisioning, regulating, and cultural ecosystem services (*sensu* 9) (SI Appendix, Table S3). Next, we created simulated landscapes by randomly drawing plots from a country to generate a “landscape” of five plots, from which γ -multifunctionality was calculated. We then explored relationships between α - and β -diversity and different measures of γ -multifunctionality: threshold-based γ -multifunctionality, quantified as the number of functions with levels above a threshold [a certain percentage of maximum functioning observed across all plots (10)] in at least one plot of the landscape (quantification is shown in Fig. 1B), and sum-based γ -multifunctionality, quantified as the sum of scaled values of all functions across all plots within a landscape (quantification is shown in Fig. 1B). To demonstrate how α - and β -diversity can promote threshold-based γ -multifunctionality, we also measured the relationships between both α - and β -diversity and threshold-based α - and β -multifunctionality (quantification is shown in Fig. 1B).

Results and Discussion

Our analyses show that relationships between α -diversity and threshold-based γ -multifunctionality varied from positive, when moderate levels of ecosystem functioning were desired (40–70% thresholds), to negative, when very high levels (90% threshold) of ecosystem functioning were required (Fig. 2C; all $P < 0.05$). In contrast, relationships between β -diversity and threshold-based γ -multifunctionality were, when significant, always positive, irrespective of the level of functioning desired (Fig. 2C; all $P < 0.05$). These positive relationships with β -diversity were generally consistent throughout countries (Fig. 3C) and largely independent of whether diversity was measured as total species richness or richness of abundant target species (SI Appendix, Fig. S3 and Tables S5–S7) and the statistical approach used to investigate diversity–multifunctionality relationships (SI Appendix, Figs. S5–S9). Thus, landscapes with a high spatial turnover in species composition had consistently more functions at high levels in at least some plots than more biotically homogeneous landscapes. This finding indicates that biotic homogenization can have detrimental consequences for threshold-based γ -multifunctionality of ecosystems, whereas management that promotes a higher spatial turnover in species composition may reverse these detrimental effects. In contrast, sum-based γ -multifunctionality was related to neither α - nor β -diversity (Fig. 2C and SI Appendix, Fig. S3).

Next, we investigated the mechanisms by which α - and β -diversity may affect threshold-based γ -multifunctionality by investigating relationships between α - and β -diversity and threshold-based α - and β -multifunctionality (i.e., local multifunctionality and turnover in functioning across plots; Fig. 1B). These analyses showed that the aforementioned relationships between α -diversity and threshold-based γ -multifunctionality were mediated by effects on threshold-based α -multifunctionality: α -Diversity was positively related to threshold-based α -multifunctionality when moderate levels (40–50%) of functioning were desired [similar to most experimental

studies (8)], but negatively related when high levels (90%) of functioning were required (Fig. 2A; all $P < 0.001$), a finding that was largely consistent throughout countries (Fig. 3A) and largely independent of whether the richness of the dominant species or the richness of all species (SI Appendix, Fig. S3) was used as an α -diversity measure. This pattern may have been caused by “statistical averaging” effects similar to the portfolio effects that drive diversity–stability relationships (39): Without strong selection or complementarity effects (40), mixed species plots will tend to have intermediate, but never extremely high or low, ecosystem function levels due to the averaging of individual species effects on function. In line with this mechanism, α -diversity did not have significant effects on sum-based γ -multifunctionality (Fig. 2C). These results suggest that although function values were, on average, not higher or lower in diverse communities than in monocultures, they tended to be less extreme (never extremely high or low) (41). This result contrasts with the results of other studies focusing on more diverse and/or experimental grassland, aquatic, or soil communities. In these studies, higher α -diversity enhances α -multifunctionality even at very high thresholds (8, 24), possibly due to strong complementarity effects. However, the diversity–ecosystem functioning literature has tended to concentrate on particular ecosystems and study designs. As a result, it is difficult to infer whether these contrasting results are caused by biological or methodological differences. In any case, our results indicate that in European forests at least, the relationship between α -diversity and both α - and γ -multifunctionality strongly depends on the desired level of functioning.

In our next analysis, we investigated the relationship between β -diversity and γ -multifunctionality. In contrast to α -diversity, the positive relationship between β -diversity and threshold-based γ -multifunctionality was independent of the desired level of ecosystem functioning. In almost all countries (Figs. 2B and 3C), and irrespective of whether target or total species richness was the diversity metric used (SI Appendix, Fig. S3), β -diversity was positively related to threshold-based β -multifunctionality when moderate or high levels of ecosystem functioning were desired, thereby increasing the number of functions that were provided at high levels in at least one part of the landscape (threshold-based γ -multifunctionality; Fig. 2B). However, we did not detect a significant relationship between β -diversity and sum-based γ -multifunctionality (Fig. 2B). This finding was likely due to trade-offs between ecosystem functions: Of the 120 possible pairwise correlations among functions, 50 were negative. This result made it impossible to




Fig. 3. Scale-dependent effects of biodiversity on forest ecosystem multifunctionality are generally consistent across countries. Bars represent the standardized regression coefficients of α -diversity (blue) and β -diversity (red) in LMMs explaining α - (A), β - (B), or γ - (C) multifunctionality. Multifunctionality was quantified at different scales using a threshold approach, with thresholds of 40%, 50%, 70%, and 90%. In addition, sum-based γ -multifunctionality was calculated as the sum of scaled (between 0 and 1) individual function values. Diversity measures were calculated based on individuals of target tree species.

achieve very high levels of all functions across the entire landscape. These results thus indicate that although β -diversity is not related to higher average levels of ecosystem functions, it is positively related to the number of functions that perform at high levels in at least part of the landscape. Hence, positive relationships between β -diversity and threshold-based γ -multifunctionality (and hence threshold-based γ -multifunctionality) are caused by the fact that different species support different functions (5, 7). For example, in Polish forests, monoculture plots of the conifer *Picea abies* are related to high levels of many functions relating to the production of quality timber (e.g., timber quality, biomass production), whereas plots of the deciduous tree *Carpinus betulus* were of higher recreational and conservation value due to a high diversity of bats and understory plants (*SI Appendix, Table S8*). Hence, forest landscapes where some locations were dominated by *P. abies* and others by *C. betulus* provided more functions at high levels than those forest landscapes where all plots had the same tree species composition.

Our finding that the relationship between biodiversity and European forest γ -multifunctionality depends strongly on the way that multifunctionality is quantified has important implications for European forest management. In short, our results suggest that different patterns of tree species distribution would achieve different management goals (or “landscape multifunctionality scenarios”). The results of the threshold-based γ -multifunctionality analysis would be most relevant to situations where managers sought to promote forest landscapes with very high levels of ecosystem functioning in at least some (but not necessarily all) local patches (*sensu* 30). As described earlier, this situation may occur when managers seek to provide different ecosystem services in different localities. An example of such a landscape is one where some localities provide recreation or cultural services, such as aesthetic beauty and a diversity of charismatic taxa (31) (*SI Appendix, section S3*), whereas other localities maximize provisioning services that are only cost-effective when delivered at very high levels, such as production of high-value timbers (42), or form hotspots of certain biogeochemical functions that need to be strategically located, such as the minimization of nutrient runoff close to water bodies. In such scenarios, threshold-based γ -multifunctionality could be promoted in forest landscapes that possess a high turnover in community composition, but a low α -diversity (top left landscape in Fig. 1C), by promoting a range of different monocultures across the landscape. When the delivery of provisioning services is cost-effective at lower levels or when cultural or regulating services do not need to be at extremely high levels, one could aim to promote landscapes with moderate levels of many functions (40% or 50% threshold γ -multifunctionality). In this scenario, threshold-based γ -multifunctionality is highest in forests with both a high spatial turnover in community composition and a high local diversity of tree species (top right landscape in Fig. 1C). This finding is in line with Gamfeldt et al. (15), who hypothesized, based on local-scale analyses, that “adjacent stands, each with multiple species but in different combinations, might be the best way to provide multiple ecosystem services at the landscape scale.” A third hypothetical scenario would be to maximize total delivery of services across the landscape (high sum-based γ -multifunctionality), rather than having highly localized specialist patches that deliver a limited number of services at very high levels. This scenario may be most relevant to cases where the primary goal of ecosystem management is to provide ecosystem services whose benefits are manifested at large scales, such as carbon sequestration (43). We found that neither α -diversity nor spatial turnover in community composition (β -diversity) had significant detectable relationships with γ -multifunctionality under this scenario. In summary, we demonstrate that the importance of different components of diversity for promoting γ -multifunctionality is likely to depend on management goals. Accordingly, stakeholder engagement is required to see where these situations apply in real forested landscapes. Further studies are also required to confirm that tree α - and β -diversity are causal drivers of the observed relationships and to see how important they are in comparison to

other potentially important factors in driving ecosystem multifunctionality in representative European forests.

In this study, some of the benefits of biodiversity for γ -multifunctionality may have been underestimated. Our study did not consider some of the spatiotemporal processes that occur in real forest landscapes [e.g., dispersal and movement of ecosystem service providers, species-environment matching, large-scale resource partitioning, spillover and subsidy of ecosystem services between neighboring patches (13, 44)]. These processes could promote ecosystem functioning in landscapes that possess a high spatial turnover in species composition even more than was detected here. For example, a forest resistant to herbivory might also reduce pest damage in adjacent forests by lowering populations of herbivores and preventing their movement into more vulnerable areas, thus strengthening the relationship between β -diversity and forest γ -multifunctionality. Future studies could explore these ideas by studying ecosystem multifunctionality in landscapes where ecological interactions between patches of differing diversity and composition are quantified.

Previous studies have demonstrated that local-scale (α -) biodiversity can boost multifunctionality in the real world, in addition to experimental ecosystems (8, 14–16). Here, we add evidence that both α - and β -diversity can also drive ecosystem multifunctionality at the landscape scale, and that the desired distribution of ecosystem functions across the landscape influences the importance of this relationship. Biotic homogenization is occurring worldwide at local, regional, and global scales (19–21). Similarly, current forest management often results in large areas of low species turnover. Our study is an important step forward in exploring the importance of this biotic homogenization for γ -multifunctionality. It shows that biotic homogenization may have negative, strong, far-reaching, and so far overlooked impacts upon the ecosystem services on which humanity depends, and that these impacts may be as strong as, or even stronger than, the impacts of local diversity loss.

Methods

Plot Selection. In total, 209 forest plots (each measuring 30 × 30 m) were established within the European FunDivEUROPE project (fundiveuropekt.boku.ac.at). Because we were interested in the effects of tree species diversity on ecosystem functioning in mature forests (38), plot selection was aimed at mimicking the design of a biodiversity experiment, in which variation in environment is minimized and diversity is not confounded with composition, as in most observational studies of diversity. Hence, the design aimed to bridge the gap between controlled but very young tree diversity experiments and observational studies, where diversity can be strongly confounded with other factors.

Plots were located in six European countries, ranging from boreal to Mediterranean zones, and with each representing a major European forest type (38): Finland (28 plots, boreal forest), Poland (43 plots, hemiboreal forest), Germany (38 plots, temperate deciduous forest), Romania (28 plots, mountainous deciduous forest), Italy (36 plots, thermophilous deciduous forest), and Spain (36 plots, Mediterranean mixed forest) (*SI Appendix, Fig. S1*). Within countries, plots were located in a single region ranging in size from 5 × 5 km (Romania) to 150 × 150 km (Finland). In each country, between three and five regionally common target species were selected, 15 in total (*SI Appendix, Table S1*). Plots were then selected to differ as much as possible in richness of target species and so that almost all possible combinations of these target species were realized, a design that emulates the designs of biodiversity experiments (38). Richness levels of one, two, three, four, and five target species were replicated 56, 67, 54, 29, and 3 times, respectively, across countries, and most possible target species compositions were realized [additional details on the selection procedure are provided in the study by Baeten et al. (38)]. To achieve this goal, some admixture of nontarget species was unavoidable. However, on average, target species accounted for 93.75% of the individuals and 91.39% of the basal area, and they were always represented by more than two individuals (*SI Appendix, Fig. S2*). We therefore focus on using the richness of target species in our analyses, but we also tested for the effect of total (target + admixed) species richness (*SI Appendix*). Plot selection strictly avoided correlations between tree species richness and soil factors, as well as any spatial autocorrelation in diversity (38), by choosing plots that differed as little as possible in environmental factors (soil texture, depth, pH, and altitude) that could potentially confound diversity effects on multifunctionality. The diversity gradient

environmental factors as covariates in the LMMs. We performed three different LMM analyses. In the first, we investigated how α -multifunctionality was driven by α -diversity, using an LMM with α -multifunctionality as the response variable, species richness as the focal fixed factor, species evenness, proportion of evergreen trees, altitude, soil depth, soil pH, soil sand and clay content, and their two-way interactions as covariates, and with country as a random factor. In the second analysis, we investigated the relationship between β -multifunctionality and β -diversity, using an LMM with β -multifunctionality as the response variable, β -diversity as the focal fixed factor, environmental heterogeneity as a covariate, and country as a random factor. In the third analysis, we investigated how γ -multifunctionality was affected by both α - and β -diversity, by first constructing a full LMM with γ -multifunctionality as the response variable; α - and β -diversity as the focal fixed factors; species evenness, proportion of evergreen trees, altitude, soil depth, soil pH, soil sand and clay content, environmental heterogeneity, and their bivariate interactions as covariates; and country as a random factor. As a result of the careful study design, diversity measures were largely independent of the covariates (38): Correlations between focal predictors and covariates were always <0.230 , whereas the correlation between α - and β -diversity was moderate ($R^2 = 0.316$, $P < 0.0001$); hence, there was no strong indication of multicollinearity. In all three analyses, we used a backward model-selection analysis to remove covariates sequentially (based on ratio-likelihood tests with a Bonferroni correction) until we reached a final model with only the focal fixed factor(s) and significant covariates. From this final model, we quantified the significance of α - and

β -diversity in driving multifunctionality using likelihood ratio tests, and we also quantified their standardized regression coefficients. All analyses were performed for all different threshold-based and sum-based multifunctionality variables. As a robustness check, we also repeated all these analyses with predictors and covariates based on all tree species, rather than on target species only (SI Appendix). In addition, to investigate how general the main patterns were across countries, we ran linear models for each country separately, with the same fixed factors in models as in the finally selected LMMs. All analyses were performed using R version 3.0.2 (51). LMMs were fitted using the "lmer" function of the "lme4" library (52). Given that effects of covariates were variable and complex, and that the main focus of this study was on biodiversity effects, the effects of covariates are not presented here (effect sizes of all covariates are shown in SI Appendix, Tables S2–S4).

In addition to LMMs, we used structural equation models to investigate relationships between biodiversity and multifunctionality, to test the sensitivity of our results to the statistical method used, and to test for indirect relationships between biodiversity and multifunctionality (SI Appendix, section S4).

ACKNOWLEDGMENTS. We thank the Hainich National Park administration as well as Felix Berthold and Carsten Beinhoff for support of this study and Gerald Kaendler and the Johann Heinrich von Thünen-Institut for providing access to the German National Forest Inventory data. The research leading to these results received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under Grant Agreement 265171.

- Hooper DU, et al. (2005) Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecol Monogr* 75(1):3–35.
- Balvanera P, et al. (2006) Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecol Lett* 9(10):1146–1156.
- Cardinale BJ, et al. (2011) The functional role of producer diversity in ecosystems. *Am J Bot* 98(3):572–592.
- Handa IT, et al. (2014) Consequences of biodiversity loss for litter decomposition across biomes. *Nature* 509(7499):218–221.
- Hector A, Bagchi R (2007) Biodiversity and ecosystem multifunctionality. *Nature* 448(7150):188–190.
- Zavaleta ES, Pasari JR, Hulvey KB, Tilman GD (2010) Sustaining multiple ecosystem functions in grassland communities requires higher biodiversity. *Proc Natl Acad Sci USA* 107(4):1443–1446.
- Isbell F, et al. (2011) High plant diversity is needed to maintain ecosystem services. *Nature* 477(7363):199–202.
- Lefcheck JS, et al. (2015) Biodiversity enhances ecosystem multifunctionality across trophic levels and habitats. *Nat Commun* 6:6936.
- Millennium Ecosystem Assessment (2005) *Ecosystems and Human Well-Being: Synthesis* (Island Press, Washington, DC).
- Garnfeldt L, Hillebrand H, Jonsson PR (2008) Multiple functions increase the importance of biodiversity for overall ecosystem functioning. *Ecology* 89(5):1223–1231.
- Green RE, Cornell SJ, Scharlemann JPW, Balmford A (2005) Farming and the fate of wild nature. *Science* 307(5709):550–555.
- Goldstein JH, et al. (2012) Integrating ecosystem-service tradeoffs into land-use decisions. *Proc Natl Acad Sci USA* 109(19):7565–7570.
- Manning P, Taylor G, Hanley ME (2015) Bioenergy, food production and biodiversity—an unlikely alliance? *Glob Change Biol Bioenergy* 7(4):570–576.
- Maestre FT, et al. (2012) Plant species richness and ecosystem multifunctionality in global drylands. *Science* 335(6065):214–218.
- Garnfeldt L, et al. (2013) Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat Commun* 4:1340.
- Allan E, et al. (2015) Land use intensification alters ecosystem multifunctionality via loss of biodiversity and changes to functional composition. *Ecol Lett* 18(8):834–843.
- Pasari JR, Levi T, Zavaleta ES, Tilman D (2013) Several scales of biodiversity affect ecosystem multifunctionality. *Proc Natl Acad Sci USA* 110(25):10219–10222.
- McKinney ML, Lockwood JL (1999) Biotic homogenization: A few winners replacing many losers in the next mass extinction. *Trends Ecol Evol* 14(11):450–453.
- Dornelas M, et al. (2014) Assemblage time series reveal biodiversity change but not systematic loss. *Science* 344(6181):296–299.
- Vellend M, et al. (2013) Global meta-analysis reveals no net change in local-scale plant biodiversity over time. *Proc Natl Acad Sci USA* 110(48):19456–19459.
- McGill BJ, Dornelas M, Gotelli NJ, Magurran AE (2015) Fifteen forms of biodiversity trend in the Anthropocene. *Trends Ecol Evol* 30(2):104–113.
- Murphy GEP, Romanuk TN (2014) A meta-analysis of declines in local species richness from human disturbances. *Ecol Evol* 4(1):91–103.
- Newbold T, et al. (2015) Global effects of land use on local terrestrial biodiversity. *Nature* 520(7545):45–50.
- Byrnes JEK, et al. (2014) Investigating the relationship between biodiversity and ecosystem multifunctionality: Challenges and solutions. *Methods Ecol Evol* 5(2):111–124.
- Bradford MA, et al. (2014) Discontinuity in the responses of ecosystem processes and multifunctionality to altered soil community composition. *Proc Natl Acad Sci USA* 111(40):14478–14483.
- Byrnes J, et al. (2014) Multifunctionality does not imply that all functions are positively correlated. *Proc Natl Acad Sci USA* 111(51):E5490.
- Bradford MA, et al. (2014) Reply to Byrnes et al.: Aggregation can obscure understanding of ecosystem multifunctionality. *Proc Natl Acad Sci USA* 111(51):E5491. (lett).
- Lavorel S, et al. (2011) Using plant functional traits to understand the landscape distribution of multiple ecosystem services. *J Ecol* 99(1):135–147.
- Grigulis K, et al. (2013) Relative contributions of plant traits and soil microbial properties to mountain grassland ecosystem services. *J Ecol* 101(1):47–57.
- Brandt J (2003) Multifunctional landscapes—perspectives for the future. *J Environ Sci (China)* 15(2):187–192.
- Chan KMA, Shaw MR, Cameron DR, Underwood EC, Daily GC (2006) Conservation planning for ecosystem services. *PLoS Biol* 4(11):e379.
- Trumper K, et al. (2009) *The Natural Fix? The Role of Ecosystems in Climate Mitigation. A UNEP Rapid Response Assessment. United Nations Environment Programme* (United Nations Environment Programme-World Conservation Monitoring Centre, Cambridge, UK).
- Loreau M, Mouquet N, Gonzalez A (2003) Biodiversity as spatial insurance in heterogeneous landscapes. *Proc Natl Acad Sci USA* 100(22):12765–12770.
- Cardinale BJ, Ives AR, Inchausti P (2004) Effects of species diversity on the primary productivity of ecosystems: Extending our spatial and temporal scales of inference. *Oikos* 104(3):437–450.
- Aerts R, Honnay O (2011) Forest restoration, biodiversity and ecosystem functioning. *BMC Ecol* 11:29.
- Fares S, Mugnozza GS, Corona P, Palahi M (2015) Sustainability: Five steps for managing Europe's forests. *Nature* 519(7544):407–409.
- Bauhus J, van der Meer P, Kanninen M (2010) *Ecosystem Goods and Services from Plantation Forests* (Earthscan, London).
- Baeten L, et al. (2013) A novel comparative research platform designed to determine the functional significance of tree species diversity in European forests. *Perspect Plant Ecol Syst* 15(5):281–291.
- Tilman D, Lehman CL, Bristow CE (1998) Diversity-stability relationships: Statistical inevitability or ecological consequence? *Am Nat* 151(3):277–282.
- Loreau M, Hector A (2001) Partitioning selection and complementarity in biodiversity experiments. *Nature* 412(6842):72–76.
- Loreau M (1998) Biodiversity and ecosystem functioning: A mechanistic model. *Proc Natl Acad Sci USA* 95(10):5632–5636.
- Phalan B, Onial M, Balmford A, Green RE (2011) Reconciling food production and biodiversity conservation: Land sharing and land sparing compared. *Science* 333(6047):1289–1291.
- Canadell JG, Raupach MR (2008) Managing forests for climate change mitigation. *Science* 320(5882):1456–1457.
- Tscharntke T, Klein AM, Kruss A, Steffan-Dewenter I, Thies C (2005) Landscape perspectives on agricultural intensification and biodiversity–ecosystem service management. *Ecol Lett* 8(8):857–874.
- Pielou EC (1966) The measurement of diversity in different types of biological collections. *J Theor Biol* 13:131–144.
- Jucker T, Bouriaud O, Avacaritei D, Coomes DA (2014) Stabilizing effects of diversity on aboveground wood production in forest ecosystems: Linking patterns and processes. *Ecol Lett* 17(12):1560–1569.
- White RE (2005) *Principles and Practice of Soil Science: The Soil as a Natural Resource* (Wiley-Blackwell, Oxford, UK), 4th Ed.
- Whittaker RH (1960) Vegetation of the Siskiyou Mountains, Oregon and California. *Ecol Monogr* 30(3):279–338.
- Lennon JJ, Koleff P, Greenwood JJD, Gaston KJ (2001) The geographical structure of British bird distributions: Diversity, spatial turnover and scale. *J Anim Ecol* 70(6):966–979.
- Wang S, Loreau M (2014) Ecosystem stability in space: α , β and γ variability. *Ecol Lett* 17(8):891–901.
- R Core Team (2013) *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria).
- Bates D, Maechler M, Bolker B, Walker S (2014) lme4: Linear mixed-effects models using Eigen and S4, R package version 1.1-7. Available at cran.r-project.org/web/packages/lme4/index.html. Accessed February 16, 2015.