



## **Thermal maturation of carbonaceous material from Mbuji-Mayi Supergroup (Kasai, Democratic Republic of Congo).**

Blaise K. Baludikay (1), Jean-Yves Storme (1), Daniel Baudet (2), Camille François (1), and Emmanuelle Javaux (1)

(1) UR GEOLOGY, Department of Geology, University of Liege, Liège, Belgium. [bkbaludikay@ulg.ac.be](mailto:bkbaludikay@ulg.ac.be), (2) Geodynamics & Mineral Resources Service, Royal Museum for Central Africa, Tervuren, Belgium

Thermal maturation of carbonaceous material from Mbuji-Mayi Supergroup (Kasai, Democratic Republic of Congo).

B.K. Baludikay<sup>1\*</sup>, J.-Y. Storme<sup>1</sup>, D. Baudet<sup>2</sup>, C. François<sup>1</sup> and E.J. Javaux<sup>1</sup>

<sup>1</sup>Palaeobiogeology-Palaeobotany-Palaeopalynology, UR GEOLOGY, Department of Geology, University of Liege, Quartier Agora, Allée du 6 Août, 14, B18, B-4000 Liège 1, Belgium. [bkbaludikay@ulg.ac.be](mailto:bkbaludikay@ulg.ac.be)

<sup>2</sup> Geodynamics & Mineral Resources Service, Royal Museum for Central Africa, 3080 Tervuren, Belgium.

The Mbuji-Mayi Supergroup is a sedimentary sequence in DRC unaffected by regional metamorphism. It consists of two distinct successions: a lower, ~500 m thick siliciclastic sequence of the BI Group and an upper, ~1000 m thick carbonate sequence with stromatolitic build-ups and black shales of the BII Group directly overlain by basaltic lavas [1]. Radiometric data suggest a Latest Meso- to Early Neoproterozoic age [2, 3, 4, and 5]. A well preserved and diversified microfossil assemblage is reported including 54 taxa belonging to 32 genera. The potential Late Mesoproterozoic-Tonian index fossil *Trachyhystrichosphaera aimika*, is reported for the first time in central Africa, and co-occurs with other eukaryotes and prokaryotes [6]. Thermal maturation calculated on macerate residues, using geothermometer for low-grade metamorphism [7] reveals thermal palaeoenvironments of organic matter, ranging from 180 to 279° C (average = 249 ± 37 °C). The range of thermal maturity is similar, in both microfossils and amorphous organic matter. Raman reflectance ( $R_{mcRo}$  %), which is also an index indicative of maturity [8], ranges from 1.05 to 2.55 % (average = 2.01 ± 0.42 %). So, organic matter from Mbuji-Mayi is likely into a maturation stage corresponding to oil window.

References: [1] Raucq (1957) Ann. MRAC, série 8, Sc. géol. 18, 427. [2] Cahen & Snelling (1966) Publ. C., Amsterdam. [3] Cahen et al. (1984) Clarendon Press, Oxford. [4] Delpomdor et al. (2013) Pal. 3 389, 4–34. [5] François et al. (in preparation). [6] Baludikay et al. (in review) Prec. Res. [7] Kouketsu et al. (2014) Island Arc 23, 33–50. [8] Liu et al. (2013) Geochemistry, Chi. Sc. Bul. 58 (11), 1285-1298.