13th Congress of the European Federation of Sexology Dubrovnik, 25-28th May 2016

A study evaluating the effectiveness of bibliotherapy for premature ejaculation (BibliothEP)

Ph. Kempeneers, R. Andrianne, S. Bauwens, S. Blairy, M. Cuddy, I. Georis, Q. Longrée, J-F Pairoux (†)

University of Liege, Belgium







Treating Premature Ejaculation (PE) Clinical Trials

Sex therapy

- Demonstrated effectiveness (Cormio et al., 2015; De Carufel & Trudel, 2006; De Sutter et al., 2002; Kempeneers et al., 2014; Jern, 2013; Melnik et al., 2009, 2011; Saint Lawrence & Madakasira, 1992; Rowland, 2011)
- Limitations: few controlled studies, relative complexity of the treatments, lack of theoretical clarity, require substantial personal and financial input on the patient's part.

Drug treatments (e.g. SSRIs – Dapoxetine)

- Efficacy demontstrated by numerous controlled studies (Althof et al., 2010; McMahon et al., 2011; Porst , 2011)
- *Limitations*: Delaying ≠ controlling ejaculation, purely symptomatic action, possible unwanted side effects

Treating PE in Real Life What Surveys Say

- PE is a very common complaint, but only a small proportion of men suffering from it seek professional help (Levinson, 2008; Porst et al., 2007)
- On the other hand, men suffering from PE are likely to gather written or online information about the problem, and they attempt a number of self-help interventions themselves, often with limited success (Porst, 2012)
- Even in case of meeting with a professional, the latter is infrequently a specialist in sexual health (Kempeneers et al., 2013; Levinson, 2008; Porst et al., 2007)
- Professional meeting often *does not lead to a satisfying outcomes* (Perelman, 2006; Porst et al., 2007).
- > PE is a common problem, but insufficiently treated

Treating PE in Real Life What Surveys Say

PE is a common problem, but insufficiently treated

Possible reasons

- Feeling uncomfortable talking about the issue (Porst et al., 2007; Rowland, 2011; Symonds et al., 2003)
- Lack of awareness in the general population about effective treatments for PE (Ibid.)
- Treatments limitations
 - Sex therapy can be a complex intervention and requires a specialist practitioner. In several countries, this form of treatment is not reimbursed by social security. This intervention requires substantial input on the patient's part, often at significant financial cost, for results that are not guaranteed.
 - Pharmacological interventions can be effective in prolonging sexual intercouse, but delaying
 ejaculation is not the same as controlling it: penetration duration might remain too short... or
 become too long. Thus, sexual satisfaction is not always optimised. Moreover the pharmacological
 agents can sometimes have undesirable side effects.

A challenge:

- ➤ How to improve the affordability of efficient treatments?
- ➤ How to improve the effectiveness of first-line interventions?

➤ BibliothEP study

BibliothEP study (Bibliotherapy for PE)

<u>Aim</u>: to develop an intervention for PE that would be simple, effective, affordable and easy to access, safe, and possibly to be used without professional guidance, or at least, without support from a specialist in sex therapy.



Steps

- Hypothesis I: sex therapy for PE could be simplified by reducing it to some main active ingredients. (Kempeneers, Bauwens & De Sutter, RFCCC, 2004)
- Hypothesis II: This simplified model of treatment could be efficiently presented in a concise self-help manual (< 16,500 words in French, see cover above). (Kempeneers, Bauwens & Andrianne, De Boeck, 2015)
 - ➤ **BibliothEP-1**: bibliotherapy vs. waiting-list (Kempeneers et al., JSM, 2012)
- Hypothesis III: The bibliotherapy could optimized as a first-line treatment with brief support from a professional not specialized in sex therapy but specifically trained (5-hour training module) to facilitate the self-help process (e.g. by providing further information, providing motivational support, helping patients to adapt the material to their own situation and demonstrating some exercises).
 - ➤ **BibliothEP-2**: assisted bibliotherapy vs. bibliotherapy alone (Kempeneers et al., submitted)

Fighting against Premature Ejaculation (Practical Guide)

Kempeneers, Bauwens, & Andrianne (2015)



CONTENTS

INTRODUCTION

- I. DEFINITION
- 1. A subjective landmark: the person's wish
- 2. Minimal stimulation of the penis
- 3. Frequency of the difficulty
- 4. Frequency of sexual activities with ejaculations
- 5. Newness of the sexual experience, newness of the partner

Conclusion: the « official definition »

II. THE CAUSES OF PREMATURE EJACULATION

Preamble

- 1. In general
- 2. Special cases
- 3. In detail
- 3.1. Nature factor
- 3.2. Culture factor

III. A TWO-STAGES BEHAVIOURAL TREATMENT

- 1. To look at the coitus in perspective and to enlarge the range of erotic behaviours
- 2. To thwart the reflexes contributing to a rapid growth of excitation
- 2.1. Masturbation "fixed wrist mobile body"
- 2.2. "Abdominal" respiration
- 2.3. To synchronize the breathing with the copulating movement
- 2.4. To have breaks
- 2.5. Application in coital situations
- 2.6. In the end


VI. EPILOGUE

- 1. The advantages of the booklet
- 2. Its limits
- 3. Do efficacious medications for premature ejaculation exist?

GLOSSARY

FOR FURTHER READING

USEFUL ADDRESSES




Clinical Results: Percieved latency time

BibliothEP-1 (N = 120)


>Outcome significantly different from baseline at p<.001 either at 6 and 12 month F-up

➤ No significant change in waiting-list condition



BibliothEP-2 (N = 71)

➤ Outcome significantly different from baseline at p<.001 ➤ No significant difference between conditions (bibliotherapy + accompaniment vs. bibliotherapy alone)



Clinical Results: Control


BibliothEP-1 (N = 120)


>Outcome significantly different from baseline at p<.001 either at 6 and 12 month F-up

➤ Waiting-list condition (2 month) different from baseline at p<.05

BibliothEP-2 (N = 71)

➤ Outcome significantly different from baseline at p<.001 ➤ No significant difference between conditions (bibliotherapy + accompaniment vs. bibliotherapy alone)






Clinical Results: Sexual Satisfaction

BibliothEP-1 (N = 120)


➤ Outcome significantly different from baseline at p<.001 either at 6 and 12 month F-up

➤ No significant change in waiting-list condition



BibliothEP-2 (N = 71)

➤ Outcome significantly different from baseline at p<.001 ➤ No significant difference between conditions (bibliotherapy + accompaniment vs. bibliotherapy alone)




Clinical Results: Distress

BibliothEP-1 (N = 120)


>Outcome significantly different from baseline at p<.001 either at 6 and 12 month F-up

➤ No significant change in waiting-list condition



BibliothEP-2 (N = 71)

 \gt Outcome significantly different from baseline at p<.001 \gt More important change in bibliotherapy + accompaniment than in bibliotherapy alone condition at p<.05




Clinical Results: Distress

BibliothEP-1 (N = 120)


>Outcome significantly different from baseline at p<.001 either at 6 and 12 month F-up

➤ No significant change in waiting-list condition



BibliothEP-2 (N = 71)

➤ Outcome significantly different from baseline at p<.001 ➤ More important change in bibliotherapy + accompaniment than in bibliotherapy alone condition at p<.05



Dysfunctional sexual cognitions

(Sexual Irationality Questionnaire)

	Pre-test	Post-test
BibliothEP-1	108 (13.3)	99* (14.6)
BibliothEP-2	106 (12.54)	99* (17.15)


^{*} Significant at .001

The sexual improvement appeared to be accompanied by a better adjustment of sexual cognitions

Feeling of improvement

BibliothEP-1

> 86% at least « slightly improved »




BibliothEP-2

> Bibliotherapy alone: 86% at least « slightly improved »

➤ Bibliotherapy + accompaniment: 94% at least « sligthly imroved »

** conditions different at p<.05



Conclusions and prospects (1)

- Bibliotherapy produces significant improvements in PE problems
- Mainly if the PE is of moderate severity, but this is not exclusive
- The indication for bibliotherapy cannot be determined on the basis of a specific profile (e.g. lifelong, acquired, subjective, severe or less severe forms of PE)
- Behavioural and cognitive factors targeted in the bibliotherapy appear to be involved in several forms of PE.
- Limitations: a possible recruitment bias and important dropout rates (50-70%) question the generality of the results.
- The cost/benefit ratio of the bibliotherapy makes it an ideal tool as a first-line intervention
- Some clinical pictures have complexities that exceed the therapeutic capacities of the bibliotherapy.
- To some extent, additional therapist support can facilitate the assimilation of bibliotherapy and thus enhance the treatment effect. However, the benefit of such an accompaniment remains statistically modest.
 - Improving the outcome by increasing the amount of therapist support?
 - Ceiling effect?

Conclusions and prospects (2)

- Future research is needed to compare the relative effectiveness of bibliotherapy with pharmacotherapy and combined treatments.
- However, bibliotherapy could compete with pharmacotherapy to become the first-line intervention in severe forms of PE

Références (1)

- 1. Althof, S.E., Abdo, C.H.N., Dean, J., Hacket, G., McCabe, M., McMahon, C.G., Rosen, R.C., et al. (2010). International Society for Sexual Medicine's guidelines for the diagnosis and treatment of premature ejaculation. *J Sex Med*, 7, 2947-69.
- 2. Cormio, L., Massenio, P., La Rocca, R., Verze, P., Mirone, V., & Carrieri, G. (2015). The combination of dapoxetine and behavioral treatment provides better results than dapoxetine alone in the management of patients with lifelong premature ejaculation. *J Sex Med*, 12, 1609-15.
- 3. De Carufel, F. (2009). The Difficulty to Prolong Intercourse: Theoretical Aspects, Evaluation and Treatment. [L'éjaculation prématurée : Compréhension et traitement par la thérapie sexofonctionnelle]. Louvain-la-Neuve, B : Presses universitaires de Louvain.
- 4. De Carufel, F., & Trudel, G. (2006). Effects of a new functional-sexological treatment for premature ejaculation. *J Sex Mar Ther*, 32, 97-114.
- 5. De Sutter, P., Reynaert, N., Van Broeck, N., & De Carufel, F. (2002). Treatment of premature ejaculation by a cognitivo-behavioral bibliotherapy. [Traitement de l'éjaculation précoce par une approche bibliothérapeutique cognitivo-comportentale sexologique]. *JTCC*. 12:131-6.
- 6. Desjardins, J. (1985). L'approche sexo-corporelle : fondements théoriques et champs d'application. *Psychothérapie*, 1, 51-8.
- 7. Jern, P. (2013). Evaluation of a behavioral treatment intervention for premature ejaculation using a handheld stimulating device. *J Sex Mar Ther, 40,* 358-366.
- 8. Kempeneers, P., Andrianne, R., Bauwens, S., Georis, I., Pairoux, J.F., & Blairy, S. (2012). Clinical outcomes of a new self-help booklet for premature ejaculation. *J Sex Med*. 9, 2419-28
- 9. Kempeneers, P., Andrianne, R., Bauwens, S., Georis, I., Pairoux, J. F., & Blairy, S. (2013). Functional and psychological characteristics of Belgian men with premature ejaculation and their partners. *Arch Sex Behav, 42*(1), 51-66.
- 10. Kempeneers, P., Andrianne, R., Cuddy, M., Georis, I., Longré, Q., Pairoux, J.F., & Blairy, S. (submitted). BibliothEP-2: a comparative study of guided vs pure self-treatment for premature ejaculation. *Submitted*.
- 11. Kempeneers, P., Andrianne, R., Lequeux, A., & Blairy, S. (2014). Premature ejaculation: a review of questions. [L'éjaculation précoce : une revue de questions]. *RFCCC.* 19, 2, 35-60.
- 12. Kempeneers P, Bauwens S & Andrianne R. (2015). Fighting against premature ejaculation. [Lutter contre l'éjaculation précoce: quide pratique]. Paris, France: De Boeck-Solal.

Références (2)

- 13. Kempeneers, P., Bauwens, S., & De Sutter, P. (2004). New prospects in the treatment of premature ejaculation. [Perspectives nouvelles dans le traitement de l'éjaculation précoce]. *RFCCC*, 9, 4, 1-8.
- 14. Levinson, S. (2008). The "difficulties" of sexual function: contexts, determinants, and significations [Les « difficultés » de la fonction sexuelle : contextes, déterminants et significations]. In N. Bajos & M. Bozon (Eds.), *Enquête sur la sexualité en France : Pratiques, genre et santé* (pp. 485-508). Paris : La Découverte.
- 15. McMahon, C.G., Althof, S.E., Kaufman, J.M., Buvat, J., Levine, S.B., Aquilina, J.W. et al. (2011). Efficacy and safety of dapoxetine for the treatment of premature ejaculation: integrated analysis of results from five phase 3 trials. *J Sex Med*, 8, 524-39.
- 16. Melnik, T., Althof, S., Atallah, A.N., Puga, M.E.D.S., Glina, S., & Riera, R. (2011). Psychosocial interventions for premature ejaculation (Review). *Cochrane Database of Systematic Reviews*, 8, CD008195.
- 17. Melnik, T., Glina, S., & Rodrigues, O.M. (2009). Psychological intervention for premature ejaculation. *Nature Reviews / Urology*, 6, 501-8.
- 18. Perelman, M. A. (2006). A new combination treatment for premature ejaculation: a sex therapist perspective. *J Sex Med,* 3, 1004-1012.
- 19. Porst, H. (2011). An overview of pharmacotherapy in premature ejaculation. *J Sex Med*, 8 (s4), 335-41.
- 20. Porst, H. (2012). Premature ejaculation. In H. Porst, & Y. Reisman (Eds.), *The European Society of Sexual Medicine syllabus of sexual medicine* (pp. 547-92). Amsterdam (NL): Medix Publishers.
- 21. Porst, H., Montorsi, F., Rosen, R.C., Gaynor, L., Grupe, S., & Alexander, J. (2007). The premature ejaculation prevalence and attitudes (PEPA) survey: prevalence, comorbidities, and professional help-seeking. *Eur Urol*, 51, 816-24.
- 22. Rowland, D.L. (2011). Psychological impact of premature ejaculation and barriers to its recognition and treatment. *Cur Med Res Op* 27, 8, 1509-18.
- 23. Saint Lawrence, J.S., & Madakasira, S. (1992). Evaluation and treatment of premature ejaculation: a critical review. *Int J Psych Med*, 22, 1, 77-97
- 24. Symonds, T., Roblin, D., Hart, K., & Althof, S. (2003). How does premature ejaculation impact a man's life? *J Sex Mar Ther*, 29, 5, 361-70.