

Unusually high sea ice cover influences resource use by benthic invertebrates in coastal Antarctica

L. N. Michel^{1,*}, P. Dubois², M. Eleaume³, J. Fournier⁴, C. Gallut⁵, P. Jane⁶ & G. Lepoint¹

¹University of Liège, Belgium. ²Université Libre de Bruxelles, Belgium. ³National Museum of Natural History, France. ⁴Concarneau Biological Station, National Museum of Natural History, France. ⁵Pierre and Marie Curie University, France. ⁶Aquarium de Paris – Cinéaqua, France. *E-mail: loicnmichel@gmail.com, publications available at <http://goo.gl/xdxaHM>



Context, objectives & methods

Download this poster

- Climate change has contrasted effects on Antarctica. West Antarctic Peninsula: T° ↗ and sea ice ↘; East Antarctica: T° → ↘ and sea ice cover ↗.
- Dumont-d'Urville station (Adélie Land, East Antarctica) recently underwent an event of high spatial and temporal sea ice coverage (no seasonal breakup during austral summers 2013-2014 and 2014-2015)




▲ Accumulations of benthic biofilm on coastal rocky bottoms near Dumont-d'Urville station at the time of sampling (December 2014).


- How will Antarctic communities respond to such environmental changes? How could increased sea ice cover impact benthic food webs?
- Use of stable isotope ratios of C and N to identify resources supporting dominant benthic invertebrates (here: focus on primary consumers and omnivores)
- Use of SIAR mixing model to quantify relative importance of 4 producers / organic matter pools: sympagic algae, suspended particulate organic matter, benthic macroalgae (*Himantothallus grandifolius*) and benthic biofilm (unusually abundant, heterogeneous mix of microalgae, amorphous material and detrital items)

Results & discussion

1) Isotopic biplot of consumers and food sources



2) Main food items according to SIAR mixing model




OV: *O. validus*; SN: *S. neumayeri*; DB: *D. brucei*; HA: *Harmothoe* sp.; FM: *F. mundata*; PO: *Polycirrus* sp.; OP: *Ophiura* sp.; PE: *Perkinsiana* sp.; TL: *T. longstaffi*; MA: *Marseniopsis* sp.; HE: *Heterocucumis* sp.; LE: *L. elliptica*; AC: *A. colbecki*; ST: *Staurocucumis* sp.

3) Comparison with literature data

Species	DDU	1	2	3	4	5	6
<i>Laternula elliptica</i>	Green	Grey	Blue	Grey	Blue	Blue	Blue
<i>Adamussium colbecki</i>	Green	Grey	Blue	Grey	Blue	Blue	Blue
<i>Stereochinus neumayeri</i>	Orange	Purple	Purple	Purple	Purple	Purple	Purple
<i>Odontaster validus</i>	Orange	Purple	Purple	Purple	Purple	Purple	Purple
<i>Staurocucumis</i> sp.	Green	Grey	Blue	Grey	Blue	Blue	Blue
<i>Harmothoe</i> sp.	Orange	Purple	Purple	Purple	Purple	Purple	Purple

Sea ice



- Main food items
- Sympagic algae / Ice POM
 - Benthic algae / Biofilm
 - Plankton / SPOM
 - Sediment POM
 - Animal-based diet
 - No data

References: 1-3: Norkko et al. 2007 Ecology 88: 2810-2820; 4: Gillies et al. 2012 Estuar Coast Shelf S 97: 44-57; 5: Dunton 2001 Amer Zool 41: 99-112; 6: Corbisier et al. 2004 Polar Biol 27: 75-82

- Main organic matter source of most consumers: sympagic algae (1, 2).
- Important differences in resource use by consumers of Adélie Land and from other locations of coastal Antarctica, including sea ice influenced sites (3).
- Trophic importance of benthic biofilm comparatively limited despite high abundance (2).
- Norkko et al. 07: benthic invertebrates consume more detrital matter in sea ice influenced locations. Our results disagree with this hypothesis (2,3).
- However, no data about dynamics of biofilm accumulation + long-lived consumers with low metabolic rates: low isotopic turnover? Is isotopic equilibrium reached?

Conclusions & perspectives

- Important sea ice cover is linked with high reliance of benthic invertebrates on sympagic algae.
- Important spatial and/or temporal discrepancies in feeding habits and benthic community response to sea ice: high trophic plasticity?
- Interpretation of data is complicated by the lack of background data ("normal" conditions) and by physiological features of studied organisms.

Acknowledgements

Authors would like to thank the French Polar Institute for welcoming us at Dumont-d'Urville Station in the framework of the REVOLTA program. GL and PD are F.R.S-FNRS research associate and director, respectively. This work is part of the VERSO project, funded by the Belgian Science Policy Office (BELSPO).