

Will climate change affect insect pheromonal communication?

Antoine Boullis¹, Claire Detrain², Frédéric Francis¹ and François J Verheggen¹



Understanding how climate change will affect species interactions is a challenge for all branches of ecology. We have only limited understanding of how increasing temperature and atmospheric CO₂ and O₃ levels will affect pheromone-mediated communication among insects. Based on the existing literature, we suggest that the entire process of pheromonal communication, from production to behavioural response, is likely to be impacted by increases in temperature and modifications to atmospheric CO₂ and O₃ levels. We argue that insect species relying on long-range chemical signals will be most impacted, because these signals will likely suffer from longer exposure to oxidative gases during dispersal. We provide future directions for research programmes investigating the consequences of climate change on insect pheromonal communication.

Addresses

¹ Entomologie Fonctionnelle et Evolutive, Gembloux Agro-Bio Tech, Université de Liège, 2 Passage des Déportés, 5030 Gembloux, Belgium
² Service d'Ecologie Sociale, Université libre de Bruxelles, Campus de la Plaine, Boulevard du Triomphe, 1050 Brussels, Belgium

Corresponding author: Verheggen, François J. (fverheggen@ulg.ac.be)

Current Opinion in Insect Science 2016, 17:87–91

This review comes from a themed issue on **Global change biology**

Edited by **Vladimir Košťál** and **Brent J Sinclair**

<http://dx.doi.org/10.1016/j.cois.2016.08.006>

2214-5745/© 2016 Elsevier Inc. All rights reserved.

Introduction

Since the 19th century, the atmospheric concentration of greenhouse gases, particularly carbon dioxide (CO₂), have drastically increased causing changes to environmental parameters at a global scale, including temperature [1]. Recent studies now highlight the impact of such modifications on the whole dynamics of life [2]. Through cascade effects, entire ecosystems are being disturbed, impacting the population dynamics of inhabiting species and altering the ways that they interact with one another. This phenomenon has been well documented for insect–plant

interactions mediated by plant secondary metabolites [3,4*].

Communication between insects relies mainly on semi-chemicals, which are organic molecules involved in the chemical interactions between organisms [5]. They include pheromones (intraspecific communication) and allelochemicals (interspecific communication). Pheromones have a variety of important roles, especially related to foraging, aggregation or sexual behaviour [6]. Using behaviour-changing pheromones (named releaser pheromones) is central to integrated pest management (IPM) [7], so predicting the impact of climate change on IPM programmes depends on understanding the impact of changes in related abiotic parameters on insect pheromonal communication. However, few studies have focused on how changes in climate will disturb each stage in the pheromone pathway from emitters to receivers (Figure 1).


A pheromone's long journey

Biosynthesis

Most insect pheromones are synthesised *de novo* and secreted in specialised glandular tissues, regulated by various enzymatic activities [8,9]. Others are sequestered and/or derived from dietary precursors and depend on the nutritive quality of the diet. Elevated temperatures will likely have pronounced effects on pheromone biosynthesis. Because insects are ectothermic and poikilothermic, changing their body temperature will influence enzymatic activities [10], and impact pheromone biosynthesis both quantitatively and qualitatively. For example, temperature modifies the ratio of compounds in the sex pheromone of the potato tuber worm moth *Phthorimaea operculella* [11]. Moths (Lepidoptera: Heterocera) differentially use the same precursors to synthesise different pheromone components, thanks to a wide variety of enzymes (*i.e.* oxidase, desaturases, reductase), allowing specific recognition [12]. Although these insects can perceive a wide range of pheromone components, the activation of neurons in their macroglomerular complexes, and the elicitation of relevant behavioural responses, is combinatorial: it will happen only when the right combination and ratio of components is perceived at the same time [6].

Developmental temperature has a strong influence on adult life history, morphology, and physiology. Furthermore, in some species, pheromone production and availability is

Figure 1



Intraspecific chemical communication in insects may be subdivided into five steps that are probably impacted by modifications to atmospheric gas composition and associated raise in ambient temperature. Graphic art by Carolina Levicek.

dependent on larval, pupation, and/or adult developmental conditions [8,13,14], hence the effect of abiotic parameters on all the insect life stages is important. In the male beewolf, *Philanthus triangulum*, an increase of 5 °C in the larval rearing temperature led adult males to produce more pheromonal secretions [13]. Moreover, warmer rearing conditions led to higher relative amounts of compounds with high molecular weight. As a consequence, a shift in temperature could weaken intraspecific relationships of these insect species by reducing the efficiency (*i.e.* specificity, activity, timing of production, *etc.*) of their chemical communication.

Increasing atmospheric CO₂ concentrations [1] could also affect the biosynthesis of insect pheromones. Changes in CO₂ concentrations affect plant biochemistry, including the synthesis of secondary metabolites [4*]. Since some phytophagous insect species produce their pheromone components based on precursors taken from their host plant, we hypothesise that phytophagous insects could be among the most vulnerable to changes in atmospheric CO₂ concentrations, through cascade effects of CO₂ on plant chemistry [15,16*]. In *Holomelina* spp. moths, leucine is the starting material for sex pheromone production [17]. In bark beetles, while pheromones are produced primarily *de novo* mainly through the mevalonate pathway, some aggregation pheromone components arise from the hydroxylation of host tree-derived secondary metabolites [18].

Emission

Few studies have specifically investigated how changes in temperature and atmospheric gas composition act on pheromone release. In the moth *Striacosta ablicosta*, increase in average temperature does not affect the calling behaviour of females, while an increasing variation between photophase and scotophase temperatures alter significantly this behaviour, as also observed in *Phyllonorycter junoniella* [19]. Ladybird larvae deposit more long-chained hydrocarbons — used as oviposition deterring pheromone — when exposed to rising temperature [20**]. An increase in atmospheric CO₂ concentration reduces the emission rate of the alarm pheromone in pea aphids (*Acyrtosiphon pisum*) (Boullis *et al.*, unpublished).

Signal dispersal

After pheromone release by the emitter, volatile pheromones may be altered by oxidative gases such as ozone on their way to the receiver. Most pheromones are simple, lipophilic and of low molecular weight, which facilitate their long-distance dispersal in the air. Other pheromones are heavier molecules, including semi-volatile pheromones and cuticular hydrocarbons (CHC), which are used in short-range or contact communication [6]. Like phyto-genic volatile organic compounds (VOCs), insect pheromones made of unsaturated terpenes may be decomposed by ozone [21–23]. Similar terpenes are constitutive of sexual, aggregation or alarm pheromones in several insect taxa, such as ladybirds [24,25], aphids

[26,27], bark beetles [28] and fruit flies [29]. As highlighted for *Drosophila melanogaster*, terpenes could lose their biological activity after short-term ozone fumigation at environmentally-realistic concentrations (ranging from 40 to 120 ppb) [30]. The lifespan of trail-pheromones and alarm-pheromones, which act in a short time window and small spatial scale [31,32], may thus be further reduced in an ozone-rich atmosphere. In addition to the effect of ozone on pheromones, temperature acts on the volatility of semiochemicals. In the case of pheromones dispersed over long distances such as sex or aggregation pheromones, temperature changes may modify the shape of scent plumes and disturb the efficiency of insects to reach their target [33].

Increased temperature may also alter heavy molecules, such as cuticular pheromones involved in contact recognition. Because of their low volatility, temperature will likely more affect the chemical composition of pheromonal blends (ratios of components). Insects' cuticular lipids exist in a solid state at ambient temperatures, but they can partially melt upon contact with the animal's surface (with higher than ambient temperature), which, in turn, induces modifications in the ratios of cuticular composition [34]. As observed in *D. melanogaster*, a 4 °C increase of ambient temperature changes cuticular hydrocarbon composition, which lead to sexual isolation and affect the stability of ecological communities [35].

Perception

Pheromone perception occurs through a complex series of events, starting when pheromones enter the sensilla lymph and ending at brain processing [36–38]. Very few data are available on how environmental changes will impact pheromone perception, but one recent study showed that the sex pheromone perception was altered in male moth *Caloptilia fraxinella* under elevated temperature [39].

Because insects are poikilotherms, changes in their body temperature may alter the affinity between a pheromone and its binding protein (PBP) that transports this molecule through the sensillum lymph to olfactory receptor neurons. In *Apis mellifera* and *A. cerana*, ASP1 acts as PBP that has a good affinity to the queen mandibular pheromone [40]. However, increasing temperature weakens the van der Waals and hydrogen bonds established between the queen mandibular pheromone and PBP, which implies that binding affinity between these molecules can be lessened, inducing a reduced efficiency of the signal transportation through the hydrophilic lymph [41].

Behavioural response

Although insect responses to pheromones are innate, they may be conditional and influenced by direct (age, sex, hormonal status, experience) and indirect (cascade effect) factors [6]. Temperature is a major abiotic factor together

with photoperiod that determines the intensity and timing of various insects' activities [42,43]. Field studies related to IPM approaches on several lepidopterans have shown that the diel periodicity of their sexual attraction is modified by both photoperiod and ambient temperature [44–46]. Moreover, the seasonal rate of capture by trap catching is generally related to the associated temperature, depending on specific seasonal degree-days that insects are subjected to [47,48]. By this logic, an increase in global surface temperature may shift the seasonal periodicity of sex-related flights in insects, requiring an adaptation of monitoring and treatment periods against these pest insects. Another example is the impact of changing temperatures on ant foraging activity. Ants that use chemical recruitment tend to forage at lower temperatures compared to those that do not [49]. Therefore, accelerated pheromone decay caused by increased temperature is expected to alter trail-following behaviour and to be more detrimental to foraging by mass-recruiting ant species [50].

In addition to a general increase in insect mobility, some specific behavioural responses to pheromones can be altered by elevated temperature. For instance, male moths *C. fraxinella* reared under increased temperature during their reproductive diapause and subsequently exposed to female sex pheromones in a wind tunnel show more pronounced sexual responses [39]. At higher temperatures, male moths also show a lower level of specificity towards their sex pheromones, due to shifts in behavioural thresholds related to plume orientation and to the elicitation of upwind flight [51].

With regards to the impact of atmospheric CO₂ concentration, the escape behaviour of aphids reared under elevated CO₂ concentrations (*i.e.* 2100 predicted levels) is lower compared to those reared under ambient CO₂ conditions [52,53,54**]. The increase in CO₂ concentration could affect the escape behaviour of aphids by reducing the enzymatic activity of acetylcholinesterase, which is involved in neuronal transmission related to alarm signal perception [55]. This altered ability of aphids to produce and/or respond to the alarm pheromone may alter their defensive behaviours under changing climatic scenarios.

Conclusions, wider context and future directions

Based on the existing literature, we suggest that pheromonal communication in insects will be disturbed by increases in temperature and atmospheric gas concentrations. Insects relying on long-range chemical signalling, involving complex blends of molecules, are likely to be more impacted, due to the possible perturbation of enzymatic properties (leading to modification in compound ratio) or signal degradation by oxidative gases during dispersal (disrupting pheromone plumes). Behavioural

aspects of chemical communication (emission of the signal and induced behaviours) could also be affected as a consequence of shifts in optimal conditions affecting phenology and/or physiology. However, because climate change effects on pheromonal communication may be masked by the daily and seasonal rhythms of behaviour and physiology, our knowledge of the specific effects of climate change on pheromone signalling is sparse.

In this review we only focused on intraspecific (pheromonal) chemical communication, although allelochemical-mediated communication in the broader sense will likely also be affected by climate change. Indeed, as already suggested in plant–insect relationships, the changes of several abiotic parameters could affect interactions between organisms from different trophic levels, and thus affect the dynamics of ecological systems. However, it is difficult to predict how climate change will impact chemical communication between insects for several reasons. Abiotic factors could affect the different stages of insect pheromone communication (Figure 1), and the response of insects to particular environmental conditions could be species-specific. Moreover, the interactive effects of elevated atmospheric ozone and CO₂ concentrations, as well as temperature increase, on chemical-mediated interactions have received limited attention, despite that all of these factors are affecting ecosystems' stability. A key solution lies in the use of mesocosms and other facilities where multiple components of climate change can be manipulated in a multispecies context [56]. This approach could be used to assess how all climatic changes associated with a predicted scenario in the coming century might interact to impact the production of plant secondary metabolites, and the associated cascade effects on the pheromone production in phytophagous insects.

In an IPM context, the efficiency of pheromone slow-release devices could be reduced following climatic modifications, since their release kinetics are sensitive to various climatic parameters including temperature [57]. Moreover, because the behavioural response of insects to pheromones and allelochemicals could also be modified as a consequence of climate change, we suggest that all semiochemical-based IPM strategies will be impacted, including mass trapping, mating disruption, monitoring, push-pull strategies and other intercropping systems.

Acknowledgements

A. Boullis was financially supported by a PhD grant from the Fonds pour la Formation à la Recherche dans l'Industrie et l'Agriculture (FRIA), Belgium.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. IPCC: **Summary for policymakers**. In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Edited by Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM. Cambridge, United Kingdom/New York, NY, USA: Cambridge University Press; 2013.
2. Hughes L: **Biological consequences of global warming: is the signal already apparent?** *Trends Ecol Evol* 2000, **15**:56-61.
3. DeLucia EH, Nabity PD, Zavala JA, Berenbaum MR: **Climate change: resetting plant–insect interactions**. *Plant Physiol* 2012, **160**:1677-1685.
4. Ode PJ, Johnson SN, Moore BD: **Atmospheric change and induced plant secondary metabolites—are we reshaping the building blocks of multi-trophic interactions?** *Curr Opin Insect Sci* 2014, **5**:57-65.
- This is a recent review highlighting the impact of elevated greenhouse gas concentrations on plant defences against insect herbivores.
5. Law JH, Regnier FE: **Pheromones**. *Annu Rev Biochem* 1971, **40**:533-548.
6. Wyatt TD: *Pheromones and Animal Behavior*. Cambridge University Press; 2014.
7. Miller JR, Gut LJ: **Mating disruption for the 21st century: matching technology with mechanism**. *Environ Entomol* 2015, **44**:427-453.
8. Blomquist GJ, Jurenka R, Schal C, Tittiger C: **Pheromone production: biochemistry and molecular biology**. In *Insect Endocrinology*. Edited by Gilbert LI. Oxford: Elsevier; 2012: 523-567.
9. Tillman JA, Seybold SJ, Jurenka RA, Blomquist GJ: **Insect pheromones—an overview of biosynthesis and endocrine regulation**. *Insect Biochem Mol Biol* 1999, **29**:481-514.
10. Neven LG: **Physiological responses of insects to heat**. *Postharvest Biol Technol* 2000, **21**:103-111.
11. Ono T: **Effect of rearing temperature on pheromone component ratio in potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae)**. *J Chem Ecol* 1993, **19**:71-81.
12. Ando T, Yamakawa R: **Analyses of lepidopteran sex pheromones by mass spectrometry**. *Trend Anal Chem* 2011, **30**:990-1002.
13. Roeser-Mueller K, Strohm E, Kalltenpoth M: **Larval rearing temperature influences amount and composition of the marking pheromone of the male beewolf, *Philanthus triangulum***. *J Insect Sci* 2010, **10**:1-16.
14. Ono T: **Effect of temperature on biosynthesis of sex pheromone components in potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae)**. *J Chem Ecol* 1994, **20**:2733-2741.
15. Bidart-Bouzat MG, Imeh-Nathaniel A: **Global change effects on plant chemical defenses against insect herbivores**. *J Integr Plant Biol* 2008, **50**:1339-1354.
16. Boullis A, Francis F, Verheggen FJ: **Climate change and tritrophic interactions: will modifications to greenhouse gas emissions increase the vulnerability of herbivorous insects to natural enemies?** *Environ Entomol* 2015, **44**:277-286.
- A comprehensive review of the effects of climate and atmospheric changes on chemical communication-mediated interactions between phytophagous insect pests, their host plants, and their natural enemies.
17. Charlton RE, Roelofs WL: **Biosynthesis of a volatile, methyl-branched hydrocarbon sex pheromone from leucine by arctiid moths (*Holomelina* spp.)**. *Arch Insect Biochem Physiol* 1991, **18**:81-97.
18. Blomquist GJ, Figueroa-Teran R, Aw M, Song M, Gorzalski A, Abbott NL, Chang E, Tittiger C: **Pheromone production in bark beetles**. *Insect Biochem Mol Biol* 2010, **40**:669-671.
19. Mozūraitis R, Būda V: **Pheromone release behaviour in females of *Phyllonorycter junoniella* (Z.) (Lepidoptera, Gracillariidae) under constant and cycling temperatures**. *J Insect Behav* 2006, **19**:129-142.

20. Sentis A, Ramon-Portugal F, Brodeur J, Hemptinne JL: **The smell of change: warming affects species interactions mediated by chemical information.** *Glob Change Biol* 2015, **21**:3586-3594.
This study demonstrates the effect of elevated temperature on infochemical-related behaviours in ladybeetles. Temperature changes induce modification in terms of emission of marking pheromones as well as shifts in laying behaviour of receiving females.
21. Holopainen JK, Himanen SJ, Yuan JS, Chen F, Stewart CN: **Ecological functions of terpenoids and climate changes.** In *Handbook of Natural Products*. Edited by Ramawat KG, Merillon JM. Springer; 2013:2913-2940.
22. Pinto DM, Tiiva P, Miettinen P, Joutsensaari J, Kokkola H, Nerg A-M, Laaksonen A, Holopainen JK: **The effects of increasing atmospheric ozone on biogenic monoterpene profiles and the formation of secondary aerosols.** *Atmos Environ* 2007, **41**:4877-4887.
23. Pinto DM, Blande JD, Nykänen R, Dong W-X, Nerg A-M, Holopainen JK: **Ozone degrades common herbivore-induced plant volatiles: does this affect herbivore prey location by predators and parasitoids?** *J Chem Ecol* 2007, **33**:683-694.
24. Fassotte B, Fischer C, Durieux D, Lognay G, Haubruge E, Francis F, Verheggen FJ: **First evidence of a volatile sex pheromone in lady beetles.** *PLOS ONE* 2014, **9**:e115011.
25. Fassotte B, Francis F, Verheggen FJ: **The scent of love: how important are semiochemicals in the sexual behavior of lady beetles?** *J Pest Sci* 2016, **89**:347-358.
26. Francis F, Vandermoten S, Verheggen F, Lognay G, Haubruge E: **Is the (E)- β -farnesene only volatile terpenoid in aphids?** *J Appl Entomol* 2005, **129**:6-11.
27. Boullis A, Verheggen F: **Chemical ecology of aphids.** In *Biology and Ecology of Aphids*. Edited by Vilcinskas A. CRC Press; 2016.
28. Taft S, Najjar A, Erbilgin N: **Pheromone production by an invasive bark beetle varies with monoterpene composition of its naïve host.** *J Chem Ecol* 2015, **41**:540-549.
29. Sarles L, Verhaeghe A, Francis F, Verheggen F: **Semiochemicals of *Rhagoletis* fruit flies: potential for integrated pest management.** *Crop Prot* 2015, **78**:114-118.
30. Arndt U: **Air pollutants and pheromones—a problem?** *Chemosphere* 1995, **30**:1023-1031.
31. Czaczkes TJ, Grüter C, Ratnieks FL: **Trail pheromones: an integrative view of their role in social insect colony organization.** *Annu Rev Entomol* 2015, **60**:581-599.
32. Joachim C, Vosteen I, Weisser WW: **The aphid alarm pheromone (E)- β -farnesene does not act as a cue for predators searching on a plant.** *Chemoecology* 2015, **25**:105-113.
33. McFrederick QS, Fuentes JD, Roulston TA, Kathilankal JC, Lerda M: **Effects of air pollution on biogenic volatiles and ecological interactions.** *Oecologia* 2009, **160**:411-420.
34. Gibbs AG: **Lipid melting and cuticular permeability: new insights into an old problem.** *J Insect Physiol* 2002, **48**:391-400.
35. Bontonou G, Denis B, Wicker-Thomas C: **Interaction between temperature and male pheromone in sexual isolation in *Drosophila melanogaster*.** *J Evol Biol* 2013, **26**:2008-2020.
36. Zhang J, Walker WB, Wang G: **Pheromone reception in moths: from molecules to behaviors.** In *Progress in Molecular Biology and Translational Science*. Edited by Glatz RB. Academic Press; 2015:109-128.
37. Zhou J-J: **Odorant-binding proteins in insects.** In *Pheromones*. Edited by Litwack G. Elsevier Academic Press; 2010:241-272.
38. Ishida Y, Leal WS: **Chiral discrimination of the Japanese beetle sex pheromone and a behavioral antagonist by a pheromone-degrading enzyme.** *Proc Natl Acad Sci U S A* 2008, **105**:9076-9080.
39. Lemmen J, Evenden M: **Environmental conditions terminate reproductive diapause and influence pheromone perception in the long-lived moth *Caloptilia fraxinella*.** *Physiol Entomol* 2015, **40**:30-42.
40. Slessor KN, Winston ML, Le Conte Y: **Pheromone communication in the honeybee (*Apis mellifera* L.).** *J Chem Ecol* 2005, **31**:2731-2745.
41. Weng C, Fu Y, Jiang H, Zhuang S, Li H: **Binding interaction between a queen pheromone component HOB and pheromone binding protein ASP1 of *Apis cerana*.** *Int J Biol Macromol* 2015, **72**:430-436.
42. Bonsignore CP, Bellamy C: **Daily activity and flight behaviour of adults of *Capnodis tenebrionis* (Coleoptera: Buprestidae).** *Eur J Entomol* 2007, **104**:425-431.
43. Seybold SJ, King JA, Harris DR, Nelson LJ, Hamud SM, Chen Y: **Diurnal flight response of the walnut twig beetle, *Pityophthorus juglandis* Blackman (Coleoptera: Scolytidae), to pheromone-baited traps in two northern California walnut habitats.** *Pan-Pac Entomol* 2012, **88**:231-247.
44. Comeau A, Cardé RT, Roelofs WL: **Relationship of ambient temperatures to diel periodicities of sex attraction in six species of Lepidoptera.** *Can Entomol* 1976, **108**:415-418.
45. Lance DR, Odell TM, Mastro VC, Schwalbe CP: **Temperature-mediated programming of activity rhythms in male gypsy moths (Lepidoptera: Lymantriidae): implications for the sterile male technique.** *Env Entomol* 1988, **17**:649-653.
46. Cardé RT, Charlton RE, Wallner WE, Baranchikov YN: **Pheromone-mediated diel activity rhythms of male Asian gypsy moths (Lepidoptera: Lymantriidae) in relation to female eclosion and temperature.** *Ann Entomol Soc Am* 1996, **89**:745-753.
47. Del Tío R, Martínez JL, Ocete R, Ocete ME: **Study of the relationship between sex pheromone trap catches of *Lobesia botrana* (Den. & Schiff.) (Lep., Tortricidae) and the accumulation of degree-days in Sherry vineyards (SW of Spain).** *J Appl Entomol* 2001, **125**:9-14.
48. Gallardo A, Ocete R, López MA, Maistrello L, Ortega F, Semedo A, Soria FJ: **Forecasting the flight activity of *Lobesia botrana* (Denis & Schiffermüller) (Lepidoptera, Tortricidae) in southwestern Spain.** *J Appl Entomol* 2009, **133**:626-632.
49. Ruano F, Tinaut A, Soler JJ: **High surface temperatures select for individual foraging in ants.** *Behav Ecol* 2000, **11**:396-404.
50. van Oudenhove L, Billoir E, Boulay R, Bernstein C, Cerdá X: **Temperature limits trail following behavior through pheromone decay in ants.** *Naturwissenschaften* 2011, **98**:1009-1017.
51. Linn CE, Campbell MG, Roelofs W: **Temperature modulation of behavioural thresholds controlling male moth sex pheromone response specificity.** *Physiol Entomol* 1986, **13**:59-67.
52. Awmack C, Woodcock C, Harrington R: **Climate change may increase vulnerability of aphids to natural enemies.** *Ecol Entomol* 1997, **22**:366-368.
53. Mondor EB, Tremblay MN, Awmack CS, Lindroth RL: **Divergent pheromone-mediated insect behaviour under global atmospheric change.** *Glob Change Biol* 2004, **10**:1820-1824.
54. Hentley WT, Vanbergen AJ, Hails RS, Jones TH, Johnson SN: **Elevated atmospheric CO₂ impairs aphid escape responses to predators and conspecific alarm signals.** *J Chem Ecol* 2014, **40**:1110-1114.
- Rare are the studies that assess the impact of elevated CO₂ on pheromonal signalling in insects. Here, the authors showed that an increase in CO₂ concentration reduce the escape behaviour in aphids when alarm pheromone is presented to the colony.
55. Sun Y, Su J, Ge F: **Elevated CO₂ reduces the response of *Sitobion avenae* (Homoptera: Aphididae) to alarm pheromone.** *Agric Ecosyst Environ* 2010, **135**:140-147.
56. Stewart RI, Dossena M, Bohan DA, Jeppesen E, Kordas RL, Ledger ME, Meerhoff M, Moss B, Mulder C, Shurin JB et al.: **Mesocosm experiments as a tool for ecological climate-change research.** *Adv Ecol Res* 2013, **48**:71-181.
57. Heuskin S, Verheggen FJ, Haubruge E, Wathelet J-P, Lognay G: **The use of semiochemical slow-release devices in integrated pest management strategies.** *Biotechnol Agron Soc* 2011, **15**:459-470.