

DEBATE

Open Access



Sarcopenia in daily practice: assessment and management

Charlotte Beaudart^{1*} , Eugène McCloskey^{2,3}, Olivier Bruyère¹, Matteo Cesari^{4,5}, Yves Rolland⁶, René Rizzoli⁷, Islène Araujo de Carvalho⁸, Jotheeswaran Amuthavalli Thiyagarajan⁸, Ivan Bautmans⁹, Marie-Claude Bertière¹⁰, Maria Luisa Brandi¹¹, Nasser M. Al-Daghri¹², Nansa Burlet¹, Etienne Cavalier¹³, Francesca Cerreta¹⁴, Antonio Cherubini¹⁵, Roger Fielding¹⁶, Evelien Gielen¹⁷, Francesco Landi¹⁸, Jean Petermans¹⁹, Jean-Yves Reginster¹, Marjolein Visser^{20,21}, John Kanis^{2,22} and Cyrus Cooper^{23,24}

Abstract

Background: Sarcopenia is increasingly recognized as a correlate of ageing and is associated with increased likelihood of adverse outcomes including falls, fractures, frailty and mortality. Several tools have been recommended to assess muscle mass, muscle strength and physical performance in clinical trials. Whilst these tools have proven to be accurate and reliable in investigational settings, many are not easily applied to daily practice.

Methods: This paper is based on literature reviews performed by members of the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO) working group on frailty and sarcopenia. Face-to-face meetings were afterwards organized for the whole group to make amendments and discuss further recommendations.

Results: This paper proposes some user-friendly and inexpensive methods that can be used to assess sarcopenia in real-life settings. Healthcare providers, particularly in primary care, should consider an assessment of sarcopenia in individuals at increased risk; suggested tools for assessing risk include the Red Flag Method, the SARC-F questionnaire, the SMI method or different prediction equations. Management of sarcopenia should primarily be patient centered and involve the combination of both resistance and endurance based activity programmes with or without dietary interventions. Development of a number of pharmacological interventions is also in progress.

Conclusions: Assessment of sarcopenia in individuals with risk factors, symptoms and/or conditions exposing them to the risk of disability will become particularly important in the near future.

Keywords: Sarcopenia, Daily practice, Assessment, Management, Tools

Background

The term sarcopenia was first coined by Rosenberg et al. in 1989 [1] as a progressive loss of skeletal muscle mass with advancing age. Since then, the definition has expanded to incorporate the notion of impaired muscle strength and/or physical performance. Currently, several definitions of sarcopenia have been proposed [2–10] but no consensus has yet been reached. Depending on the definition used, the prevalence of sarcopenia is reported to be up to 29 % for older community-dwelling adults and up to 33 % for individuals living in long-term care

institutions [11, 12]. Sarcopenia is associated with morbidity and mortality from linked physical disability, falls, fractures, poor quality of life, depression and hospitalization [13–19].

Current research is focusing on nutritional exercise/activity based and other novel interventions for improving the quality and quantity of skeletal muscle in older people. Some studies demonstrated that resistance training combined with nutritional supplements can improve muscle function [11, 20–22]. A number of pharmacological interventions are in development but no single agent has been shown to be clinically effective, without unwanted effects, in maintaining or increasing skeletal muscle mass or function. With the prospect of effective interventions, the identification and assessment of sarcopenia will become

* Correspondence: c.beaudart@ulg.ac.be

¹Department of Public Health, Epidemiology and Health Economics, University of Liège, Avenue Hippocrate 13, CHU B23, 4000 Liège, Belgium
Full list of author information is available at the end of the article

particularly important to prevent disability and other negative health outcome in the near future.

The challenge in clinical practice will be in the assessment of sarcopenia to identify those who might benefit most from these interventions. Among the 1 current definitions of sarcopenia [3, 7, 8], there is a general agreement on the need for muscle mass measurement with varying recommendations on the roles of muscle strength assessment and/or physical performance. Currently, several well validated tools exist to measure these parameters, which have been reviewed recently [18, 23, 24]. Whereas they have been used for sarcopenia case finding in the research setting, their use is not always feasible in daily clinical practice. The purpose of this paper is to discuss different approaches in the assessment of sarcopenia and potential management strategies in clinical practice.

Methods

As in previous initiatives and publications [25–35], the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO) working group on frailty and sarcopenia consists of clinical scientists and experts in the field of musculoskeletal diseases. Different members of the ESCEO working group were asked to prepare a review of the literature on 1) the general tools for the assessment of sarcopenia, both in research and in clinic (CC); 2) the assessment of physical performance in daily practice (MC); 3) the role of imaging in the diagnosis of sarcopenia in daily practice (MV); 4) the role of biochemical markers in the diagnosis of sarcopenia in daily practice (EC) and 5) the role of primary versus secondary care physicians in the evaluation of sarcopenia (AC). A brief summary of the management of sarcopenia in daily practice was also proposed and discussed. Randomized controlled studies, prospective studies, systematic reviews and meta-analyses published before September 2015 were searched on PubMed and Embase using the following search terms : 1) Sarcopenia, Clinical, Evaluation, Assessment, Management; 2) Physical function, Physical performance, Gait, Walk, Walking, Strength; 3) Elderly, Muscle mass, Sarcopenia, Dual x-ray absorptiometry/DXA/DEXA, Computer tomography/CT, Magnetic resonance imaging/MRI, Bioelectrical impedance/BIA; 4) Frailty, Sarcopenia, Biomarker, Biochemical marker, and 5) Primary care, Specialist care, Secondary care, Sarcopenia, Management, Screening, Questionnaire. Additional studies were identified by a manual search of bibliographic references of relevant articles and existing reviews. Each member prepared a list of the most important papers based on their review of the literature and then made a set of preliminary recommendations. The subsequent step was a face-to-face meeting for the whole group to make amendments and discuss further recommendations. The plan of the manuscript was also discussed and shared

conclusions were reached. The views expressed in this article are the personal views of the authors and may not be understood or quoted as being made on behalf of or reflecting the position of the EMA or one of its committees or working parties.

Results

How to assess sarcopenia in clinical practice?

Despite a relatively large number of tools being available to measure muscle mass, muscle strength and physical performance [36, 37], some of them are likely to be of greater validity and utility for the assessment of sarcopenia in clinical practice than in clinical research and are summarised in Table 1. Whereas some biochemical markers of muscle metabolism (e.g. activin, n-terminal propeptide of procollagen III and myostatin) are being investigated for their ability to indicate muscle mass or strength, current data suggest that it is premature to recommend their use in daily practice [38].

Table 1 Applicability of the existing tools for the assessment of muscle mass, muscle strength and physical performance in research and clinical settings

	Applicable in research settings	Applicable in specialist clinical settings	Applicable in primary care settings
Assessment of muscle mass			
<i>DXA</i>	+++	+++	+
<i>Anthropometric measurements</i>	+	++	++
<i>CT-scan</i>	+++	++	+
<i>MRI</i>	+++	++	+
<i>BIA</i>	++	++	+
Assessment of muscle strength			
<i>Handgrip strength</i>	+++	+++	+++
<i>Lower limb muscle strength</i>	+++	++	+
<i>Repeated chair stands test</i>	+	+	++
Assessment of physical performance			
<i>Gait speed</i>	+++	+++	+++
<i>Timed Up and Go test</i>	++	+	+
<i>Balance test</i>	+	+	+
<i>6-min walk test</i>	++	+	+
<i>400 m walk test</i>	++	+	+
<i>Stair climb test</i>	++	+	+
<i>SPPB test</i>	+++	++	+

SPPB Short Physical Performance Battery

Nb. The group has chosen to attribute to each tool +++ (best recommended tool) or ++ (best alternative tool) or + (less recommended tool) based on the availability and the costs of the tool, the required time for the examination and the availability of robust cut-off points

There are currently a number of approaches to the definition of sarcopenia in clinical practice [3, 7, 36]. However, these are usually more suited to research studies than wider clinical practice. Additionally, some of the available methodologies for the assessment of sarcopenia utilise methods for measuring muscle mass, strength and physical function that are more suited to secondary care, than primary care settings. We therefore tabulated our preferences according to feasibility, complexity, required time for the examination, availability of robust cut-off points and cost, in each of these three contexts: research, specialist settings and primary care (Table 1).

Assessment of muscle mass

The widespread use of magnetic resonance imaging (MRI) and computed tomography (CT) scan for the non-invasive assessment of muscle mass [39] is limited in primary care settings by difficulties in access, costs, the lack of portable equipment and the requirement of highly specialized personnel.

Dual-energy x-ray absorptiometry (DXA) is a well-established, low-radiation technique used to assess body composition and provides reproducible estimates of appendicular skeletal lean mass [40, 41]. It is acknowledged that the accuracy of DXA for assessing muscle mass in people of different ages and different pathological conditions may vary. Moreover, DXA (in contrast to CT-scan and MRI) cannot assess intra-muscular fat, which turns out to be of increasing importance in terms of the quality of muscle and associations with clinical outcomes. Bearing these limitations in mind, DXA is still considered as the procedure of choice for routine clinical assessment. Using DXA, appendicular skeletal lean mass (ALM) is measured as the sum of the non-bone and non-fat mass of the four limbs. To adjust for body size, a skeletal muscle index (SMI) is derived as $ALM/height^2$. Thresholds of SMI at two standard deviations below the mean SMI of young male and female reference groups have been proposed as gender-specific cut-off points for sarcopenia. This results in two thresholds, proposed by the EWGSOP [3], the first of 5.5 kg/m^2 for women and 7.26 kg/m^2 [8] for men and the second of 5.67 kg/m^2 for women and 7.25 kg/m^2 for men [42], depending on the reference group on which these cut-off have been established. Using a different approach, the FNIH sarcopenia project [7] has also recently defined cut-offs for appendicular lean mass adjusted for body mass index (BMI), giving values of <0.512 for women and <0.789 for men. However, it should be pointed that these cut-offs might also be modified according to ethnicity [43].

If clinicians have no access to DXA, they can use anthropometric measurements. Indeed, a recent survey [44] showed that anthropometric data are currently the most widely used methods in clinical practice (57.5 % of

clinicians that measure muscle mass in their practice use anthropometric data) followed by DXA (45.9 %). Several anthropometric measurements exist (i.e. body mass index, calf circumference, mid-upper arm circumference and skinfold thickness). Moreover, mid-arm muscle and calf circumferences have been shown to be correlated with appendicular muscle mass and reflect both health and nutritional status and predict performance, health and survival in older people [45–47]. However, with advancing age, changes in the distribution of fat and loss of skin elasticity are such that circumference and skinfold measures incur a loss of accuracy and precision in older people [47, 48]. Some studies suggest that an adjustment of anthropometric measurements for age, sex or BMI results in a better correlation with DXA-measured lean mass [49–51]. Anthropometric measurements are simple clinical prediction tools that can be easily applied for sarcopenia since they offer the most portable, commonly applicable, inexpensive and non-invasive technique for assessing size, proportions and composition of the human body [50]. However, their validity is limited when applied to individuals due to large prediction errors and because cut-off points, to identify low muscle mass, still need to be defined. Therefore, if a patient is identified as at risk of having sarcopenia by anthropometric measurements, an additional measurement of muscle mass with DXA would still be recommended.

Finally, bio-electrical impedance analysis (BIA) is a method which estimates the volume of fat and lean body mass based on the relationship between the volume of a conductor and its electrical resistance. The method is not expensive, requires no specialized staff and is relatively easy to use in clinical practice, both on ambulatory subjects or on hospitalized patients. Moreover, reference values have been established for older individuals [3]. Even if the method's accuracy has been challenged and has been reported to overestimate muscle mass and underestimate fat mass [52–54], it is possible to use some adjustment equations to obtain valid measurements [55].

In summary, we would propose assessing primarily muscle mass by DXA, if this tool is available, and if not, anthropometry measurements can easily be used, in primary care settings, as a first screening of patients with low muscle mass. These patients can then be referred for an additional evaluation in specialist clinical settings.

Assessment of muscle strength

Handgrip strength appears to be the most widely used method for the measurement of muscle strength. A recent survey indicated that clinicians, both from the fields of geriatric medicine and rheumatology, prefer the use of grip strength over chest press and lower limb isokinetic

dynamometry as a measure of overall muscle strength [44]. In general, isometric handgrip strength shows a good correlation with leg strength [56] and also with lower extremity power, knee extension torque and calf cross-sectional muscle area [15, 57]. The measurement is easy to perform, inexpensive and does not require a specialist trained staff. Standardized conditions for the test [58] include seating the subject in a standard chair with their forearms resting flat on the armchairs. Clinicians should demonstrate the use of the dynamometer and show that gripping very tightly registers the best score. Six measures should be taken, 3 with each arm. Ideally, the patients should be encouraged to squeeze as hard and as tightly as possible during 3–5 seconds for each of the 6 trials; usually the highest reading of the 6 measurements is reported as the final result. The Jamar dynamometer, or similar hydraulic dynamometer, is the gold standard for this measurement. However, for patients with advanced arthritis, the design of this dynamometer may be a limitation [59]. A pneumatic dynamometer, such as the Martin vigorimeter, may be a good alternative. With this device, patients try to squeeze rubber balls (available in three sizes) with the same protocol as that used for the Jamar dynamometer. A variety of thresholds of grip strength have been proposed to characterize low muscle strength, ranging from 16 to 20 kg for women and 26–30 kg for men [7, 15, 60, 61]. Lower limb muscle strength, most frequently of the quadriceps, can also be measured. Commercial dynamometers can enable isometric and/or isokinetic measurements of strength. Even if these measurements are feasible in frail people [62, 63], they are often limited in clinical practice by their relative expense, the need to purchase dedicated equipment, the lack of trained staff and limited data in older populations. However, the repeated chair stand test, which is a timed test requiring participants to rise from a chair without using their arms and return to the seated position, consecutively, for five times, has been shown to be able to provide a reasonably reliable and valid indication of lower body strength [64].

In summary, we would recommend to measure muscle strength by handgrip strength in clinical practice (Table 1). For primary care settings where the availability of a handgrip dynamometer is not systematic, the repeated chair stand test could be used as an alternative measure of muscle strength.

Assessment of physical performance

The most widely used tool in clinical practice for the assessment of physical performance is the gait speed measurement, employed by almost two-thirds (63.3 %) of clinicians that assess physical performance (among 255 clinicians who took part in an international online survey; 87.8 % of medical doctors with geriatrics (57.6 %

and rheumatology (18.8 %) as major fields of interest) [44]. The test is highly acceptable for participants and health professionals in clinical settings [65]. No special equipment is required as it only needs a flat floor devoid of obstacles. In the 4-m gait speed test, which is recommended by the EWGSOP for the assessment of sarcopenia, men and women with a gait speed <0.8 m/s are described as having a poor physical performance [15]. The average extra-time added to the consultation by measuring the 4-metre gait speed was only 95 ± 20 s.

Gait speed can be performed alone or as part of a test battery, the most popular of which is the Short Physical Performance Battery (SPPB). The SPPB is a test scored to a maximum of 12 points comprising an assessment of gait speed (over 3–4 m), a balance test and a repeated chair stand test. These tests focus on lower extremity function, as the latter has been shown to correlate with mobility, disability and patient outcomes including hospitalization, institutionalization, and mortality. The SPPB takes about 10 min to complete [66]. Participants presenting a score ≤ 8 points have been described as having a poor physical performance [3].

Other standalone tests can be performed to assess physical performance. In the Timed Up and Go (TUG) test, individuals are asked to rise from a standard armchair, walk to a marker 3 m away, turn, walk back and sit down again. The 6-min walk distance or 400 m walk time can be used to measure aerobic capacity. The stair climb power test also shows good correlation with other measures of leg power and physical performance, but is mostly restricted to use in research settings [67].

In summary, we would propose that physical performance is primarily assessed in clinical practice by measuring gait speed. The SPPB test may be limited by the time of administration but might also be useful to identify men and women with low physical performance (Table 1).

The role of primary care physicians

In view of the current lack of a consensus concerning the definition of sarcopenia and also of the practical issues related to time constraints and limited access to assessment tools in the primary care setting, the group believes that the role of primary care physicians should be to identify patients who are at risk of sarcopenia and to refer them to specialists in the field. Some interesting methods that might be suitable for screening purpose are presented in the following section.

Consideration of possible sarcopenia should be undertaken in older individuals (e.g. > 65 years) with signs or symptoms suggestive of the condition both in primary care and in specialized clinical settings. Several methods can be proposed to perform a simple, rapid and inexpensive

identification of those at risk. However, none of them has received an extensive validation, and therefore further research in this area is urgently needed.

The red flag method

The purpose of the red flag method is to understand, during a standard medical consultation (or health assessment) the clinical presentation of individuals with particular regard to physical manifestations of sarcopenia such as general weakness or loss of muscle mass. The subject can also be asked about symptoms such as loss of weight, loss of muscle strength, loss of energy, falls, etc. (Table 2). An assessment of nutrition habits should also be performed to check, for example, if the subject has sufficient protein intake. The Mini-Nutritional Assessment could also be used for a rapid and easy assessment of malnutrition or, at least, risk of malnutrition [68]. Finally, clinicians can also assess physical activity. Indeed physical inactivity or high levels of sedentary behaviour may be considered a red flag. If the screening identifies any red flag suggesting the presence of sarcopenia, more sophisticated assessment procedures of sarcopenia can be implemented. Red flags have been identified through reviewed papers identified by members of the group and are presented in Table 2.

The SARC-F questionnaire

The SARC-F questionnaire [69] was developed as a possible rapid screening test for sarcopenia. This questionnaire could enable healthcare providers to quickly and easily assess the risk of sarcopenia during a standard health consultation. The subject is asked 5 questions addressing strength, assistance in walking, rising from a

chair, stair climbing and falls. Each component is scored from 0 to 2 points, giving a global score of the SARC-F between 0 and 10 points. A score ≥ 4 points is reported to be predictive of sarcopenia and poor outcomes and could be a trigger for a more detailed assessment of sarcopenia.

Despite a questionable sensitivity [70], the SARC-F questionnaire is considered as one of the best available tools to be used in primary care for raising awareness of the diagnosis of sarcopenia. Similarly to the red flag method, a result ≥ 4 for the SARC-F questionnaire could be an incentive to send the subject to a complete assessment of sarcopenia.

Prediction of low muscle mass according to age and BMI

Recently, a study [71] has been performed with the purpose to identify predictors of low skeletal muscle mass in older adults toward development of a practical clinical assessment tool for use by clinicians to identify individuals requiring DXA screening for muscle mass. For this purpose, ALM was calculated from DXA scans and SMI defined as the ratio of ALM divided by height in square centimetres. Older participants (from 65 to 85 years) were classified as having low muscle mass if their SMI was 1 standard deviation below the mean SMI of young adults. This model was validated on a sample of 200 subjects of the NHANES population. Results of the validation analysis revealed that age and BMI were strongly associated with a low SMI and may be an informative predictor in the primary care settings. Consequently, two models were proposed, one for men and one for women and consist of two tables presenting the probability of low muscle mass by age and BMI. In a 200-person validation, the model sensitivity was 81.6 % for men and 90.6 % for women and the model specificity was 66.1 % for men and 66.2 % for women.

Anthropometric prediction equation in combination with a measure of muscle function

Other authors developed gender specific anthropometric equations, based on age, weight, BMI values, to estimate appendicular skeletal muscle mass [72]. To validate these prediction equations, muscle mass was assessed using DXA in three cohorts of older Australian subjects [72] (appendicular skeletal muscle mass prediction equation: $10.05 + 0.35(\text{weight}) - 0.62(\text{BMI}) - 0.02(\text{age}) + 5.10$ (if male)). The results showed a strong correlation between the equations and the muscle mass measured using DXA, with an adjusted R^2 of 0.869. In a subsequent research the prediction equations were evaluated in combination with assessment of hand grip strength as a screening method to identify older patients who should undergo DXA evaluation for sarcopenia. The best strategy to reduce the number of DXA was to apply the equation

Table 2 The Red Flag method

	Red flags
Clinician's observation	General weakness of the subject
	Visual identification of loss of muscle mass
	Low walking speed
Subject's presenting features	Loss of weight
	Loss of muscle strength, in arms or in legs
	General weakness
	Fatigue
	Falls
	Mobility impairment
	Loss of energy
Clinician's assessment	Difficulties in physical activities or activities of daily living
	Nutrition
	Body weight
	Physical activity

Nb. Red flags have been identified through reviewed papers identified by members of the group

first, to assess hand grip strength in those with low estimated muscle mass and to proceed to DXA only in individuals with low grip strength [50].

Prediction of sarcopenia using age, handgrip strength and calf circumference

In 2014, Ishii et al. [73] developed a new screening tool for sarcopenia in a sample of almost 2000 autonomous community-dwelling older subjects in Japan. Sarcopenia was defined on the basis of low muscle mass measured by BIA and either low muscle strength characterized by handgrip or low physical performance characterized by slow gait speed. Using a database including demographic variables, albumin, chronic diseases, physical activity information and anthropometric measurements, the authors developed a gender specific model including three variables, i.e. age, handgrip strength and calf circumference. Based on the model, the authors constructed a gender specific score chart that had an excellent discrimination ability, with an area under the curve of 0.939 for men and 0.909 for women. The formula to calculate the scores are as follows: score in men, $0.62 \times (\text{age}-64) - 3.09 \times (\text{grip strength}-50) - 4.64 \times (\text{calf circumference}-42)$; score in women, $0.80 \times (\text{age}-64) - 5.09 (\text{grip strength}-34) - 3.28 \times (\text{calf circumference}-42)$. The corresponding probabilities of sarcopenia were calculated as: probability in men, $1 / [1 + e^{-(\text{sum score}/10-11.9)}]$; probability in women, $1/[1 + e^{-(\text{sum score}/10-12.5)}]$. This model still requires further validation in independent cohorts, before its use in clinical practice can be promoted.

How to manage sarcopenia in daily practice?

Identification of comorbidities

Sarcopenia is frequently found in association with comorbidities, e.g. osteoporosis, osteopenia, obesity, type II diabetes mellitus, breast cancer, etc. [74, 75]. In such cases, sarcopenia may be considered as a secondary consequence of the co-existing pathological condition. The impact of management of these conditions (e.g. better diabetic control, reduction of inflammatory status, or weight loss in obesity due to an energy-restricted diet) on the accompanying sarcopenia is unclear [76].

Physical activity

Physical activity interventions and progressive resistance training have been suggested to have a predominant effect on muscle strength, muscle mass and physical performance in older people [77].

However, so far, studies mainly focusing on well-defined sarcopenia with standardization of the physical intervention are still missing. Hence, it is still difficult to give a patient-specific physical activity prescription for the management of sarcopenia. However, healthcare providers can nevertheless give some general recommendations in order to improve other common conditions in older

adults (WHO recommendation: http://www.who.int/diet-physicalactivity/factsheet_olderadults/en/). Moreover, in their review, Cruz Jentoft et al [11]. forwarded two recommendations regarding the management of physical activity interventions in older people. First, to obtain an impact on muscle function, the duration of the intervention should be for at least 3 months. Second, supervised resistance exercise or multicomponent/combined exercise programs should be recommended for frail or sedentary community-dwelling people.

Nutrition

Although nutrition is considered as a major point for the management of sarcopenia, evidence of the effect of nutrition on muscle function is often derived from short-term studies in specifically selected sample and large clinical trials are still lacking. Currently, there is no robust evidence for nutritional recommendations for subjects with sarcopenia.

However, even if randomized controlled trials are inconsistent regarding the effects of protein supplementation on muscle function, several observational studies have suggested that maintaining adequate protein intake may help preserve muscle mass and strength in both adults and older people [78, 79]. Bauer et al. [80] recommended increasing protein intake to 1.2 g/kg body weight/day either by diet or by protein supplementation in older adults because of blunted muscle protein synthetic response and blunted post-prandial inhibition of muscle protein breakdown (anabolic resistance). Frail older adults or older who have acute or chronic diseases need higher dietary protein (i.e. 1.2–1.5 g/kg body weight/d) [80]. Recent evidence suggests that the recommended dietary allowance for protein is inadequate in older people [81]. Some other nutritional supplements, such as β -hydroxy β -methylbutyrate, creatine and vitamin D have been suggested to have an effect on muscle function. Indeed, β -hydroxy β -methylbutyrate supplements appear to increase muscle mass whilst its effects on muscle strength and physical performance are inconsistent [11, 20, 21]. Supplementation with creatine, protein or leucine combined with resistance exercises seems to have a positive impact on muscle mass, muscle strength and physical performance [22, 82, 83]. Finally, a recent meta-analysis has suggested that vitamin D supplementation could increase lower limb muscle strength [84]. Based on this evidence, dietary protein caloric intake, protein quality, as well as the vitamin D status of older individuals could be checked by clinicians and/or dieticians and individual prescription of nutritional supplements could be considered.

Pharmacological management

Currently, no drug is registered for the treatment of sarcopenia. However, several new chemical entities are

Table 3 Pharmacological agents in development with potential for treating sarcopenia

Mechanism of action	Drug name	Drug Developer	Indication sought	Study phase
I. Myostatin Antagonists				
Activin receptor trap	ACE-031	Acceleron	Duchenne muscular dystrophy	Phase 3 (trial terminated early)
Myostatin antibody	REGN-1033	Regeneron/Sanofi	Sarcopenia	Phase 2
	LY-2495655	Eli Lilly	Hip arthroplasty	Phase 2
			Elderly Fallers	
			Cancer Cachexia	
Activin receptor inhibitor	PF-06252616	Pfizer	Inclusion body myositis	Phase 1
	Bimagrumab (BMY338)	Novartis	Sarcopenia	Phase 2 and 3
			Hip fracture	Phase 2
			Cancer and COPD cachexia	
II. Selective Androgen Receptor Modulators	Enobasarm (Ostarine)	GTx	Cancer Cachexia	Phase 3 (did not meet primary endpoint)
III. Skeletal Troponin Activators	Tirasemtiv	Cytokinetics	ALS	Phase 2,3
	CK-2017357		Myasthenia Gravis	

currently at various stages of development. These are summarized in Table 3 with their potential future indications and their current phase of development.

Discussion and general consensus

The ESCEO Experts group agreed on some general recommendations to be implemented in clinical practice:

- Several tools are currently available for the measurement of muscle mass, muscle strength and physical performance, with a potential use for the diagnosis and follow-up of sarcopenia but they are not fully adapted for widespread use in clinical daily practice. The recommended tools for the diagnosis of sarcopenia in specialist clinical practice are DXA for the measurement of appendicular muscle mass, grip strength for the measurement of muscle strength and gait speed for the measurement of physical performance. Thresholds previously recommended in the literature can be applied to distinguish normal from abnormal;
- Healthcare providers, particularly in primary care, should consider an assessment of sarcopenia in individuals at increased risk; suggested tools for assessing risk include the SARC-F questionnaire, the SMI method or different prediction equations based on anthropometric data associated with the measurement of handgrip strength, although all of them require further validation;
- Whereas further studies are required to provide a full evidence-based guidance to clinicians, current management can include physical activity advice, particularly progressive resistance training, treatment and prevention of vitamin D deficiency and adequate energy and dietary protein intake.

The Expert group also emphasizes the importance of education and increased awareness of clinicians to the potential deleterious outcomes of sarcopenia.

Conclusions

Physicians and other health professionals have an important role to play in the assessment and management of sarcopenia to reduce its impact on individuals' well-being, the development of disability, and on health resources utilization.

Acknowledgements

None.

Funding

This meeting was founded by the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis, a Belgian not-for-profit organisation.

Availability of data and materials

Not applicable.

Authors' contributions

JYR organised the meeting. CC, MC, MV, EC, AC and OB performed the literature review. CB has drafted the manuscript. All authors (CB, EM, OB, MC, YR, RR, IAC, JAT, IB, MCB, MLB, NMAD, NB, EC, FC, AC, RF, EG, FL, JP, JYR, MV, JAK, CC) have taken part in the discussion and meeting and have critically analysed and approved the final manuscript.

Competing interests

Charlotte Beaudart: none; Eugene McCloskey has been involved with the development of the FRAX tool and with its continual update; has played a consultant/advisory role, and/or speaker and/or research funding for ActiveSignal, Amgen, ARUK, AstraZeneca, Consilient Healthcare, GSK, Hologic, i3 Innovus, Internis, IOF, Lilly, Medtronic, Merck, MRC, Novartis, Pfizer, Roche, Sanofi-Aventis, Servier, Synexus, Tethys, UCB, Unilever, and Warner Chilcott. Bruce H. Mitlak is an employee and has stock ownership in Eli Lilly and Company; Olivier Bruyère has received grant support from IBSA, Merck Sharp and Dohme, Nutraveris, Novartis, Pfizer, Rottapharm, Servier, and Theramex; lecture fees from IBSA, Rottapharm, Servier, and SMB; Matteo Cesari has received a research grant from Pfizer. Moreover, he has received honoraria for presentation at scientific meetings from Nestlé and Pfizer; Yves Rolland has received consulting and/or grant recipient from Amgen, Eli Lilly, Novartis and Lactalis; René Rizzoli has received consulting and

lecture fees from Merck Sharp and Dohme, Eli Lilly, Amgen, Novartis, Servier, Nycomed, Nestlé and Danone; Islène Araujo de Carvalho: none; Jotheeswaran Amuthavalli Thiyagarajan: none; Ivan Bautmans: none; Marie-Claude Berti  re : none; Maria Luisa Brandi has received consulting and/or grant recipient from Amgen, Eli Lilly, MSD, Novartis, NPS, Roche, and Servier; Nasser Al-Daghri: none; Nansa Burlet: none; Etienne Cavalier is a consultant for DiaSorin and IDS and has received lecture fees from IDS, DiaSorin, Roche, Abbott, Pfizer and Amgen; Francesca Cerreta: none; Antonio Cherubini is a partner of the SPRINT Consortium, which is partly funded by the European Federation of Pharmaceutical Industries and Associations (EFPIA); Roger A. Fielding has had remuneration, has played a consultant/advisory role, and has stock ownership of or funding from Eli Lilly, Dairy Management, Abbott, Pronutria, Segterra, Ammonett, Bristol Myers Squibb, Cytokinetics, Regeneron, Pfizer, Astellas, and Nestec; Evelien Gielen: none; Francesco Landi is a partner of the SPRINT Consortium, which is partly funded by the European Federation of Pharmaceutical Industries and Associations (EFPIA); Jean Petermans is the Treasurer of the EUGMS; Jean-Yves Reginster has received consulting fees or paid advisory boards for Servier, Novartis, Negma, Lilly, Wyeth, Amgen, GlaxoSmithKline, Roche, Merckle, NycomedTakeda, NPS, IBSA-Genevri  r, Theramex, UCB, Asahi Kasei, Endocyte; lecture fees from Merck Sharp and Dohme, Lilly, Rottapharm, IBSA, Genevri  r, Novartis, Servier, Roche, GlaxoSmithKline, Merckle, Teijin, Teva, Analis, Theramex, Nycomed, NovoNordisk, Ebewee Pharma, Zodiac, Danone, Will Pharma, Amgen; Grant support from Bristol Myers Squibb, Merck Sharp & Dohme, Rottapharm, Teva, Roche, Amgen, Lilly, Novartis, GlaxoSmithKline, Servier, Pfizer, Theramex, Danone, Organon, Therabel, Boehringer, Chiltern, Galapagos; Marjolein Visser: none; John A. Kanis has worked with and received funding from many companies and non-governmental organizations dealing with skeletal metabolism including research funding from the Health Technology Assessment NHS R&D HTA Programme of the UK; he has been involved with the development of the FRAX tool and with its continual update; Cyrus Cooper has received honoraria and consulting fees from Amgen, Glaxo SmithKline, ABBH, Merck Sharpe and Dohme, Eli Lilly, Pfizer, Novartis, Servier, Medtronic and Roche.

Consent for publication

Not applicable.

Ethical approval and consent to participate

This is a paper based on literature reviews and meetings. No participants were involved and the ethical approval is therefore not applicable.

Author details

¹Department of Public Health, Epidemiology and Health Economics, University of Li  ge, Avenue Hippocrate 13, CHU B23, 4000 Li  ge, Belgium. ²Centre for Metabolic Bone Diseases, University of Sheffield, Sheffield, UK. ³MRC and Arthritis Research UK Centre for Integrated research in Musculoskeletal Ageing (CIMA), London, UK. ⁴G  rontop  le, University Hospital of Toulouse, Toulouse, France. ⁵INSERM UMR1027, University of Toulouse III Paul Sabatier, Toulouse, France. ⁶G  rontop  le of Toulouse, University of Toulouse III, CHU Purpan, Toulouse, France. ⁷Service of Bone Diseases, Faculty of Medicine, Geneva University Hospitals, Geneva, Switzerland. ⁸World Health Organization, Geneva, Switzerland. ⁹Gerontology and Frailty in Ageing Research Department, Vrije Universiteit Brussel (VUB), Brussels, Belgium. ¹⁰Centre de Recherche et d'information Nutritionnelles, Paris, France. ¹¹Department of Surgery and Translational Medicine, University of Florence, viale Pieraccini 6, 59139 Florence, Italy. ¹²Prince Mutaib Chair for Biomarkers of Osteoporosis, Biochemistry Department, College of Science, King Saud University, Riyadh 11451, Saudi Arabia. ¹³Department of Clinical Chemistry, University of Li  ge, CHU Sart-Tilman, B  t B35, 4000 Li  ge, Belgium. ¹⁴Human Medicines Research and Development Support Division, Scientific Advice, London, UK. ¹⁵Geriatrics and Geriatric Emergency Care, IRCCS-INRCA, Ancona, Italy. ¹⁶Nutrition, Exercise Physiology and Sarcopenia Laboratory, Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, USA. ¹⁷Department of Geriatric Medicine, Katholieke Universiteit Leuven, Leuven, Belgium. ¹⁸Department of Geriatrics, Neurosciences and Orthopedics, Catholic University of the Sacred Heart Rome, Milano, Italy. ¹⁹Geriatric Department, CHU Sart-Tilman, B  t B35, 4000 Li  ge, Belgium. ²⁰Department of Health Sciences, VU University Amsterdam, Amsterdam, Netherlands. ²¹Department of Nutrition and Dietetics, Internal Medicine, VU University Medical Center, Amsterdam, Netherlands. ²²Institute for Health and Aging, Catholic University of Australia, Melbourne, Australia.

²³MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, England, UK. ²⁴NIHR Musculoskeletal Biomedical Research Unit, University of Oxford, Oxford, UK.

Received: 24 March 2016 Accepted: 29 September 2016

Published online: 05 October 2016

References

- Rosenberg IH. Sarcopenia: origins and clinical relevance. *J Nutr.* 1997; 127:990S–15S.
- Muscaritoli M, et al. Consensus definition of sarcopenia, cachexia and pre-cachexia: joint document elaborated by Special Interest Groups (SIG) 'cachexia-anorexia in chronic wasting diseases' and 'nutrition in geriatrics'. *Clin Nutr.* 2010;29:154–9.
- Cruz-Jentoft AJ, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing.* 2010;39:412–23.
- Fielding RA, et al. Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc.* 2011;12:249–56.
- Dam T-T, et al. An evidence-based comparison of operational criteria for the presence of sarcopenia. *J Gerontol A Biol Sci Med Sci.* 2014;69:584–90.
- Morley JE, et al. Sarcopenia with limited mobility: an International Consensus. *J Am Med Dir Assoc.* 2011;12:403–9.
- Studenski SA, et al. The FNIH sarcopenia project: rationale, study description, conference recommendations, and final estimates. *J Gerontol A Biol Sci Med Sci.* 2014;69:547–58.
- Baumgartner RN, et al. Epidemiology of sarcopenia among the elderly in New Mexico. *Am J Epidemiol.* 1998;147:755–63.
- Cooper C, et al. Frailty and sarcopenia: definitions and outcome parameters. *Osteoporos Int.* 2012;23:1839–48.
- Chen L-K, et al. Sarcopenia in Asia: Consensus Report of the Asian Working Group for Sarcopenia. *J Am Med Dir Assoc.* 2014;15:95–101.
- Cruz-Jentoft AJ, et al. Prevalence of and interventions for sarcopenia in ageing adults: a systematic review. Report of the International Sarcopenia Initiative (EWG SOP and IWGS). *Age Ageing.* 2014. doi:10.1093/ageing/afu115
- Bischoff-Ferrari HA, et al. Comparative performance of current definitions of sarcopenia against the prospective incidence of falls among community-dwelling seniors age 65 and older. *Osteoporos Int.* 2015. doi:10.1007/s00198-015-3194-y.
- Beaudart C, Rizzoli R, Bruy  re O, Reginster JY, Biver E. Sarcopenia: Burden and challenges for Public Health. 2014. *Archives of Public Health. Arch Public Heal.* 2014;72:45.
- Rizzoli R, et al. Quality of life in sarcopenia and frailty. *Calcif Tissue Int.* 2013;93:101–20.
- Lauretani F, et al. Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J Appl Physiol.* 2003;95:1851–60.
- Visser M, Schaap LA. Consequences of sarcopenia. *Clin Geriatr Med.* 2011;27:387–99.
- Lang T, et al. Sarcopenia: etiology, clinical consequences, intervention, and assessment. *Osteoporos Int.* 2010;21:543–59.
- Landi F, et al. Sarcopenia and frailty: From theoretical approach into clinical practice. 2016. <http://emvmsa1a.jouue-hdi.com/en/article/1029882>.
- Landi F, et al. Sarcopenia as a risk factor for falls in elderly individuals: results from the iLSIRENTE study. *Clin Nutr.* 2012;31:652–8.
- Denison HJ, Cooper C, Sayer AA, Robinson SM. Prevention and optimal management of sarcopenia: a review of combined exercise and nutrition interventions to improve muscle outcomes in older people. *Clin Interv Aging.* 2015;10:859–69.
- Molfino A, Gioia G, Rossi Fanelli F, Muscaritoli M. Beta-hydroxy-beta-methylbutyrate supplementation in health and disease: a systematic review of randomized trials. *Amino Acids.* 2013;45:1273–92.
- Devries MC, Phillips SM. Creatine supplementation during resistance training in older adults: a meta-analysis. *Med Sci Sports Exerc.* 2014;46:1194–203.
- Cooper C, et al. Tools in the assessment of sarcopenia. *Calcif Tissue Int.* 2013;93:201–10.
- Mijnarends DM, et al. Validity and reliability of tools to measure muscle mass, strength, and physical performance in community-dwelling older people: a systematic review. *J Am Med Dir Assoc.* 2013;14:170–8.

25. Hiligsmann M, et al. A reference case for economic evaluations in osteoarthritis: An expert consensus article from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Semin Arthritis Rheum*. 2014;44:271–82.
26. Rizzoli R, et al. The role of dietary protein and vitamin D in maintaining musculoskeletal health in postmenopausal women: a consensus statement from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Maturitas*. 2014;79:122–32.
27. Bruyère O, et al. An algorithm recommendation for the management of knee osteoarthritis in Europe and internationally: a report from a task force of the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Semin Arthritis Rheum*. 2014;44:253–63.
28. Rizzoli R, et al. Management of osteoporosis of the oldest old. *Osteoporos Int*. 2014;25:2507–29.
29. Arden N, et al. Can We Identify Patients with High Risk of Osteoarthritis Progression Who Will Respond to Treatment? A Focus on Biomarkers and Frailty. *Drugs Aging*. 2015;32:525–35.
30. Cooper C, et al. How to define responders in osteoarthritis. *Curr Med Res Opin*. 2013;29:719–29.
31. Rizzoli R, et al. Vitamin D supplementation in elderly or postmenopausal women: a 2013 update of the 2008 recommendations from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Curr Med Res Opin*. 2013;29:305–13.
32. Kaufman J-M, et al. Treatment of osteoporosis in men. *Bone*. 2013;53:134–44.
33. Rizzoli R, et al. Guidance for the prevention of bone loss and fractures in postmenopausal women treated with aromatase inhibitors for breast cancer: an ESCEO position paper. *Osteoporos Int*. 2012;23:2567–76.
34. Reginster J-Y, et al. Recommendations for an update of the 2010 European regulatory guideline on clinical investigation of medicinal products used in the treatment of osteoarthritis and reflections about related clinically relevant outcomes: expert consensus statement. *Osteoarthritis Cartilage*. 2015. doi:10.1016/j.joca.2015.07.001.
35. Reginster J-Y, et al. Recommendations for the conduct of clinical trials for drugs to treat or prevent sarcopenia. *Aging Clin Exp Res*. 2015;28:47–58.
36. Hunter S, White M, Thompson M. Techniques to evaluate elderly human muscle function: a physiological basis. *J Gerontol Biol Sci*. 1998;53:2–4.
37. Beaudart C, et al. Estimation of sarcopenia prevalence using various assessment tools. *Exp Gerontol*. 2014;61C:31–7.
38. Cavalier E, Beaudart C, Buckinx F, Bruyère O, Reginster J-Y. Critical analytical evaluation of promising markers for sarcopenia. 2015. <http://www.em-consulte.com/en/article/1021209>.
39. Mitsiopoulos N, et al. Cadaver validation of skeletal muscle measurement by magnetic resonance imaging and computerized tomography. *J Appl Physiol*. 1998;85:115–22.
40. Levine JA, et al. Measuring leg muscle and fat mass in humans: comparison of CT and dual-energy X-ray absorptiometry. *J Appl Physiol*. 2000;88:452–6.
41. Visser M, Fuerst T, Lang T, Salamone L, Harris TB. Validity of fan-beam dual-energy X-ray absorptiometry for measuring fat-free mass and leg muscle mass. Health, Aging, and Body Composition Study–Dual-Energy X-ray Absorptiometry and Body Composition Working Group. *J Appl Physiol*. 1999;87:1513–20.
42. Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc*. 2002;50:889–96.
43. Rush EC, Freitas J, Plank LD. Body size, body composition and fat distribution: comparative analysis of European, Maori, Pacific Island and Asian Indian adults. *Br J Nutr*. 2009;102:632–41.
44. Bruyère O, et al. Assessment of muscle mass, muscle strength and physical performance in clinical practice: An international survey. *Eur Geriatr Med*. 2016;7:243–6.
45. Landi F, et al. Midarm muscle circumference, physical performance and mortality: results from the aging and longevity study in the Sirente geographic area (ISIRENTE study). *Clin Nutr*. 2010;29:441–7.
46. Landi F, et al. Calf circumference, frailty and physical performance among older adults living in the community. *Clin Nutr*. 2014;33:539–44.
47. Rolland Y, et al. Sarcopenia, calf circumference, and physical function of elderly women: a cross-sectional study. *J Am Geriatr Soc*. 2003;51:1120–4.
48. Heymsfield SB, McManus C, Smith J, Stevens V, Nixon DW. Anthropometric measurement of muscle mass: revised equations for calculating bone-free arm muscle area. *Am J Clin Nutr*. 1982;36:680–90.
49. Kulkarni B, et al. Development and validation of anthropometric prediction equations for estimation of lean body mass and appendicular lean soft tissue in Indian men and women. *J Appl Physiol*. 2013;115:1156–62.
50. Yu S, et al. An anthropometric prediction equation for appendicular skeletal muscle mass in combination with a measure of muscle function to screen for sarcopenia in primary and aged care. *J Am Med Dir Assoc*. 2015;16:25–30.
51. Villani AM, et al. Appendicular skeletal muscle in hospitalised hip-fracture patients: development and cross-validation of anthropometric prediction equations against dual-energy X-ray absorptiometry. *Age Ageing*. 2014;43:857–62.
52. Reiss J, et al. Case finding for sarcopenia in geriatric inpatients: performance of bioimpedance analysis in comparison to dual X-ray absorptiometry. *BMC Geriatr*. 2016;16:52.
53. Kim M, Kim H. Accuracy of segmental multi-frequency bioelectrical impedance analysis for assessing whole-body and appendicular fat mass and lean soft tissue mass in frail women aged 75 years and older. *Eur J Clin Nutr*. 2013;67:395–400.
54. Kim M, Shinkai S, Murayama H, Mori S. Comparison of segmental multifrequency bioelectrical impedance analysis with dual-energy X-ray absorptiometry for the assessment of body composition in a community-dwelling older population. *Geriatr Gerontol Int*. 2015;15:1013–22.
55. Buckinx F, et al. Concordance between muscle mass assessed by bioelectrical impedance analysis and by dual energy X-ray absorptiometry: a cross-sectional study. *BMC Musculoskelet Disord*. 2015;16:60.
56. Stevens PJ, et al. Is grip strength a good marker of physical performance among community-dwelling older people? *J Nutr Heal Aging*. 2012;16:769–74.
57. Bohannon RW, Magasi SR, Bubela DJ, Wang Y-C, Gershon RC. Grip and knee extension muscle strength reflect a common construct among adults. *Muscle Nerve*. 2012;46:555–8.
58. Roberts HC, et al. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age Ageing*. 2011;40:423–9.
59. Bean JF, et al. The relationship between leg power and physical performance in mobility-limited older people. *J Am Geriatr Soc*. 2002;50:461–7.
60. Dodds RM, et al. Grip strength across the life course: normative data from twelve British studies. *PLoS One*. 2014;9:e113637.
61. Anjum SN, Choudary P, Dimri R, Ankarath S. Comparative evaluation of grip and pinch strength in an Asian and European population. *Hand Ther*. 2012;17:11–4.
62. Brown M, Sinacore DR, Binder EF, Kohrt WM. Physical and performance measures for the identification of mild to moderate frailty. *J Gerontol A Biol Sci Med Sci*. 2000;55:M350–5.
63. Callahan D, Phillips E, Carabello R, Frontera WR, Fielding RA. Assessment of lower extremity muscle power in functionally-limited elders. *Aging Clin Exp Res*. 2007;19:194–9.
64. Jones CJ, Rikli RE, Beam WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exerc Sport*. 1999;70:113–9.
65. Karpman C, Lebrasseur NK, Depew ZS, Novotny PJ, Benzo RP. Measuring gait speed in the out-patient clinic: methodology and feasibility. *Respir Care*. 2014;59:531–7.
66. National Institute on Aging, Laboratory of Epidemiology, Demography, and Biometry. <https://www.nia.nih.gov/>.
67. Bean JF, Kiely DK, LaRose S, Alian J, Frontera WR. Is Stair Climb Power a Clinically Relevant Measure of Leg Power Impairments in At-Risk Older Adults? *Arch Phys Med Rehabil*. 2007;88:604–9.
68. Guigoz Y, Vellas B, Garry PJ. Assessing the nutritional status of the elderly: The Mini Nutritional Assessment as part of the geriatric evaluation. *Nutr Rev*. 1996;54:S59–65.
69. Malmstrom TK, Morley JE. SARC-F: a simple questionnaire to rapidly diagnose sarcopenia. *J Am Med Dir Assoc*. 2013;14:531–2.
70. Woo J, Leung J, Morley JE. Validating the SARC-F: a suitable community screening tool for sarcopenia? *J Am Med Dir Assoc*. 2014;15:630–4.
71. Goodman MJ, et al. Development of a practical screening tool to predict low muscle mass using NHANES 1999–2004. *J Cachexia Sarcopenia Muscle*. 2013;4:187–97.
72. Visvanathan R, et al. Appendicular skeletal muscle mass: development and validation of anthropometric prediction equations. 2012. <https://digital.library.adelaide.edu.au/dspace/handle/2440/76238>.
73. Ishii S, et al. Development of a simple screening test for sarcopenia in older adults. *Geriatr Gerontol Int*. 2014;14 Suppl 1:93–101.

74. Sayer AA, et al. Type 2 Diabetes, Muscle Strength, and Impaired Physical Function: The tip of the iceberg? *Diabetes Care*. 2005;28:2541–2.
75. Reginster J-Y, Beaudart C, Buckinx F, Bruyère O. Osteoporosis and sarcopenia: two diseases or one? *Curr Opin Clin Nutr Metab Care*. 2015. doi:10.1097/MCO.0000000000000230.
76. Kalyani RR, Corriere M, Ferrucci L. Age-related and disease-related muscle loss: the effect of diabetes, obesity, and other diseases. *Lancet Diabetes Endocrinol*. 2014;2:819–29.
77. Peterson MD, Sen A, Gordon PM. Influence of resistance exercise on lean body mass in aging adults: a meta-analysis. *Med Sci Sport Exerc*. 2011;43:249–58.
78. Sahni S, Mangano KM, Hannan MT, Kiel DP, McLean RR. Higher Protein Intake Is Associated with Higher Lean Mass and Quadriceps Muscle Strength in Adult Men and Women. *J Nutr*. 2015;145:1569–75.
79. Genaro Pde S, Pinheiro Mde M, Szejnfeld VL, Martini LA. Dietary protein intake in elderly women: association with muscle and bone mass. *Nutr Clin Pract*. 2015;30:283–9.
80. Bauer J, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. *J Am Med Dir Assoc*. 2013;14:542–59.
81. Deer RR, Volpi E. Protein intake and muscle function in older adults. *Curr Opin Clin Nutr Metab Care*. 2015;18:248–53.
82. Gryson C, et al. Four-month course of soluble milk proteins interacts with exercise to improve muscle strength and delay fatigue in elderly participants. *J Am Med Dir Assoc*. 2014;15:958.e1–9.
83. Finger D, et al. Effects of protein supplementation in older adults undergoing resistance training: a systematic review and meta-analysis. *Sports Med*. 2015;45:245–55.
84. Beaudart C, et al. The effects of vitamin D on skeletal muscle strength, muscle mass and muscle power: a systematic review and meta-analysis of randomized controlled trials. *J Clin Endocrinol Metab* jc20141742 (2014). doi:10.1210/jc.2014-1742

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

