
Physiology and Molecular Biology of Trace Element Hyperaccumulation

Sylvain Merlot, Vanesa Sanchez Garcia de la Torre,
and Marc Hanikenne

Abstract

Some trace elements are essential for plants but become toxic at high concentration. Remarkably, about 500 plant species worldwide are able to accumulate tremendous amounts of metals in their leaves and are therefore called metal hyperaccumulators. In the context of sustainable development, there is a regain of interest for metal hyperaccumulation mechanisms that may become instrumental for improving metal phytoextraction from contaminated soils to produce metals with a lower net impact on the environment. In addition, studying the molecular mechanisms of hyperaccumulation in diverse plant species is necessary in order to understand the evolution of this extreme and complex adaptation trait. Our current knowledge of metal hyperaccumulation is mostly based on the analysis of a few species from the Brassicaceae family, and suggests that the underlying mechanisms result from an exaggeration of basic mechanisms involved in metal homeostasis. However, the development of Next Generation Sequencing technologies opens today the possibility for studying new hyperaccumulator species that therefore may reveal more diversity in these mechanisms. The goal of this chapter is to provide background information on metal hyperaccumulation and give a clear picture of what we know currently about the molecular mechanisms involved in this trait. We also attempt to outline for the reader the future scientific challenges that this field of research is facing.

S. Merlot (✉) • V. Sanchez Garcia de la Torre
Institute for Integrative Biology of the Cell (I2BC), CEA,
CNRS, Univ. Paris-Sud, Université Paris-Saclay, Gif-sur-
Yvette cedex, France
e-mail: sylvain.merlot@i2bc.paris-saclay.fr

M. Hanikenne
InBioS - Functional Genomics and Plant Molecular
Imaging, and PhytoSYSTEMS, University of Liège,
Liège, Belgium

1 Introduction

Investigating the mechanisms involved in metal hyperaccumulation allows us to observe extreme adaptation of metal homeostasis networks in plants and identify key players in metal distribution and tolerance in plant tissues. The study of metal hyperaccumulator species also permits examination of mechanisms underlying the evolution of this extreme adaptive trait (Shahzad et al. 2010; Hanikenne and Nouet 2011; Hanikenne et al. 2013). Several comprehensive reviews have been published recently on this topic (Verbruggen et al. 2009b; Krämer 2010; Hanikenne and Nouet 2011) and we refer readers to those reviews. Whereas hyperaccumulator species typically accumulate specifically one metal when growing in their natural environment, some species have the ability to tolerate and accumulate several metals when grown *ex situ*. This is well documented for the hyperaccumulator species of the Brassicaceae family *Noccaea caerulescens*, in which ultramafic-adapted accessions such as Puy de Wolf (France), Monte Prinzera (Italy), or Puente Basadre (Spain) are able to accumulate Ni but also Zn and Cd (Assunção et al. 2003; Peer et al. 2003; Escarré et al. 2013; Gonneau et al. 2014; Callahan et al. 2016). This ability to tolerate and accumulate several metals likely reflects the relatively low specificity of some mechanisms involved in metal transport and chelation.

Several metals that are accumulated in hyperaccumulator species (e.g. Zn, Ni, Mn) are essential nutrients but become toxic at high concentrations for most plants (i.e. non-accumulating species). Therefore, all plant species have developed mechanisms to regulate essential metal homeostasis according to their needs and metal availability in soils (Burkhead et al. 2009; Palmer and Guerinot 2009; Thomine and Vert 2013). Our current knowledge suggests that the molecular mechanisms involved in metal hyperaccumulation are derived essentially from the mechanisms involved in metal homeostasis. In several examples, genes involved in metal homeostasis are differentially expressed in hyperaccumulators compared to related non-accumulator species, as a result of

gene duplication and/or changes in promoter activity (Talke et al. 2006; van de Mortel et al. 2006; Krämer et al. 2007; Hanikenne et al. 2008; Shahzad et al. 2010). However, specific genes linked to hyperaccumulation may be discovered as molecular analysis of hyperaccumulation will extend in the future to additional non-model species from various plant families owing to the development of high-throughput sequencing technologies (Verbruggen et al. 2013; Halimaa et al. 2014b; Merlot et al. 2014). For most metals, the hyperaccumulation trait appeared independently in distant plant families. Therefore, some of the mechanisms involved in metal hyperaccumulation may be specific to a plant family or a species, whereas others may be convergent among distant hyperaccumulators. For instance, several examples of convergent evolution have been identified between the Brassicaceae *Arabidopsis halleri* and *N. caerulescens* (see below and Krämer et al. 2007; Hanikenne et al. 2008; O’Lochlainn et al. 2011; Craciun et al. 2012), suggesting important functional constraints in the metal homeostasis network.

Because of its singularity, metal hyperaccumulation may appear as an exception having minimal relevance. However, from a scientific point of view, metal hyperaccumulation in plants is fascinating, and understanding the mechanisms involved in this trait may provide tools for producing metals with lower impact on the environment in the near future. The goal in this chapter is to outline our current knowledge about the molecular mechanisms of metal hyperaccumulation in plants and to highlight possible future developments in this important field of research.

2 Molecular Physiology of Metal Hyperaccumulation

2.1 Main Steps of Metal Hyperaccumulation

When exposed to excess metals, most plant species adopt a so-called excluder strategy to prevent metal accumulation in photosynthetically active shoot tissues (Krämer 2010). This goal

can be achieved by limiting metal absorption by roots, increasing metal efflux from root tissues, and/or increasing metal storage in root cell walls and vacuoles. In contrast, achieving metal hyperaccumulation and hypertolerance requires modifications at specific nodes of the metal homeostasis network to ensure that the metal flux in the plant is directed towards shoot tissues (Fig. 1; Clemens et al. 2002). At the

physiological level, these alterations include some or all of the following steps:

1. An enhanced metal mobilization and uptake in roots;
2. An efficient radial metal transport towards the root vascular tissues; this includes a reduction of metal storage in root vacuoles;
3. An increased transport of metal from the root to the shoot, with efficient xylem loading; this




Fig. 1 Model of the physiology of metal hyperaccumulation and hypertolerance. Enhanced metal uptake and radial transport in roots, xylem loading/unloading and vacuolar storage in shoots all make major contributions to the traits. In roots, it is possible that the metal can travel in the apoplast up to the endodermis cell layer before cellular uptake (not represented). Note that

the tissue (epidermis or mesophyll) involved in metal storage varies depending on the species and the metal (see text). *co* cortex, *en* endodermis, *ep* epidermis, *me* mesophyll, *Zinc-NA* Zinc-Nicotianamine chelates, *pc* pericycle, *vac* vacuole, *xp* xylem parenchyma, *xy* xylem. Figure modified from Hanikenne and Nouet (2011)

step contributes to metal tolerance by enabling metal storage (and thus detoxification) in shoot tissues;

4. An efficient mechanism for xylem unloading and metal distribution in shoots together with a high vacuolar storage capacity.

Processes occurring in both roots and shoots thus contribute to metal tolerance and hyperaccumulation. Root-based processes play an essential role in hyperaccumulation whereas shoot-based processes are required for hypertolerance, as suggested from grafting experiments between *N. caerulescens* and the non-Zn-accumulating *Microthlaspi perfoliatum* (Guimarães et al. 2009).

2.2 Metal Distribution in Shoots

Metal distribution in shoot tissues is specific to both the species and the metal considered. This topic has been extensively reviewed recently (Fernando et al. 2013; Leitenmaier and Küpper 2013). Briefly, in most cases, metals (Zn, Cd, Ni or Se) accumulate at the base of the trichomes and in the vacuoles of epidermal cells. In contrast, mesophyll cells that are the main site of photosynthesis accumulate lower amounts of metals (Küpper et al. 1999, 2001; Lombi et al. 2002; Cosio et al. 2005). In the vacuoles of epidermal cells, metals can reach very high concentrations (e.g. several hundred mM; Küpper et al. 1999; Fernando et al. 2006b). There are, however, exceptions. For instance, Zn and Cd are stored in the vacuoles of mesophyll cells in both Zn- and Cd-hyperaccumulators *A. halleri* and *Sedum alfredii*, in which Zn is mostly bound to malate (Küpper et al. 2000; Sarret et al. 2002, 2009; Tian et al. 2011; Lu et al. 2014; Isaure et al. 2015). In the hyperaccumulator *Sedum plumbizincola*, Zn accumulates mostly in leaf epidermal cells, but also in large amounts in mesophyll cells of young leaves (Cao et al. 2014). In this species, Cd is mostly bound to cell walls in leaves (Peng et al. 2017). Accumulation of Mn in mesophyll

cells is also observed in several Mn-hyperaccumulators (Fernando et al. 2006a, b, 2013).

3 Identification of Molecular Processes Involved in Metal Hyperaccumulation

In the last 15 years, a number of complementary approaches have been used to identify the molecular actors underlying hyperaccumulation and hypertolerance. These approaches included:

- I. Screens of cDNA libraries in yeast in order to isolate genes contributing to metal transport and tolerance (e.g. Lasat et al. 2000; Pence et al. 2000; Bernard et al. 2004; Papoyan and Kochian 2004);
- II. Quantitative genetics analyses aiming to identify Quantitative Traits Loci (QTLs) co-segregating with the traits in progenies of crosses between an hyperaccumulator and a related non-accumulator species (e.g. Dräger et al. 2004; Deniau et al. 2006; Filatov et al. 2007; Courbot et al. 2007; Willems et al. 2007, 2010; Frérot et al. 2010; Baliardini et al. 2015).
- III. Transcriptomic studies comparing gene expression levels in hyperaccumulator and related non-accumulator species (e.g. Becher et al. 2004; Weber et al. 2004, 2006; Chiang et al. 2006; Craciun et al. 2006; Filatov et al. 2006; Hammond et al. 2006; Talke et al. 2006; van de Mortel et al. 2006; van de Mortel et al. 2008; Gao et al. 2013; Han et al. 2015).

These candidate genes are mostly involved in metal transport, metal chelator synthesis, or metal-induced oxidative stress response. Note that the last is not discussed in this chapter. Several candidate genes were further characterized functionally (e.g. Pence et al. 2000; Persans et al. 2001; Dräger et al. 2004; Kim et al. 2004; Hanikenne et al. 2008; Gustin et al. 2009; Lin et al. 2009; Shahzad et al. 2010; Ueno et al. 2011; Milner et al. 2012; Deinlein et al. 2012; Merlot et al. 2014; Baliardini et al. 2015; Nouet et al. 2015; Charlier et al. 2015). However, only a few

candidates (e.g. *HMA4*, see below) were confirmed by reverse genetics in hyperaccumulator species as major players in metal hyperaccumulation and tolerance; their functions are described in detail in the following sections.

A large part of our knowledge on metal hyperaccumulation comes from the study of two model Zn and Cd hyperaccumulating species of the Brassicaceae family, *A. halleri* and *N. caerulescens*, which are related to the sensitive and non-accumulating species *A. thaliana* (Yogeeswaran et al. 2005; Clauss and Koch 2006). These two species have been instrumental in successfully improving our understanding of the physiological, molecular, and genetic bases of metal hyperaccumulation and associated hypertolerance (see Krämer et al. 2007; Milner and Kochian 2008; Pauwels et al. 2008; Roosens et al. 2008; Verbruggen et al. 2009b; Krämer 2010; Hanikenne and Nouet 2011). Those successes relied on availability of the *A. thaliana* genome sequence (The Arabidopsis Genome Initiative 2000), and on dedicated tools and resources combined with relatively high gene sequence conservation among Brassicaceae species (94% and 88% identity with *A. thaliana* for *A. halleri* and *N. caerulescens*, respectively) (Talke et al. 2006; van de Mortel et al. 2006). It is expected that our knowledge will rapidly broaden with new species becoming accessible to molecular and genomic analyses (Gao et al. 2013; Verbruggen et al. 2013; Merlot et al. 2014).

4 Mechanisms of Zn and Cd Hyperaccumulation

As mentioned above, most of our knowledge about Zn and Cd hyperaccumulation was acquired using *A. halleri* and *N. caerulescens* (Fig. 2). More recent models include (i) the Crassulaceae *S. alfredii* and *S. plumbizincola* from Asia, which are the only species reported to hyperaccumulate Cd other than the Brassicaceae (Yang et al. 2004, 2006; Deng et al. 2007; Krämer 2010; Wu et al. 2013; Cao et al. 2014; Ma et al. 2015; Peng et al. 2017); and

(ii) the Amaranthaceae *Gomphrena claussenii* from South America, which is highly tolerant to Zn and Cd and presents indicator levels of Zn and Cd accumulations (Villafort Carvalho et al. 2013, 2015).

Arabidopsis halleri and *N. caerulescens* display constitutive Zn hyperaccumulation and hypertolerance, although intraspecific variation for those traits has been reported (Bert et al. 2000, 2002; Reeves et al. 2001; Assunção et al. 2003; Molitor et al. 2005; Besnard et al. 2009). Similarly, hyperaccumulation of Cd shows substantial intraspecific variation (Escarré et al. 2000; Bert et al. 2002; Roosens et al. 2003; Verbruggen et al. 2013; Meyer et al. 2015). Metal hyperaccumulation evolved independently in the two species (Krämer 2010). However, both share a set of alterations of their metal homeostasis networks in comparison to the non-accumulator *A. thaliana*, which is described in detail below.

4.1 Uptake of Zn and Cd

Prior to uptake, it is suggested that metals are actively mobilized from the soil, by acidification and/or chelate secretion (Clemens et al. 2002). A recent report, however, suggested that the roots of *A. halleri* secrete elevated levels of nicotianamine (NA), a metal chelator able to form NA-Zn complexes (Curie et al. 2009; Clemens et al. 2013), which may reduce root Zn uptake and increase tolerance (Tsednee et al. 2014). Another report indicated higher organic acid levels and Zn mobilization in the dissolved organic matter within the rhizosphere of hyperaccumulator compared to non-accumulator accessions of *S. alfredii* (Li et al. 2012).

Several divalent metal transporters of the ZIP (Zrt-Irt-like Protein) family are highly expressed in roots and/or shoots of both *A. halleri* and *N. caerulescens* (Talke et al. 2006; Krämer et al. 2007; Lin et al. 2009, 2016). It presumably results in enhanced rates of root metal uptake or mobilization from root storage sites. By contributing to Zn radial transport towards the xylem in roots, it also may contribute to metal partitioning between root and shoot tissues. Several




Fig. 2 Model for Zn hyperaccumulation and hypertolerance in the Brassicaceae *A. halleri* and *N. caerulescens*. Enhanced functions of ZIP transporters in cellular uptake, of the P-type ATPase HMA4 in xylem loading/unloading and of MTP1 in vacuolar storage all make major contributions to the traits. The exact functions and localizations of individual ZIPs are unknown. In roots, the metal chelator nicotianamine (NA) possibly favours Zn radial transport towards the xylem by symplastic inter-cellular mobility of Zn by either allowing Zn movement through plasmodesmata and/or preventing vacuolar storage. Vacuolar storage in

shoots occurs in the epidermis in *N. caerulescens* and in the mesophyll in *A. halleri*. Additional metal homeostasis genes that are highly expressed in both hyperaccumulators are discussed in the text. Note that similar mechanisms have been identified as more active in a metal hyperaccumulating population of the Crassulaceae *S. alfredii* compared to a non-accumulating population. *co* cortex, *en* endodermis, *ep* epidermis, *me* mesophyll, *Zinc-NA* Zinc-Nicotianamine chelates, *pc* pericycle, *vac* vacuole, *xp* xylem parenchyma, *xy* xylem. Figure modified from Hanikenne and Nouet (2011)

ZIP genes are induced by Zn deficiency under the control of the bZIP19 and bZIP23 transcription factors in *A. thaliana* (Assunção et al. 2010). Their high expression in *A. halleri* and *N. caerulescens* roots could be the direct

consequence of the high activity of HMA4 (Heavy Metal ATPase 4; see below), which depletes Zn in roots (Talke et al. 2006; Hanikenne et al. 2008; Gustin et al. 2009). Note that several ZIP genes are also highly expressed

in *S. plumbizincola* (Peng et al. 2017). Further work will be required to determine their individual function in Zn hyperaccumulation and to assess whether Cd uptake is determined by specific mechanisms or via Zn- and/or Fe-transport mechanisms (Meyer and Verbruggen 2012).

4.2 Root-to-Shoot Transfer of Zn and Cd

Increased rate of root-to-shoot metal transfer is key to achieving metal hyperaccumulation in shoots. It requires enhanced radial transport to xylem, decreased vacuolar storage in root cells, and efficient xylem loading. Several *NAS* (nicotianamine synthase) genes are highly expressed in *A. halleri* and *N. caerulescens* (Weber et al. 2004; van de Mortel et al. 2006; Deinlein et al. 2012). *NAS* transcript levels are also higher in roots of a hyperaccumulator accession compared to those of a non-hyperaccumulator accession in *S. alfredii* (Liang et al. 2014). Elevated levels of NA have been measured in roots of *A. halleri* compared to *A. thaliana* (Weber et al. 2004; Deinlein et al. 2012). It was further shown, using *A. halleri* RNAi lines, that high expression of the *NAS2* gene provides increased NA levels for Zn symplastic mobility towards the xylem and for controlling the rate of Zn xylem loading in roots (Deinlein et al. 2012; Cornu et al. 2015). The amino-acid histidine (His), whose concentration weakly correlates with Zn content in *N. caerulescens*, was shown to enhance Zn xylem loading and thus contribute to reduce Zn storage in roots (Callahan et al. 2007; Kozhevnikova et al. 2014).

In *A. halleri*, Zn and Cd loading into the xylem is driven by the HMA4 protein (Talke et al. 2006; Courbot et al. 2007; Hanikenne et al. 2008), which is a plasma membrane P-Type ATPase pump that uses the energy released from the hydrolysis of Adenosine triphosphate (ATP) to transport metal against the electro-chemical gradient (Hussain et al. 2004; Wong and Cobbett 2009; Pedersen et al. 2012; Hanikenne and Baurain 2014). The *HMA4* gene co-segregates with QTLs for Zn and Cd tolerance and accumulation (Courbot et al. 2007; Willems et al. 2007, 2010; Frérot et al.

2010; Meyer et al. 2016). High expression of *HMA4* is required for both hyperaccumulation and hypertolerance in *A. halleri* (Talke et al. 2006; Hanikenne et al. 2008). Increased gene dosage of *HMA4* was selected during the evolutionary history of *A. halleri* and evolved through tandem triplication and activation in *cis* of the promoters of all three copies (Hanikenne et al. 2008, 2013). The *A. halleri* *HMA4* locus was shaped by positive selection, resulting in a selective sweep and in ectopic gene conversion (Hanikenne et al. 2013). The three *HMA4* copies are active mainly in vascular tissues of *A. halleri*, which allows acting in xylem metal loading in roots and possibly in metal distribution in leaves. It also may ensure metal exclusion from metal-sensitive tissues (e.g. root tip, cambium). By controlling highly active Zn xylem loading, HMA4 also acts as a physiological regulator: it depletes the root Zn pool, which triggers a Zn-deficiency response resulting in high expression of several *ZIP* genes (Hanikenne et al. 2008). In agreement, modeling of the Zn supply-dependent spatio-temporal evolution of Zn concentration in root symplast and apoplast of *A. thaliana* predicted that slight changes in *HMA4* transcript levels have a major impact on the radial distribution of Zn in roots and the root-to-shoot Zn gradient (Claus et al. 2013). It was further shown that a certain extent of functional differentiation exists among the three *AhHMA4* copies when expressed in *A. thaliana*, stemming from differences in expression levels rather than in expression profile. Interestingly, *AhHMA4* copy 3 was subjected to the strongest, possibly most recent, positive selection during the evolutionary history of *A. halleri* (Hanikenne et al. 2013), thus linking sequence diversity patterns and function *in vivo* (Hanikenne et al. 2013; Nouet et al. 2015).

HMA4 is also highly expressed in *N. caerulescens* as well as in Zn- and Cd-hyperaccumulator accessions of *S. alfredii* and *S. plumbizincola*, where it very likely plays similar roles to the *A. halleri* *HMA4* (Bernard et al. 2004; Papoyan and Kochian 2004; van de Mortel et al. 2006; O’Lochlainn et al. 2011; Craciun et al. 2012; Zhang et al. 2016; Peng et al. 2017). Moreover, the gene coding for the ZIP transporter *ZNT1* of *N. caerulescens* is highly

expressed in cortex, endodermis, and pericycle root cells. When expressed in *A. thaliana*, it contributes to Zn and Cd tolerance and accumulation. The *NcZNT1* gene may therefore be involved in Zn and Cd influx into cells responsible for xylem loading, providing metals for transport by HMA4 (Milner et al. 2012; Lin et al. 2016). The ortholog of *ZNT1* in *A. halleri*, *ZIP4*, is also highly expressed and may contribute to a similar function (Talke et al. 2006). Note that NRAMP1 (Natural Resistance-Associated Macrophage Protein 1) may also play a similar role for Cd in *N. caerulescens* (Milner et al. 2014). Moreover, the vacuolar metal efflux transporters NRAMP3 and NRAMP4 are highly expressed in *N. caerulescens* and *A. halleri* roots and were proposed to limit vacuolar storage and increase metal mobility (Weber et al. 2004; Oomen et al. 2009). NRAMP3 is also highly expressed in *S. plumbizincola* (Peng et al. 2017).

Once in the xylem sap, metals are transported to the shoot owing to the evapo-transpiration stream. In this compartment, Zn is mainly bound to organic acids such as malate and citrate (Monsant et al. 2011; Lu et al. 2013; Cornu et al. 2015).

4.3 Storage of Zn and Cd in Leaves

It is suggested that HMA4 and ZIP transporters play an important role in Zn unloading and distribution in shoot tissues (Krämer et al. 2007; Hanikenne and Nouet 2011). However, their exact contribution, as well as that of metal ligands or other transporters, to these processes remains to be detailed. Zn storage in vacuoles is most likely ensured by the MTP1 (Metal Tolerance Protein 1) protein in *A. halleri* (Dräger et al. 2004; Talke et al. 2006; Shahzad et al. 2010), although its role remains to be formally established through the analysis of knock-down plants. MTP1 is a vacuolar transporter implicated in Zn tolerance (Krämer 2005). The MTP1 gene is constitutively highly expressed in both root and shoot of *A. halleri*, and is present in four to five copies that are located on three distinct linkage groups in the genome (Dräger et al. 2004;

Talke et al. 2006; Willems et al. 2007; Shahzad et al. 2010). The two most highly expressed copies each co-segregate with QTLs for Zn tolerance (Dräger et al. 2004; Talke et al. 2006; Willems et al. 2007; Shahzad et al. 2010).

MTP1 is also highly expressed in *Noccaea* (formerly *Thlaspi*) *goesingense*, another Zn and Ni hyperaccumulator, in *N. caerulescens*, and in Zn-hyperaccumulating populations of *S. alfredii* (Milner and Kochian 2008; Gustin et al. 2009; Zhang et al. 2011). It likely plays in these species a similar role as in *A. halleri*.

Finally, no detailed information is currently available on the molecular mechanisms of Cd storage in *A. halleri* shoot vacuoles (Meyer and Verbruggen 2012). Indeed, MTP1 is not associated with high Cd tolerance or accumulation in *A. halleri* (Courbot et al. 2007; Willems et al. 2010), although MTP1-related proteins were shown to transport Cd in several species, including the Ni hyperaccumulator *N. goesingense* (Persans et al. 2001; Migocka et al. 2015). In *N. caerulescens*, HMA3 (Heavy Metal ATPase 3) may contribute to this process (Ueno et al. 2011). In addition, differential regulation of *MTP1* by Cd in related *Noccaea* species suggests that it may contribute to Cd handling (Martos et al. 2016).

4.4 Additional Candidate Genes for Zn and Cd Accumulation and Tolerance

A few additional candidate genes for a role in Zn or Cd tolerance have been functionally characterized. In *A. halleri*, the PDF1.1 (Plant Defensin 1.1) protein was identified through a cDNA screen in yeast as a contributor to Zn tolerance. It also confers Zn tolerance when ectopically over-expressed in *A. thaliana* and is more highly expressed in shoots of *A. halleri* compared to *A. thaliana* (Mirouze et al. 2006). It localizes in intracellular compartments (Oomen et al. 2011). PDFs were initially known as secreted antifungal proteins and are characterized by a cysteine-stabilized, α -helix β -sheet, three-dimensional structure (De Coninck et al. 2013; van der

Weerden and Anderson 2013). Family wide comparison of *A. halleri* and *A. thaliana* *PDF1* genes revealed that the molecular function of the *A. thaliana* and *A. halleri* proteins in Zn tolerance and antifungal activity is conserved, and that functional differences in the two species may result from differential expression levels and regulation (Shahzad et al. 2013; Nguyen et al. 2014).

The fine mapping of a QTL in *A. halleri* allowed the identification of *CAX1* (cation/hydrogen exchanger 1) as a candidate gene for Cd tolerance (Courbot et al. 2007; Baliardini et al. 2015). *CAX1* is localized in the vacuolar membrane and plays a key role in Ca homeostasis (Conn et al. 2011). *CAX1* is more expressed in the roots of *A. halleri* compared to those of *A. thaliana*, and high expression of *CAX1* co-segregated with Cd tolerance in a back-cross 1 population of an *A. halleri*/*A. lyrata* cross. The *CAX1* QTL is conditional on Ca supply in the medium and is detected at low Ca supply only. The *A. thaliana cax1* mutant analysis suggests that, at low Ca supply, *CAX1* may be required to tolerate Cd-induced oxidative stress (Baliardini et al. 2015, 2016).

5 Mechanisms of Ni Hyperaccumulation

Today, more than 400 nickel (Ni) hyperaccumulator species have been identified worldwide. These species are scattered in more than 40 plant families, mostly dicotyledons (Krämer 2010; van der Ent et al. 2013; Cappa and Pilon-Smits 2014). Despite this large diversity and the interest in understanding the underlying mechanisms, as they may be instrumental to improve Ni phytoextraction, only a limited number of studies have focused on the molecular mechanisms of Ni hyperaccumulation. Ni is an essential microelement for plants because it is required for urease activity (Polacco et al. 2013). Plants have, therefore, evolved mechanisms for the regulation of Ni homeostasis and Ni hyperaccumulation that likely derive from these mechanisms (Fig. 3). In *A. thaliana*, the mechanisms involved in Ni homeostasis are strongly linked to Fe homeostasis (Schaaf

et al. 2006; Morrissey et al. 2009; Nishida et al. 2011). Interestingly, in Ni hyperaccumulators of the *Alyssum* genus, it was shown that Mn treatment reduces Ni accumulation, thus suggesting that in some species Ni hyperaccumulation may also use mechanisms primary involved in Mn homeostasis (Leigh Broadhurst et al. 2009; Ghaderian et al. 2015).

5.1 Uptake of Ni

Efficient uptake of Ni by the roots of hyperaccumulators requires divalent metal importers (e.g. ZIP, NRAMP) or transporters able to carry conjugated forms of Ni (e.g. Yellow Stripe-Like (YSL) family). However, the identity of the transporters involved in Ni uptake in hyperaccumulators is still not clearly established.

In *A. thaliana*, the metal transporter IRT1 that is required for the uptake of Fe from soil was shown to also be involved in Ni uptake (Vert et al. 2002; Nishida et al. 2011, 2012). Interestingly, the high expression of the *IRT1* ortholog in the roots of *N. caerulescens* (*NcIRT1*) is correlated with Ni hyperaccumulation in the Monte Prinzer accession (Halimaa et al. 2014b). In addition, *de novo* sequencing of *NcIRT1* in Monte Prinzer revealed sequence polymorphism in the large cytoplasmic loop of IRT1 that may have a role in transport specificity and/or regulation (Halimaa et al. 2014a). However, in other Ni hyperaccumulator accessions of *N. caerulescens* (i.e. Puy de Wolf and Bergenbach), we were unable to detect correlation between *NcIRT1* expression and Ni hyperaccumulation (V. S. Garcia de la Torre, S. Merlot, unpublished data). These data suggest that diverse transporters may be involved in the uptake of Ni in hyperaccumulators. Indeed, several members of the ZIP and NRAMP families from *Noccaea* species have been linked to Ni transport or accumulation, but further studies will be required to support their implication in an efficient uptake of Ni (Mizuno et al. 2005, 2007; Wei et al. 2009; Halimaa et al. 2014b).




Fig. 3 Proposed mechanisms of Ni transport in Hyperaccumulators. Efficient Ni uptake is mediated by metal transporters (ZIP/IRT, NRAMP...) located at the plasma membrane of root epidermal cells (ep). Ni is then transported through the cortex (co) and the endodermis (en) by a combination of Ni export and import transporter activities. During this step, Ni is chelated [e.g. by nicotianamine (Ni-NA)] to reduce its reactivity in the cytoplasm. Binding to His might prevent vacuolar sequestration

to favor radial transport. In the pericycle (pc), Ni is loaded in the xylem (xy) together with chelator molecules (e.g. NA, Citrate...) by YSL and MATE transporters and transported to the shoot. The mechanisms involved in xylem unloading and transport to the epidermal cell are not well known but may be similar as the one involved in Ni transport in roots. In epidermal cell, Ni is transported and stored in the vacuole (vac) by IREG transporters. Figure modified from Hanikenne and Nouet (2011)

5.2 Root-to-Shoot Transfer of Ni

The long-distance transport of Ni from roots to shoots requires several steps that involve metal transporters and chelators that are able to bind Ni in different pH environments. In hyperaccumulators, a large proportion of Ni is found as

complexes with carboxylic acids including citrate and malate (for reviews see Callahan et al. 2006; Sarret et al. 2013). These organic acid complexes are stable in acidic compartments such as vacuoles and xylem. In particular, citrate-Ni was identified in the xylem sap of the Ni hyperaccumulator *Alyssum serpyllifolium*

(Alves et al. 2011). Interestingly, an ortholog of the *A. thaliana* citrate transporter FRD3 of the Multidrug And Toxic compound Extrusion family (MATE) is more expressed in the hyperaccumulator *N. caerulescens* than in the related non-accumulator *A. thaliana* (van de Mortel et al. 2006). AtFRD3 and its orthologue in rice, OsFRDL1, are involved in the translocation of Fe from roots to shoots (Rogers and Gueriot 2002; Yokosho et al. 2009). Therefore, high expression of MATE transporters in the root pericycle of hyperaccumulators would increase the loading of xylem with citrate and therefore favor the translocation of Ni-citrate complex from root to shoot. However, to date no direct correlation has been established between MATE transporters and Ni hyperaccumulation. It is interesting to note that *FRD3* is also highly expressed in *A. halleri*, which hyperaccumulates Zn. FRD3 transporters may, therefore, have a general function in metal hyperaccumulation, favouring long-distance transport of metal from root to shoot (Talke et al. 2006; Charlier et al. 2015). NA also has a strong affinity for Ni over a wide pH range and is proposed to bind Ni in more neutral compartments such as cytoplasm or phloem (Callahan et al. 2006; Rellán-Alvarez et al. 2008; Alvarez-Fernandez et al. 2014). Accordingly, over-expression of NA synthase in transgenic *A. thaliana* increases Ni tolerance but is not sufficient to improve Ni accumulation (Pianelli et al. 2005). The NA-Ni complex was identified in the xylem sap of *N. caerulescens* (Mari et al. 2006), in the latex of the Ni hyperaccumulator *Pycnanthus acuminata* (Schaumörfel et al. 2003), and in extracts of several hyperaccumulator species (Callahan et al. 2012). Transporters of the YSL family have been shown to transport NA-metal complexes (Curie et al. 2009; Conte and Walker 2012). Several genes coding for YSL transporters are more expressed in the hyperaccumulator *N. caerulescens* than in the related non-accumulator *A. thaliana* (Gendre et al. 2007). Among these transporters, NcYSL3 that is able to transport the NA-Ni complex is expressed in the vasculature of roots and leaves, suggesting a role in long-distance Ni transport.

Finally, the amino acid histidine (His), whose concentration in some hyperaccumulators of the *Alyssum* and *Noccaea* genera correlates with Ni accumulation, is proposed to play a role in the radial transport of Ni (Krämer et al. 1996; Richau et al. 2009). His has a strong affinity for Ni and the Ni-His complex, and has been identified in samples from *Alyssum* and *Noccaea* hyperaccumulators (Krämer et al. 1996; Persans et al. 1999; Callahan et al. 2006; McNear et al. 2010). In the Ni hyperaccumulator *A. lesbiacum*, genes involved in His biosynthesis are constitutively and highly expressed compared to the non-accumulator *A. montanum*. The over-expression of the first enzyme of the His biosynthetic pathway, ATP-phosphoribosyltransferase, in *Arabidopsis thaliana* increases Ni tolerance but not Ni content, thus suggesting that other mechanisms are necessary for accumulation of this metal (Wycisk et al. 2004; Ingle et al. 2005). Treatment of plants with Ni-His increases xylem loading and inhibits Ni uptake from root vacuoles (Richau et al. 2009). It was therefore proposed that the high concentration of His in roots of Ni hyperaccumulators prevents vacuolar storage, favoring radial transport and xylem loading (Kerkeb and Krämer 2003; Richau et al. 2009). However, the mechanisms responsible for the inhibition of Ni vacuolar sequestration by His are still unknown. Moreover, it is currently not known if this strategy is widely conserved in Ni hyperaccumulators and if this is relevant in natural environments (e.g. ultramafic-derived soil), where N is limiting (Alves et al. 2011; Centofanti et al. 2013).

5.3 Storage of Ni in Leaves

In most hyperaccumulators that have been studied, Ni is stored in the vacuole of leaf epidermal cells (for review see Sarret et al. 2013). Several lines of evidence indicate that Ferroportin (FPN)/Iron Regulated (IREG) transporters play an essential role in the sequestration of Ni in vacuoles.

In *A. thaliana*, *AtIREG2* is expressed in roots in response to Fe-starvation and the *AtIREG2*

protein localization on the vacuole. The analysis of the *ireg2* mutant indicated that AtIREG2 is involved in the storage of Ni excess in the vacuole of root cells (Schaaf et al. 2006; Morrissey et al. 2009). On the contrary, a second IREG transporter in *A. thaliana*, named FPN1/IREG1, is localized on the plasma membrane and is proposed to play a role in xylem loading of metals in roots (Morrissey et al. 2009). Interestingly, the ortholog of AtIREG2 in *A. lyrata* is genetically linked to ultramafic adaptation (Turner et al. 2010). Recent comparative transcriptomic analysis using RNA-Seq technology revealed that a high expression of the ortholog of AtIREG2 in roots of *N. caerulea* is linked to Ni accumulation (Halimaa et al. 2014b). Independent RNA-Seq analyses also indicate that NcIREG2 is strongly and constitutively expressed in leaves of the *N. caerulea* accessions Puy de Wolf and Bergenbach (V. S. Garcia de la Torre, S. Merlot, unpublished results).

The PgIREG1 transporter from the Ni hyperaccumulator *Psychotria gabriellae* (Rubiaceae) localizes in the vacuolar membrane and is able to transport Ni when expressed in yeast. PgIREG1 therefore seems to be the functional homolog of AtIREG2. Interestingly, PgIREG1 is highly expressed in leaves of *P. gabriellae* in environmental conditions and is more expressed than in the closely related, non-accumulator *P. semperflorens* (Merlot et al. 2014). These data further suggest that the high expression of IREG transporters in leaves is a convergent mechanism for Ni hyperaccumulation. However, whereas over-expression of AtIREG2 and PgIREG1 in transgenic *Arabidopsis* plants significantly increases Ni tolerance, it does not increase Ni accumulation, further supporting the hypothesis that other mechanisms are required for hyperaccumulation (Schaaf et al. 2006; Merlot et al. 2014). Other families of divalent metal exporters such as MTP transporters could mediate the transport of Ni in vacuoles, but their role in hyperaccumulation needs to be further supported (Persans et al. 2001).

6 Hyperaccumulation of Other Trace Elements

Species that are able to hyperaccumulate Mn have been identified in more than 10 genera mostly in the Myrtaceae (e.g. *Gossia*) and Proteaceae (e.g. *Virotia*) families (Fernando et al. 2013; Losfeld et al. 2015). As for other metals, Mn hyperaccumulation likely evolved from basic mechanisms involved in Mn homeostasis; however, these mechanisms are poorly investigated to date in Mn hyperaccumulators (Pittman 2005; Fernando et al. 2013; Socha and Guerinet 2014). In hyperaccumulators, Mn was found to accumulate in the vacuole of non-photosynthetic epidermal cells, but also more surprisingly in photosynthetic palisade mesophyll cells in the hyperaccumulator *Virotia neurophylla* (Fernando et al. 2012). These differences in Mn localization suggest that some mechanisms involved in Mn accumulation and detoxification might be divergent among hyperaccumulators. The high concentration of Mn found in leaves of several species is linked to their strategy to acquire P. For example, Proteaceae species excrete carboxylates in their rhizosphere that solubilize not only P but also micronutrients including Mn (Lambers et al. 2015). Several families of metal transporters, such as NRAMP, ZIP, YSL, and MTP have been shown to be able to transport Mn in plants, but their role in hyperaccumulation is not clearly established (Fernando et al. 2013; Socha and Guerinet 2014). Most NRAMP transporters are able to transport Mn in the cytoplasm, either from the external medium or the vacuole, and therefore could participate in several steps of Mn hyperaccumulation. In particular, the NRAMP1 transporter from *A. thaliana* was shown to be the main transporter involved in Mn uptake in roots (Cailliatte et al. 2010). The ShMTP8 transporter (previously known as ShMTP1) was isolated from the Mn-tolerant species *Stylosanthes hamata* (Fabaceae) as a transporter that confers Mn resistance when expressed in yeast, and was proposed to mediate accumulation of Mn in the vacuoles of plant cells (Delhaize et al. 2003). Further molecular studies on Mn hyperaccumulators will be

required to identify those mechanisms that are key to Mn hyperaccumulation.

In addition to the above-mentioned metals, mechanisms involved in the hyperaccumulation of the metalloid element Se in plants are extensively studied because of their relevance for improving Se phytoremediation and biofortification. We only briefly address Se hyperaccumulation here and refer interested readers to thorough reviews on Se homeostasis and hyperaccumulation (Zhu et al. 2009; Barillas et al. 2011; Schiavon and Pilon-Smits 2016; White 2016). Selenium hyperaccumulation has been described for 45 taxa scattered among six families. More than half of Se hyperaccumulators have been described in the genus *Astragalus* (Fabaceae), but other well-described Se hyperaccumulators have been found in the genera *Stanleya* (Brassicaceae), *Oenopsis*, and *Xylorhiza* (Asteraceae). The distribution of Se hyperaccumulators among plant families suggests that Se hyperaccumulation likely evolved independently at least six times (Cappa and Pilon-Smits 2014). Selenium is available to plants mostly as selenate (SeO_4^{2-}), a structural homologue of sulphate, or selenite (SeO_3^{2-}), depending on the nature of the soil (Elrashidi et al. 1987). In cultivated soils, selenate uptake by root cells is catalyzed by high-affinity sulphate transporters of the SULTR family (Shibagaki et al. 2002; El Kassis et al. 2007; Barberon et al. 2008). Transporters of this family are highly and constitutively expressed in Se hyperaccumulators of the *Astragalus* and *Stanleya* genera (Freeman et al. 2010; Cabannes et al. 2011; Schiavon et al. 2015). The Se hyperaccumulation trait is also linked to the capacity of Se hyperaccumulators to preferentially take up selenate over sulphate. Sequence analysis of SULTR1 transporters from *Astragalus* identified a Gly to Ala polymorphism linked to the hyperaccumulation trait; however, it is not demonstrated yet if this difference explains the preferential uptake of selenate over sulphate (Cabannes et al. 2011). In rice growing in anaerobic soils (e.g. paddy fields), selenite forms are transported in root cells by the phosphate transporters OsPT2 (Zhang et al. 2014) and

by aquaporins (Zhao et al. 2010). Long-distance transport of selenate to the shoot is also proposed to be mediated by SULTR transporters (Takahashi et al. 2000). In shoots, the main fraction of selenate is metabolized into organoselenium compounds (SeCys and SeMet) in chloroplast (Zhu et al. 2009; Barillas et al. 2011; White 2016). In hyperaccumulator species, genes involved in the synthesis of organoselenium compounds and in the methylation of SeCys to produce the non-toxic form MeSeCys were shown to be constitutively more expressed than in non-accumulator species (Pickering et al. 2003; Freeman et al. 2010; Schiavon et al. 2015), indicating that the metabolism of Se is key for its hyperaccumulation.

7 Perspectives and Conclusions

7.1 Interaction of Hyperaccumulators with Biotic Environment

The so-called ‘elemental defense’ hypothesis proposes that metal hyperaccumulation provides a defense against pathogens and/or herbivores by direct toxicity (Boyd and Martens 1992; Hörger et al. 2013). Indeed, it has been shown that Ni and Zn accumulation can protect different Brassicaceae species from bacterial and fungal infection (Boyd et al. 1994; Ghaderian et al. 2000; Fones et al. 2010). However, it is also well documented that in environmental conditions, specific populations of bacteria are associated with the root system of metal hyperaccumulators (Aboudrar et al. 2012; Cabello-Conejo et al. 2014; Lucisine et al. 2014; Muehe et al. 2015). Endophytic bacteria have also been identified in the shoots and roots of metal hyperaccumulators (Idris et al. 2004; Mengoni et al. 2009a; Nonnoi et al. 2012). However, since the majority of these bacteria are not cultivable, current studies mostly describe endophytic populations using metagenomics approaches (Chen et al. 2014; Luo et al. 2011; Sessitsch et al. 2012; Visioli et al. 2014). Interestingly, it was shown that the inoculation of

N. caerulescens and *A. serpyllifolium* with culturable endophytic bacteria increases Ni translocation to shoots (Ma et al. 2011b; Visioli et al. 2015). Little is known about the interactions between metal hyperaccumulators and associated bacteria. Metal hyperaccumulators represent an extreme niche for metal-tolerant bacteria (Mengoni et al. 2009b). On the other hand, metal-tolerant bacteria can improve plant growth and confer protection against abiotic stress by the production of hormones (auxins, cytokinins, etc.), or protect the host plant against other pathogens by the production of antagonistic substances or by competition for space and nutrients (Ma et al. 2011b; Reinhold-Hurek and Hurek 2011). Metal-tolerant bacteria can also produce organic acids and metal chelators that can favor metal solubility, transport, and tolerance (Idris et al. 2006; Ma et al. 2011a; Visioli et al. 2015). These observations suggest intimate interactions between hyperaccumulators and associated bacteria; however, mechanisms and genes involved in these interactions are mostly unknown. The development of next-generation sequencing technologies and associated meta-transcriptomic (or dual-transcriptomics) analyses will uncover those genes expressed by plants and associated bacteria involved in their interaction (Camilios-Neto et al. 2014; Pankiewicz et al. 2016). Understanding these mechanisms will be instrumental to improving metal phytoextraction or to producing secondary metabolites such as metal chelators of bacterial origin that can be used in metal-based therapies (Franz 2013).

7.2 Evolution of Hyperaccumulation Mechanisms

After this brief review of our current knowledge of the molecular mechanisms of Zn, Cd, or Ni hyperaccumulation, it is apparent that several candidate genes involved in these processes are involved in the control of metal homeostasis in non-accumulator plants. These genes display an enhanced function in hyperaccumulators, through gene-copy number amplification and/or altered regulation, which profoundly modifies

the metal flux in the plants towards shoot accumulation. Many examples have also been presented that highlight the high level of convergent evolution between *A. halleri*, *N. caerulescens*, and *S. alfredii*. This convergent evolution likely reflects—and sheds light on—functional constraints of the metal homeostasis network (Krämer et al. 2007; Krämer 2010; Verbruggen et al. 2009a; Hanikenne and Nouet 2011). The key function of HMA4 in several hyperaccumulator species represents a potent example of this convergent evolution (Hanikenne et al. 2008; O’Lochlainn et al. 2011; Craciun et al. 2012).

Metal hyperaccumulation and associated tolerance in plants are complex traits that have required the fine-tuning of multiple mechanisms during the course of evolution. If several key players have now been identified, how tolerance and hyperaccumulation traits evolved remains an open ‘chicken or egg’ question. Hence, Bayesian inference suggested that speciation between *A. halleri* and *A. lyrata* closely coincided with *HMA4* duplication (Roux et al. 2011). The complex signature of selection detected at the *HMA4* locus of *A. halleri* further supports the key role of this gene in the evolution of the hyperaccumulation trait (Hanikenne et al. 2013). Recent adaptations to anthropogenic metal-polluted sites possibly occurred independently within distinct phylogeographic units of the *A. halleri* European distribution (Pauwels et al. 2012). Hypertolerance of metallicolous populations thus potentially evolved using a variety of genetic mechanisms (Meyer et al. 2009, 2010; Pauwels et al. 2012). Moreover, the recent study by Meyer et al. (2016) suggests that, if HMA4 contributes to Zn tolerance in both metallicolous and non-metallicolous populations of *A. halleri* (Hanikenne et al. 2013), the function of MTP1 in Zn tolerance may have evolved later in metallicolous populations that colonized recently polluted soils. Indeed, the co-segregation of *MTP1* with Zn tolerance is only observed in a back-cross 1 population of a cross between a French metallicolous individual (i.e. living on metal-polluted soil) of *A. halleri* and *A. lyrata*, and is lost when a Slovakian non-metallicolous

(i.e. living on non-polluted soil) *A. halleri* individual is used as parent (Meyer et al. 2016). In contrast, the co-segregation of *HMA4* with Zn tolerance is independent of the edaphic origin of the *A. halleri* populations used in the analysis (Willems et al. 2007; Meyer et al. 2016). This interpretation partially contradicts the hypothesis that MTP1 is required for metal detoxification accommodating the high HMA4-dependent metal flux into *A. halleri* shoots, which was proposed based on the observation that expression of *AhHMA4* in non-accumulator plants resulted in increased sensitivity to excess Zn (Hanikenne et al. 2008; Barabasz et al. 2010).

The study of metal hyperaccumulation in distant plant families combined with the comparison of distinct accessions with contrasting accumulation capabilities within a species will likely shed light on the evolution of the hyperaccumulation and hypertolerance traits. Such study may indeed reveal the commonalities and differences in the mechanisms underlying these traits, highlighting evolutionary divergence and convergence. It also may reveal evolutionary ‘intermediates’ (i.e. genotypes that do not display the full extent of hyperaccumulation or tolerance), which may allow ordering the evolutionary events that took place during the adaptation of the metal homeostasis network.

7.3 How Can Phytoextraction Technologies Benefit from Molecular Knowledge?

To date, most of our knowledge on metal hyperaccumulation arises from studies on a few model hyperaccumulator species (i.e. *A. halleri* and *N. caerulescens*) of the Brassicaceae family. In the future, it will be necessary to pursue molecular studies and improve genetic manipulation of these species, in order to identify and demonstrate the role of key mechanisms involved in metal hyperaccumulation. However, these species have a low biomass and a relatively restricted distribution worldwide. Therefore, one of the coming challenges for the develop-

ment of agromining/phytoextraction will be to transfer knowledge of the mechanisms involved in metal hyperaccumulation to species having a high potential for phytoextraction. As mentioned above, the development of Next Generation Sequencing technologies opens the possibility for study of ‘non-model’ species at the genomic and transcriptomic levels.

As for other crop plants, this molecular knowledge will be instrumental for the selection of genotypes having the best potential for metal phytoextraction. The level of expression of key genes involved in metal hyperaccumulation can be used as markers to predict metal accumulation capacities. These marker genes can also be used to study the interaction between metal accumulation and agricultural practices (e.g. fertilization), for improving biomass production while maintaining efficient metal accumulation.

Finally, genome-editing technologies such as CRISPR-CAS9 are currently implemented in plants to specifically modify the sequence of target genes (Doudna and Charpentier 2014). This technology offers several advantages compared to traditional transformation technologies used to produce Genetically Modified Organisms (GMO), and therefore could be better accepted by the civil society and political stakeholders to engineer crop plants for metal phytoextraction. Using this recent technology, we can, for example, imagine specifically introducing point mutations in the sequence of genes involved in metal accumulation in order to increase their activity, improve specificity, or modify selectivity towards metals of interest (Rogers et al. 2000; Menguer et al. 2013; Pottier et al. 2015).

Acknowledgments We thank members of our laboratories for critical reading of the manuscript. The research of SM and VSGT is supported by the French National Research Agency (ANR-13-ADAP-0004) and by CNRS (Defi Environnements Gene-4-Chem). Funding to MH is from the “Fonds de la Recherche Scientifique-FNRS” (FRFC-2.4583.08, PDR-T.0206.13, MIS-F.4511.16, CDR J.0009.17), the University of Liège (SFRD-12/03) and the Belgian Program on Interuniversity Poles of Attraction (IAP no. P6/19). MH is a Research Associate of the FNRS.

References

- Aboudrar W, Schwartz C, Morel JL, Boularbah A (2012) Effect of nickel-resistant rhizosphere bacteria on the uptake of nickel by the hyperaccumulator *Nocca caerulea* under controlled conditions. *J Soils Sediments* 13:501–507
- Alvarez-Fernandez A, Diaz-Benito P, Abadia A, Lopez-Millan AF, Abadia J (2014) Metal species involved in long distance metal transport in plants. *Front Plant Sci* 5:105. doi:10.3389/fpls.2014.00105
- Alves S, Nabais C, Simoes Goncalves Mde L, Correia Dos Santos MM (2011) Nickel speciation in the xylem sap of the hyperaccumulator *Alyssum serpyllifolium* ssp. *lusitanicum* growing on serpentine soils of north-east Portugal. *J Plant Physiol* 168:1715–1722
- Assunção AGL, Bookum WM, Nelissen HJM, Vooijs R, Schat H, Ernst WHO (2003) Differential metal-specific tolerance and accumulation patterns among *Thlaspi caerulescens* populations originating from different soil types. *New Phytol* 159:411–419
- Assunção AGL, Herrero E, Lin YF, Huettel B, Talukdar S, Smaczniak C, Immink RG, van Eldik M, Fiers M, Schat H, Aarts MG (2010) *Arabidopsis thaliana* transcription factors bZIP19 and bZIP23 regulate the adaptation to zinc deficiency. *Proc Natl Acad Sci USA* 107:10296–10301
- Baliardini C, Meyer C-L, Salis P, Saumitou-Laprade P, Verbruggen N (2015) CAX1 co-segregates with Cd tolerance in the metal hyperaccumulator *Arabidopsis halleri* and plays a role in limiting oxidative stress in *Arabidopsis* spp. *Plant Physiol* 169:549–559
- Baliardini C, Corso M, Verbruggen N (2016) Transcriptomic analysis supports the role of CATION EXCHANGER 1 in cellular homeostasis and oxidative stress limitation during cadmium stress. *Plant Signal Behav* 11:e1183861
- Barabasz A, Krämer U, Hanikenne M, Rudzka J, Antosiewicz DM (2010) Metal accumulation in tobacco expressing *Arabidopsis halleri* metal hyperaccumulation gene depends on external supply. *J Exp Bot* 61:3057–3067
- Barberon M, Berthomieu P, Clairotte M, Shibagaki N, Davidian JC, Gosti F (2008) Unequal functional redundancy between the two *Arabidopsis thaliana* high-affinity sulphate transporters SULTR1;1 and SULTR1;2. *New Phytol* 180:608–619
- Barillas JRV, Quinn CF, Pilon-Smits EAH (2011) Selenium accumulation in plants—phytotechnological applications and ecological implications. *Int J Phytoremediation* 13:166–178
- Becher M, Talke IN, Krall L, Krämer U (2004) Cross-species microarray transcript profiling reveals high constitutive expression of metal homeostasis genes in shoots of the zinc hyperaccumulator *Arabidopsis halleri*. *Plant J* 37:251–268
- Bernard C, Roosens N, Czernic P, Lebrun M, Verbruggen N (2004) A novel CPx-ATPase from the cadmium hyperaccumulator *Thlaspi caerulescens*. *FEBS Lett* 569:140–148
- Bert V, MacNair MR, De Laguérie P, Saumitou-Laprade P, Petit D (2000) Zinc tolerance and accumulation in metalicolous and non metalicolous populations of *Arabidopsis halleri* (Brassicaceae). *New Phytol* 146:225–233
- Bert V, Bonnin I, Saumitou-Laprade P, De Laguérie P, Petit D (2002) Do *Arabidopsis halleri* from non metalicolous populations accumulate zinc and cadmium more effectively than those from metalicolous populations? *New Phytol* 155:47–57
- Besnard G, Basic N, Christin PA, Savova-Bianchi D, Galland N (2009) *Thlaspi caerulescens* (Brassicaceae) population genetics in western Switzerland: is the genetic structure affected by natural variation of soil heavy metal concentrations? *New Phytol* 181:974–984
- Boyd R, Martens SN (1992) The vegetation of ultramafic (serpentine) soils. Intercept, Andover, Hampshire
- Boyd RS, Shaw JJ, Martens SN (1994) Nickel hyperaccumulation defends *Streptanthus polygaloides* (Brassicaceae) against pathogens. *Am J Bot* 81:294–300
- Burkhead JL, Reynolds KA, Abdel-Ghany SE, Cohu CM, Pilon M (2009) Copper homeostasis. *New Phytol* 182:799–816
- Cabannes E, Buchner P, Broadley MR, Hawkesford MJ (2011) A comparison of sulfate and selenium accumulation in relation to the expression of sulfate transporter genes in *Astragalus* species. *Plant Physiol* 157:2227–2239
- Cabello-Conejo MI, Becerra-Castro C, Prieto-Fernández A, Monterroso C, Saavedra-Ferro A, Mench M, Kidd PS (2014) Rhizobacterial inoculants can improve nickel phytoextraction by the hyperaccumulator *Alyssum pintodasilvae*. *Plant Soil* 379:35–50
- Cailliatte R, Schikora A, Briat JF, Mari S, Curie C (2010) High-affinity manganese uptake by the metal transporter NRAMP1 is essential for *Arabidopsis* growth in low manganese conditions. *Plant Cell* 22:904–917
- Callahan DL, Baker AJM, Kolev SD, Wedd AG (2006) Metal ion ligands in hyperaccumulating plants. *J Biol Inorg Chem* 11:2–12
- Callahan DL, Kolev SD, O'Hair RA, Salt DE, Baker AJM (2007) Relationships of nicotianamine and other amino acids with nickel, zinc and iron in *Thlaspi* hyperaccumulators. *New Phytol* 176:836–848
- Callahan DL, Roessner U, Dumontet V, De Livera AM, Doronila A, Baker AJM, Kolev SD (2012) Elemental and metabolite profiling of nickel hyperaccumulators from New Caledonia. *Phytochemistry* 81:80–89
- Callahan DL, Hare DJ, Bishop DP, Doble PA, Roessner U (2016) Elemental imaging of leaves from the metal hyperaccumulating plant *Nocca caerulea* shows different spatial distribution of Ni, Zn and Cd. *RSC Adv* 6:2337–2344

- Camilios-Neto D, Bonato P, Wassem R, Tadra-Sfeir MZ, Brusamarello-Santos LC, Valdameri G, Donatti L, Faoro H, Weiss VA, Chubatsu LS, Pedrosa FO, Souza EM (2014) Dual RNA-seq transcriptional analysis of wheat roots colonized by *Azospirillum brasilense* reveals up-regulation of nutrient acquisition and cell cycle genes. *BMC Genomics* 15:378
- Cao D, Zhang H, Wang Y, Zheng L (2014) Accumulation and distribution characteristics of zinc and cadmium in the hyperaccumulator plant *Sedum plumbizincicola*. *Bull Environ Contam Toxicol* 93:171–176
- Cappa JJ, Pilon-Smits EAH (2014) Evolutionary aspects of elemental hyperaccumulation. *Planta* 239:267–275
- Centofanti T, Sayers Z, Cabello-Conejo MI, Kidd P, Nishizawa NK, Kakei Y, Davis AP, Sicher RC, Chaney RL (2013) Xylem exudate composition and root-to-shoot nickel translocation in *Alyssum* species. *Plant Soil* 373:59–75
- Charlier JB, Polese C, Nouet C, Carnol M, Bosman B, Krämer U, Motte P, Hanikenne M (2015) Zinc triggers a complex transcriptional and post-transcriptional regulation of the metal homeostasis gene *FRD3* in *Arabidopsis* relatives. *J Exp Bot* 66:3865–3878
- Chen L, Luo S, Chen J, Wan Y, Li X, Liu C, Liu F (2014) A comparative analysis of endophytic bacterial communities associated with hyperaccumulators growing in mine soils. *Environ Sci Pollut Res Int* 21:7538–7547
- Chiang HC, Lo JC, Yeh KC (2006) Genes associated with heavy metal tolerance and accumulation in Zn/Cd hyperaccumulator *Arabidopsis halleri*: a genomic survey with cDNA microarray. *Environ Sci Technol* 40:6792–6798
- Claus J, Bohmann A, Chavarría-Krauser A (2013) Zinc uptake and radial transport in roots of *Arabidopsis thaliana*: a modelling approach to understand accumulation. *Ann Bot* 112:369–380
- Clauss MJ, Koch MA (2006) Poorly known relatives of *Arabidopsis thaliana*. *Trends Plant Sci* 11:449–459
- Clemens S, Palmgren MG, Krämer U (2002) A long way ahead: understanding and engineering plant metal accumulation. *Trends Plant Sci* 7:309–315
- Clemens S, Deinlein U, Ahmadi H, Höreth S, Uruguchi S (2013) Nicotianamine is a major player in plant Zn homeostasis. *Biomaterials* 26:623–632
- Conn SJ, Gilliam M, Athman A, Schreiber AW, Baumann U, Moller I, Cheng N-H, Stancombe MA, Hirschi KD, Webb AAR, Burton R, Kaiser BN, Tyerman SD, Leigh RA (2011) Cell-specific vacuolar calcium storage mediated by CAX1 regulates apoplastic calcium concentration, gas exchange, and plant productivity in *Arabidopsis*. *Plant Cell* 23:240–257
- Conte SS, Walker EL (2012) Genetic and biochemical approaches for studying the yellow stripe-like transporter family in plants. *Curr Top Membr* 69:295–322
- Cornu J, Deinlein U, Höreth S, Braun M, Schmidt H, Weber M, Persson DP, Husted S, Schjoerring JK, Clemens S (2015) Contrasting effects of nicotianamine synthase knockdown on zinc and nickel tolerance and accumulation in the zinc/cadmium hyperaccumulator *Arabidopsis halleri*. *New Phytol* 206:738–750
- Cosio C, DeSantis L, Frey B, Diallo S, Keller C (2005) Distribution of cadmium in leaves of *Thlaspi caerulescens*. *J Exp Bot* 56:765–775
- Courbot M, Willems G, Motte P, Arvidsson S, Roosens N, Saumitou-Laprade P, Verbruggen N (2007) A major QTL for Cd tolerance in *Arabidopsis halleri* co-localizes with *HMA4*, a gene encoding a heavy metal ATPase. *Plant Physiol* 144:1052–1065
- Craciun AR, Courbot M, Bourgis F, Salis P, Saumitou-Laprade P, Verbruggen N (2006) Comparative cDNA-AFLP analysis of Cd-tolerant and -sensitive genotypes derived from crosses between the Cd hyperaccumulator *Arabidopsis halleri* and *Arabidopsis lyrata* ssp. *petraea*. *J Exp Bot* 57:2967–2983
- Craciun AR, Meyer C-L, Chen J, Roosens N, De Groodt R, Hilson P, Verbruggen N (2012) Variation in *HMA4* gene copy number and expression among *Noccaea caerulescens* populations presenting different levels of Cd tolerance and accumulation. *J Exp Bot* 63:4179–4189
- Curie C, Cassin G, Couch D, Divol F, Higuchi K, Le Jean M, Misson J, Schikora A, Czernic P, Mari S (2009) Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters. *Ann Bot* 103:1–11
- De Coninck B, Cammue BPA, Thevissen K (2013) Modes of antifungal action and in planta functions of plant defensins and defensin-like peptides. *Fungal Biol Rev* 26:109–120
- Deinlein U, Weber M, Schmidt H, Rensch S, Trampczynska A, Hansen TH, Husted S, Schjoerring JK, Talke IN, Krämer U, Clemens S (2012) Elevated nicotianamine levels in *Arabidopsis halleri* roots play a key role in zinc hyperaccumulation. *Plant Cell* 24:708–723
- Delhaize E, Kataoka T, Hebb DM, White RG, Ryan PR (2003) Genes encoding proteins of the cation diffusion facilitator family that confer manganese tolerance. *Plant Cell* 15:1131–1142
- Deng DM, Shu WS, Zhang J, Zou HL, Lin Z, Ye ZH, Wong MH (2007) Zinc and cadmium accumulation and tolerance in populations of *Sedum alfredii*. *Environ Pollut* 147:381–386
- Deniau AX, Pieper B, Ten Bookum WM, Lindhout P, Aarts MG, Schat H (2006) QTL analysis of cadmium and zinc accumulation in the heavy metal hyperaccumulator *Thlaspi caerulescens*. *Theor Appl Genet* 113:907–920
- Doudna JA, Charpentier E (2014) Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science* 346:1258096
- Dräger DB, Desbrosses-Fonrouge AG, Krach C, Chardonnens AN, Meyer RC, Saumitou-Laprade P, Krämer U (2004) Two genes encoding *Arabidopsis halleri* MTP1 metal transport proteins co-segregate

- with zinc tolerance and account for high MTP1 transcript levels. *Plant J* 39:425–439
- El Kassis E, Cathala N, Rouached H, Fourcroy P, Berthomieu P, Terry N, Davidian JC (2007) Characterization of a selenate-resistant *Arabidopsis* mutant. Root growth as a potential target for selenate toxicity. *Plant Physiol* 143:1231–1241
- Elrashidi MA, Adriano DC, Workman SM, Lindsay WL (1987) Chemical-equilibria of selenium in soils—a theoretical development. *Soil Sci* 144:141–152
- Escarré J, Lefèbvre C, Gruber W, Leblanc M, Lepart J, Rivière Y, Delay B (2000) Zinc and cadmium hyperaccumulation by *Thlaspi caerulescens* from metalliferous and nonmetalliferous sites in the Mediterranean area: implications for phytoremediation. *New Phytol* 145:429–437
- Escarré J, Lefèbvre C, Frérot H, Mahieu S, Noret N (2013) Metal concentration and metal mass of metallicolous, non metallicolous and serpentine *Noccaea caerulescens* populations, cultivated in different growth media. *Plant Soil* 370:197–221
- Fernando DR, Bakkaus EJ, Perrier N, Baker AJ, Woodrow IE, Batianoff GN, Collins RN (2006a) Manganese accumulation in the leaf mesophyll of four tree species: a PIXE/EDAX localization study. *New Phytol* 171:751–757
- Fernando DR, Batianoff GN, Baker AJM, Woodrow IE (2006b) In vivo localization of manganese in the hyperaccumulator *Gossia bidwillii* (Benth.) N. Snow & Guymer (Myrtaceae) by cryo-SEM/EDAX. *Plant Cell Environ* 29:1012–1020
- Fernando DR, Woodrow IE, Baker AJM, Marshall AT (2012) Plant homeostasis of foliar manganese sinks: specific variation in hyperaccumulators. *Planta* 236:1459–1470
- Fernando DR, Marshall A, Baker AJM, Mizuno T (2013) Microbeam methodologies as powerful tools in manganese hyperaccumulation research: present status and future directions. *Front Plant Sci* 4:319
- Filatov V, Dowdle J, Smirnov N, Ford-Lloyd B, Newbury HJ, Macnair MR (2006) Comparison of gene expression in segregating families identifies genes and genomic regions involved in a novel adaptation, zinc hyperaccumulation. *Mol Ecol* 15:3045–3059
- Filatov V, Dowdle J, Smirnov N, Ford-Lloyd B, Newbury HJ, Macnair MR (2007) A quantitative trait loci analysis of zinc hyperaccumulation in *Arabidopsis halleri*. *New Phytol* 174:580–590
- Fones H, Davis CAR, Rico A, Fang F, Smith JAC, Preston GM (2010) Metal hyperaccumulation armors plants against disease. *PLoS Pathog* 6:e1001093
- Franz KJ (2013) Clawing back: broadening the notion of metal chelators in medicine. *Curr Opin Chem Biol* 17:143–149
- Freeman JL, Tamaoki M, Stushnoff C, Quinn CF, Cappa JJ, Devonshire J, Fakra SC, Marcus MA, McGrath SP, Van Hoewyk D, Pilon-Smits EA (2010) Molecular mechanisms of selenium tolerance and hyperaccumulation in *Stanleya pinnata*. *Plant Physiol* 153:1630–1652
- Frérot H, Faucon MP, Willems G, Gode C, Courseaux A, Darracq A, Verbruggen N, Saumitou-Laprade P (2010) Genetic architecture of zinc hyperaccumulation in *Arabidopsis halleri*: the essential role of QTL x environment interactions. *New Phytol* 187:355–367
- Gao J, Sun L, Yang X, Liu J-X (2013) Transcriptomic analysis of cadmium stress response in the heavy metal hyperaccumulator *Sedum alfredii* Hance. *PLoS One* 8:e64643
- Gendre D, Czernic P, Conejero G, Pianelli K, Briat JF, Lebrun M, Mari S (2007) TcYSL3, a member of the YSL gene family from the hyper-accumulator *Thlaspi caerulescens*, encodes a nicotianamine-Ni/Fe transporter. *Plant J* 49:1–15
- Ghaderian YSM, Lyon AJE, Baker AJM (2000) Seedling mortality of metal hyperaccumulator plants resulting from damping off by *Pythium spp.* *New Phytol* 146:219–224
- Ghaderian SM, Ghasemi R, Hajhashemi F (2015) Interaction of nickel and manganese in uptake, translocation and accumulation by the nickel-hyperaccumulator plant, *Alyssum bracteatum* (Brassicaceae). *Aust J Bot* 63:47–55
- Gonneau C, Genevois N, Frérot H, Sirguey C, Sterckeman T (2014) Variation of trace metal accumulation, major nutrient uptake and growth parameters and their correlations in 22 populations of *Noccaea caerulescens*. *Plant Soil* 384:271–287
- Guimarães MD, Gustin JL, Salt DE (2009) Reciprocal grafting separates the roles of the root and shoot in zinc hyperaccumulation in *Thlaspi caerulescens*. *New Phytol* 184:323–329
- Gustin JL, Loureiro ME, Kim D, Na G, Tikhonova M, Salt DE (2009) MTP1-dependent Zn sequestration into shoot vacuoles suggests dual roles in Zn tolerance and accumulation in Zn hyperaccumulating plants. *Plant J* 57:1116–1127
- Halimaa P, Blande D, Aarts MG, Tuomainen M, Tervahauta A, Karenlampi S (2014a) Comparative transcriptome analysis of the metal hyperaccumulator *Noccaea caerulescens*. *Front Plant Sci* 5:213. doi:10.3389/fpls.2014.00213
- Halimaa P, Lin YF, Ahonen VH, Blande D, Clemens S, Gyenesei A, Haikio E, Karenlampi SO, Laiho A, Aarts MG, Pursiheimo JP, Schat H, Schmidt H, Tuomainen MH, Tervahauta AI (2014b) Gene expression differences between *Noccaea caerulescens* ecotypes help to identify candidate genes for metal phytoremediation. *Environ Sci Technol* 48:3344–3353
- Hammond JP, Bowen H, White PJ, Mills V, Pyke KA, Baker AJM, Whiting SN, May ST, Broadley MR (2006) A comparison of *Thlaspi caerulescens* and *Thlaspi arvense* shoot transcriptomes. *New Phytol* 170:239–260
- Han X, Yin H, Song X, Zhang Y, Liu M, Sang J, Jiang J, Li J, Zhuo R (2015) Integration of small RNAs, degradome and transcriptome sequencing in

- hyperaccumulator *Sedum alfredii* uncovers a complex regulatory network and provides insights into cadmium phytoremediation. *Plant Biotechnol J* 14: 1470–1483
- Hanikenne M, Baurain D (2014) Origin and evolution of metal p-type ATPases in Plantae (Archaeplastida). *Front Plant Sci* 4:544
- Hanikenne M, Nouet C (2011) Metal hyperaccumulation and hypertolerance: a model for plant evolutionary genomics. *Curr Opin Plant Biol* 14:252–259
- Hanikenne M, Talke IN, Haydon MJ, Lanz C, Nolte A, Motte P, Kroymann J, Weigel D, Krämer U (2008) Evolution of metal hyperaccumulation required *cis*-regulatory changes and triplication of *HMA4*. *Nature* 453:391–395
- Hanikenne M, Kroymann J, Trampczynska A, Bernal M, Motte P, Clemens S, Krämer U (2013) Hard selective sweep and ectopic gene conversion in a gene cluster affording environmental adaptation. *PLoS Genet* 9: e1003707
- Hörger AC, Fones HN, Preston GM (2013) The current status of the elemental defense hypothesis in relation to pathogens. *Front Plant Sci* 4:395. doi:10.3389/fpls.2013.00395
- Hussain D, Haydon MJ, Wang Y, Wong E, Sherson SM, Young J, Camakaris J, Harper JF, Cobbett CS (2004) P-type ATPase heavy metal transporters with roles in essential zinc homeostasis in *Arabidopsis*. *Plant Cell* 16:1327–1339
- Idris R, Trifonova R, Puschenreiter M, Wenzel WW, Sessitsch A (2004) Bacterial communities associated with flowering plants of the Ni hyperaccumulator *Thlaspi goesingense*. *Appl Environ Microbiol* 70: 2667–2677
- Idris R, Kuffner M, Bodrossy L, Puschenreiter M, Monchy S, Wenzel WW, Sessitsch A (2006) Characterization of Ni-tolerant methylobacteria associated with the hyperaccumulating plant *Thlaspi goesingense* and description of *Methylobacterium goesingense* sp. nov. *Syst Appl Microbiol* 29:634–644
- Ingle RA, Mugford ST, Rees JD, Campbell MM, Smith JAC (2005) Constitutively high expression of the histidine biosynthetic pathway contributes to nickel tolerance in hyperaccumulator plants. *Plant Cell* 17: 2089–2106
- Isaure MP, Huguet S, Meyer CL, Castillo-Michel H, Testemale D, Vantelon D, Saumitou-Laprade P, Verbruggen N, Sarret G (2015) Evidence of various mechanisms of Cd sequestration in the hyperaccumulator *Arabidopsis halleri*, the non-accumulator *Arabidopsis lyrata*, and their progenies by combined synchrotron-based techniques. *J Exp Bot* 66:3201–3214
- Kerkeb L, Krämer U (2003) The role of free histidine in xylem loading of nickel in *Alyssum lesbiacum* and *Brassica juncea*. *Plant Physiol* 131:716–724
- Kim D, Gustin JL, Lahner B, Persans MW, Baek D, Yun DJ, Salt DE (2004) The plant CDF family member TgMTP1 from the Ni/Zn hyperaccumulator *Thlaspi goesingense* acts to enhance efflux of Zn at the plasma membrane when expressed in *Saccharomyces cerevisiae*. *Plant J* 39:237–251
- Kozhevnikova AD, Seregin IV, Erlikh NT, Shevyreva TA, Andreev IM, Verweij R, Schat H (2014) Histidine-mediated xylem loading of zinc is a species-wide character in *Noccaea caerulea*. *New Phytol* 203:508–519
- Krämer U (2005) MTP1 mops up excess zinc in *Arabidopsis* cells. *Trends Plant Sci* 10:313–315
- Krämer U (2010) Metal hyperaccumulation in plants. *Annu Rev Plant Biol* 61:517–534
- Krämer U, Cotter-Howells JD, Charnock JM, Baker AJM, Smith JAC (1996) Free histidine as a metal chelator in plants that accumulate nickel. *Nature* 379:635–638
- Krämer U, Talke IN, Hanikenne M (2007) Transition metal transport. *FEBS Lett* 581:2263–2272
- Küpper H, Zhao FJ, McGrath SP (1999) Cellular compartmentation of zinc in leaves of the hyperaccumulator *Thlaspi caerulescens*. *Plant Physiol* 119: 305–311
- Küpper H, Lombi E, Zhao FJ, McGrath SP (2000) Cellular compartmentation of cadmium and zinc in relation to other elements in the hyperaccumulator *Arabidopsis halleri*. *Planta* 212:75–84
- Küpper H, Lombi E, Zhao FJ, Wieshammer G, McGrath SP (2001) Cellular compartmentation of nickel in the hyperaccumulators *Alyssum lesbiacum*, *Alyssum bertolonii* and *Thlaspi goesingense*. *J Exp Bot* 52: 2291–2300
- Lambers H, Hayes PE, Laliberte E, Oliveira RS, Turner BL (2015) Leaf manganese accumulation and phosphorus-acquisition efficiency. *Trends Plant Sci* 20:83–90
- Lasat MM, Pence NS, Garvin DF, Ebbs SD, Kochian LV (2000) Molecular physiology of zinc transport in the Zn hyperaccumulator *Thlaspi caerulescens*. *J Exp Bot* 51:71–79
- Leigh Broadhurst C, Tappero R, Mangel T, Erbe E, Sparks D, Chaney R (2009) Interaction of nickel and manganese in accumulation and localization in leaves of the Ni hyperaccumulators *Alyssum murale* and *Alyssum corsicum*. *Plant Soil* 314:35–48
- Leitenmaier B, Küpper H (2013) Compartmentation and complexation of metals in hyperaccumulator plants. *Front Plant Sci* 4:374. doi:10.3389/fpls.2013.00374
- Li T, Xu Z, Han X, Yang X, Sparks DL (2012) Characterization of dissolved organic matter in the rhizosphere of hyperaccumulator *Sedum alfredii* and its effect on the mobility of zinc. *Chemosphere* 88:570–576
- Liang J, Shohag MJI, Yang X, Tian S, Zhang Y, Feng Y, He Z (2014) Role of sulfur assimilation pathway in cadmium hyperaccumulation by *Sedum alfredii* Hance. *Ecotoxicol Environ Saf* 100:159–165
- Lin YF, Liang HM, Yang SY, Boch A, Clemens S, Chen CC, JF W, Huang JL, Yeh KC (2009) *Arabidopsis* IRT3 is a zinc-regulated and plasma membrane localized zinc/iron transporter. *New Phytol* 182: 392–404
- Lin YF, Hassan Z, Talukdar S, Schat H, Aarts MG (2016) Expression of the ZNT1 zinc transporter from the metal hyperaccumulator *Noccaea caerulea*

- confers enhanced zinc and cadmium tolerance and accumulation to *Arabidopsis thaliana*. PLoS One 11: e0149750
- Lombi E, Zhao F-J, Fuhrmann M, Ma LQ, McGrath SP (2002) Arsenic distribution and speciation in the fronds of the hyperaccumulator *Pteris vittata*. New Phytol 156:195–203
- Losfeld G, L'Huillier L, Fogliani B, Mc Coy S, Grison C, Jaffré T (2015) Leaf-age and soil-plant relationships: key factors for reporting trace-elements hyperaccumulation by plants and design applications. Environ Sci Pollut Res Int 22:5620–5632
- Lu L, Tian S, Zhang J, Yang X, Labavitch JM, Webb SM, Latimer M, Brown PH (2013) Efficient xylem transport and phloem remobilization of Zn in the hyperaccumulator plant species *Sedum alfredii*. New Phytol 198:721–731
- Lu L, Liao X, Labavitch J, Yang X, Nelson E, Du Y, Brown PH, Tian S (2014) Speciation and localization of Zn in the hyperaccumulator *Sedum alfredii* by extended X-ray absorption fine structure and micro-X-ray fluorescence. Plant Physiol Biochem 84: 224–232
- Lucisine P, Echevarria G, Sterckeman T, Vallance J, Rey P, Benizri E (2014) Effect of hyperaccumulating plant cover composition and rhizosphere-associated bacteria on the efficiency of nickel extraction from soil. Appl Soil Ecol 81:30–36
- Luo SL, Chen L, Chen JL, Xiao X, TY X, Wan Y, Rao C, Liu CB, Liu YT, Lai C, Zeng GM (2011) Analysis and characterization of cultivable heavy metal-resistant bacterial endophytes isolated from Cd-hyperaccumulator *Solanum nigrum* L. and their potential use for phytoremediation. Chemosphere 85: 1130–1138
- Ma Y, Prasad MNV, Rajkumar M, Freitas H (2011a) Plant growth promoting rhizobacteria and endophytes accelerate phytoremediation of metalliferous soils. Biotechnol Adv 29:248–258
- Ma Y, Rajkumar M, Luo Y, Freitas H (2011b) Inoculation of endophytic bacteria on host and non-host plants—effects on plant growth and Ni uptake. J Hazard Mater 195:230–237
- Ma Y, Oliveira RS, Nai F, Rajkumar M, Luo Y, Rocha I, Freitas H (2015) The hyperaccumulator *Sedum plumbizincicola* harbors metal-resistant endophytic bacteria that improve its phytoextraction capacity in multi-metal contaminated soil. J Environ Manage 156: 62–69
- Mari S, Gendre D, Pianelli K, Ouerdane L, Lobinski R, Briat JF, Lebrun M, Czerniec P (2006) Root-to-shoot long-distance circulation of nicotianamine and nicotianamine-nickel chelates in the metal hyperaccumulator *Thlaspi caerulescens*. J Exp Bot 57: 4111–4122
- Martos S, Gallego B, Saez L, Lopez-Alvarado J, Cabot C, Poschenrieder C (2016) Characterization of zinc and cadmium hyperaccumulation in three *Noccaea* (Brassicaceae) populations from non-metalliferous sites in the eastern Pyrenees. Front Plant Sci 7:128. doi:10.3389/fpls.2016.00128
- McNear DH Jr, Chaney RL, Sparks DL (2010) The hyperaccumulator *Alyssum murale* uses complexation with nitrogen and oxygen donor ligands for Ni transport and storage. Phytochemistry 71:188–200
- Mengoni A, Pini F, Huang LN, Shu WS, Bazzicalupo M (2009a) Plant-by-plant variations of bacterial communities associated with leaves of the nickel hyperaccumulator *Alyssum bertolonii*. Desv. Microb Ecol 58:660–667
- Mengoni A, Schat H, Vangronsveld J (2009b) Plants as extreme environments? Ni-resistant bacteria and Ni-hyperaccumulators of serpentine flora. Plant Soil 331:5–16
- Menguer PK, Farthing E, Peaston KA, Ricachenevsky FK, Fett JP, Williams LE (2013) Functional analysis of the rice vacuolar zinc transporter OsMTP1. J Exp Bot 64:2871–2883
- Merlot S, Hannibal L, Martins S, Martinelli L, Amir H, Lebrun M, Thomine S (2014) The metal transporter PgIREG1 from the hyperaccumulator *Psychotria gabriellae* is a candidate gene for nickel tolerance and accumulation. J Exp Bot 65:1551–1564
- Meyer CL, Verbruggen N (2012) The use of the model species *Arabidopsis halleri* towards phytoextraction of cadmium polluted soils. Nat Biotechnol 30:9–14
- Meyer CL, Vitalis R, Saumitou-Laprade P, Castric V (2009) Genomic pattern of adaptive divergence in *Arabidopsis halleri*, a model species for tolerance to heavy metal. Mol Ecol 18:2050–2062
- Meyer CL, Kostecka AA, Saumitou-Laprade P, Creach A, Castric V, Pauwels M, Frerot H (2010) Variability of zinc tolerance among and within populations of the pseudometallophyte species *Arabidopsis halleri* and possible role of directional selection. New Phytol 185:130–142
- Meyer CL, Juraniec M, Huguet S, Chaves-Rodriguez E, Salis P, Isaure M-P, Goormaghtigh E, Verbruggen N (2015) Intraspecific variability of cadmium tolerance and accumulation, and cadmium-induced cell wall modifications in the metal hyperaccumulator *Arabidopsis halleri*. J Exp Bot 66:3215–3227
- Meyer CL, Pauwels M, Briset L, Godé C, Salis P, Bourceaux A, Souleman D, Frerot H, Verbruggen N (2016) Potential preadaptation to anthropogenic pollution: evidence from a common quantitative trait locus for zinc and cadmium tolerance in metallicolous and nonmetallicolous accessions of *Arabidopsis halleri*. New Phytol 212:934–943
- Migocka M, Kosieradzka A, Papierniak A, Maciaszczyk-Dziubinska E, Posyniak E, Garbiec A, Filleur S (2015) Two metal-tolerance proteins, MTP1 and MTP4, are involved in Zn homeostasis and Cd sequestration in cucumber cells. J Exp Bot 66:1001–1015
- Milner MJ, Kochian LV (2008) Investigating heavy-metal hyperaccumulation using *Thlaspi caerulescens* as a model system. Ann Bot 102:3–13
- Milner MJ, Craft E, Yamaji N, Koyama E, Ma JF, Kochian LV (2012) Characterization of the high affinity Zn transporter from *Noccaea caerulescens*, NcZNT1, and dissection of its promoter for its role

- in Zn uptake and hyperaccumulation. *New Phytol* 195:113–123
- Milner MJ, Mitani-Ueno N, Yamaji N, Yokosho K, Craft E, Fei Z, Ebbs S, Zambrano MC, Ma JF, Kochian LV (2014) Root and shoot transcriptome analysis of two ecotypes of *Noccaea caerulea* uncovers the role of *NcNramp1* in Cd hyperaccumulation. *Plant J* 78:398–410
- Mirouze M, Sels J, Richard O, Czernic P, Loubet S, Jacquier A, Francois IE, Cammue BP, Lebrun M, Berthomieu P, Marques L (2006) A putative novel role for plant defensins: a defensin from the zinc hyper-accumulating plant, *Arabidopsis halleri*, confers zinc tolerance. *Plant J* 47:329–342
- Mizuno T, Usui K, Horie K, Nosaka S, Mizuno N, Obata H (2005) Cloning of three ZIP/Nramp transporter genes from a Ni hyperaccumulator plant *Thlaspi japonicum* and their Ni²⁺-transport abilities. *Plant Physiol Biochem* 43:793–801
- Mizuno T, Usui K, Nishida S, Unno T, Obata H (2007) Investigation of the basis for Ni tolerance conferred by the expression of TjZnt1 and TjZnt2 in yeast strains. *Plant Physiol Biochem* 45:371–378
- Molitor M, Dechamps C, Gruber W, Meerts P (2005) *Thlaspi caerulescens* on nonmetalliferous soil in Luxembourg: ecological niche and genetic variation in mineral element composition. *New Phytol* 165: 503–512
- Monsant AC, Kappen P, Wang Y, Pigram PJ, Baker AJM, Tang C (2011) In vivo speciation of zinc in *Noccaea caerulea* in response to nitrogen form and zinc exposure. *Plant Soil* 348:167–183
- Morrissey J, Baxter IR, Lee J, Li L, Lahner B, Grotz N, Kaplan J, Salt DE, Gueriot ML (2009) The ferro-transport metal efflux proteins function in iron and cobalt homeostasis in *Arabidopsis*. *Plant Cell* 21:3326–3338
- Muehe EM, Weigold P, Adaktylou IJ, Planer-Friedrich B, Krämer U, Kappler A, Behrens S (2015) Rhizosphere microbial community composition affects cadmium and zinc uptake by the metal-hyperaccumulating plant *Arabidopsis halleri*. *Appl Environ Microbiol* 81:2173–2181
- Nguyen NNT, Ranwez V, Vile D, Soulie MC, DellagiAlia ED, Gosti F (2014) Evolutionary tinkering of the expression of PDF1s suggests their joint effect on zinc tolerance and the response to pathogen attack. *Front Plant Sci* 5:70. doi:10.3389/fpls.2014.00070
- Nishida S, Tsuzuki C, Kato A, Aisu A, Yoshida J, Mizuno T (2011) AtIRT1, the primary iron uptake transporter in the root, mediates excess nickel accumulation in *Arabidopsis thaliana*. *Plant Cell Physiol* 52: 1433–1442
- Nishida S, Aisu A, Mizuno T (2012) Induction of IRT1 by the nickel-induced iron-deficient response in *Arabidopsis*. *Plant Signal Behav* 7:329–331
- Nonnoi F, Chinnaswamy A, García de la Torre VS, Coba de la Peña T, Lucas MM, Pueyo JJ (2012) Metal tolerance of rhizobial strains isolated from nodules of herbaceous legumes (*Medicago* spp. and *Trifolium* spp.) growing in mercury-contaminated soils. *Appl Soil Ecol* 61:49–59
- Nouet C, Charlier JB, Carnol M, Bosman B, Farnir F, Motte P, Hanikenne M (2015) Functional analysis of the three *HMA4* copies of the metal hyperaccumulator *Arabidopsis halleri*. *J Exp Bot* 66:5783–5795
- O’Lochlainn S, Bowen HC, Fray RG, Hammond JP, King GJ, White PJ, Graham NS, Broadley MR (2011) Tandem quadruplication of *HMA4* in the zinc (Zn) and cadmium (Cd) hyperaccumulator *Noccaea caerulea*. *PLoS One* 6:e17814
- Oomen RJ, Wu J, Lelievre F, Blanchet S, Richaud P, Barbier-Brygoo H, Aarts MG, Thomine S (2009) Functional characterization of NRAMP3 and NRAMP4 from the metal hyperaccumulator *Thlaspi caerulescens*. *New Phytol* 181:637–650
- Oomen RJ, Seveno-Carpentier E, Ricodeau N, Bournaud C, Conejero G, Paris N, Berthomieu P, Marques L (2011) Plant defensin AhPDF1.1 is not secreted in leaves but it accumulates in intracellular compartments. *New Phytol* 192:140–150
- Palmer CM, Gueriot ML (2009) Facing the challenges of Cu, Fe and Zn homeostasis in plants. *Nat Chem Biol* 5:333–340
- Pankiewicz VC, Camilios-Neto D, Bonato P, Balsanelli E, Tadra-Sfeir MZ, Faoro H, Chubatsu LS, Donatti L, Wajnberg G, Passetti F, Monteiro RA, Pedrosa FO, Souza EM (2016) RNA-seq transcriptional profiling of *Herbaspirillum seropedicae* colonizing wheat (*Triticum aestivum*) roots. *Plant Mol Biol* 90:589–603
- Papoyan A, Kochian LV (2004) Identification of *Thlaspi caerulescens* genes that may be involved in heavy metal hyperaccumulation and tolerance. Characterization of a novel heavy metal transporting ATPase. *Plant Physiol* 136:3814–3823
- Pauwels M, Roossens N, Frérot H, Saumitou-Laprade P (2008) When population genetics serves genomics: putting adaptation back in a spatial and historical context. *Curr Opin Plant Biol* 11:129–134
- Pauwels M, Vekemans X, Gode C, Frérot H, Castric V, Saumitou-Laprade P (2012) Nuclear and chloroplast DNA phylogeography reveals vicariance among European populations of the model species for the study of metal tolerance, *Arabidopsis halleri* (Brassicaceae). *New Phytol* 193:916–928
- Pedersen CNS, Axelsen KB, Harper JF, Palmgren MG (2012) Evolution of plant P-type ATPases. *Front Plant Sci* 3:31. doi:10.3389/fpls.2012.00031
- Peer WA, Mamoudian M, Lahner B, Reeves RD, Murphy AS, Salt DE (2003) Identifying model metal hyper-accumulating plants: germplasm analysis of 20 Brassicaceae accessions from a wide geographical area. *New Phytol* 159:421–430
- Pence NS, Larsen PB, Ebbs SD, Letham DL, Lasat MM, Garvin DF, Eide D, Kochian LV (2000) The molecular physiology of heavy metal transport in the Zn/Cd hyperaccumulator *Thlaspi caerulescens*. *Proc Natl Acad Sci USA* 97:4956–4960
- Peng JS, Wang YJ, Ding G, Ma HL, Zhang YJ, Gong JM (2017) A pivotal role of cell wall in Cd accumulation in the Crassulaceae hyperaccumulator *Sedum plumbizincicola*. *Mol Plant*. doi:10.1016/j.molp.2016.12.007

- Persans MW, Yan X, Patnoe JM, Krämer U, Salt DE (1999) Molecular dissection of the role of histidine in nickel hyperaccumulation in *Thlaspi goesingense* (Halácsy). *Plant Physiol* 121:1117–1126
- Persans MW, Nieman K, Salt DE (2001) Functional activity and role of cation-efflux family members in Ni hyperaccumulation in *Thlaspi goesingense*. *Proc Natl Acad Sci USA* 98:9995–10000
- Pianelli K, Mari S, Marques L, Lebrun M, Czernic P (2005) Nicotianamine over-accumulation confers resistance to nickel in *Arabidopsis thaliana*. *Transgenic Res* 14:739–748
- Pickering IJ, Wright C, Bubner B, Ellis D, Persans MW, EY Y, George GN, Prince RC, Salt DE (2003) Chemical form and distribution of selenium and sulfur in the selenium hyperaccumulator *Astragalus bisulcatus*. *Plant Physiol* 131:1460–1467
- Pittman JK (2005) Managing the manganese: molecular mechanisms of manganese transport and homeostasis. *New Phytol* 167:733–742
- Polacco JC, Mazzafera P, Tezotto T (2013) Opinion: nickel and urease in plants: still many knowledge gaps. *Plant Sci* 199–200:79–90
- Pottier M, Oomen R, Picco C, Giraudat J, Scholz-Starke J, Richaud P, Carpaneto A, Thomine S (2015) Identification of mutations allowing Natural Resistance Associated Macrophage Proteins (NRAMP) to discriminate against cadmium. *Plant J* 83:625–637
- Reeves RD, Schwartz C, Morel JL, Edmondson J (2001) Distribution and metal-accumulating behavior of *Thlaspi caerulescens* and associated metallophytes in France. *Int J Phytoremediation* 3:145–172
- Reinhold-Hurek B, Hurek T (2011) Living inside plants: bacterial endophytes. *Curr Opin Plant Biol* 14:435–443
- Rellán-Alvarez R, Abadía J, Alvarez-Fernandez A (2008) Formation of metal-nicotianamine complexes as affected by pH, ligand exchange with citrate and metal exchange. A study by electrospray ionization time-of-flight mass spectrometry. *Rapid Commun Mass Spectrom* 22:1553–1562
- Richau KH, Kozhevnikova AD, Seregin IV, Vooijs R, Koevoets PL, Smith JA, Ivanov VB, Schat H (2009) Chelation by histidine inhibits the vacuolar sequestration of nickel in roots of the hyperaccumulator *Thlaspi caerulescens*. *New Phytol* 183:106–116
- Rogers EE, Gueriot ML (2002) FRD3, a member of the multidrug and toxin efflux family, controls iron deficiency responses in *Arabidopsis*. *Plant Cell* 14:1787–1799
- Rogers EE, Eide DJ, Gueriot ML (2000) Altered selectivity in an *Arabidopsis* metal transporter. *Proc Natl Acad Sci USA* 97:12356–12360
- Roosens N, Verbruggen N, Meerts P, Ximénez de Embún P, Smith JAC (2003) Natural variation in cadmium hyperaccumulation and its relationship to metal hyperaccumulation for seven populations of *Thlaspi caerulescens* from western Europe. *Plant Cell Environ* 26:1657–1672
- Roosens NH, Willems G, Saumitou-Laprade P (2008) Using *Arabidopsis* to explore zinc tolerance and hyperaccumulation. *Trends Plant Sci* 13:208–215
- Roux C, Castric V, Pauwels M, Wright SI, Saumitou-Laprade P, Vekemans X (2011) Does speciation between *Arabidopsis halleri* and *Arabidopsis lyrata* coincide with major changes in a molecular target of adaptation? *PLoS One* 6:e26872
- Sarret G, Saumitou-Laprade P, Bert V, Proux O, Hazemann JL, Traverse A, Marcus MA, Manceau A (2002) Forms of zinc accumulated in the hyperaccumulator *Arabidopsis halleri*. *Plant Physiol* 130:1815–1826
- Sarret G, Willems G, Isaure MP, Marcus MA, Fakra SC, Frerot H, Pairis S, Geoffroy N, Manceau A, Saumitou-Laprade P (2009) Zinc distribution and speciation in *Arabidopsis halleri* x *Arabidopsis lyrata* progenies presenting various zinc accumulation capacities. *New Phytol* 184:581–595
- Sarret G, Smits E, Michel HC, Isaure MP, Zhao FJ, Tappero R (2013) Use of synchrotron-based techniques to elucidate metal uptake and metabolism in plants. In: Sparks DL (ed) *Advances in agronomy*, vol 119. Elsevier Academic, San Diego
- Schaaf G, Honsbein A, Meda AR, Kirchner S, Wipf D, von Wiren N (2006) AtIREG2 encodes a tonoplast transport protein involved in iron-dependent nickel detoxification in *Arabidopsis thaliana* roots. *J Biol Chem* 281:25532–25540
- Schaumölfel D, Ouerdane L, Bouyssi  re B, Lobinski R (2003) Speciation analysis of nickel in the latex of a hyperaccumulating tree *Sebertia acuminata* by HPLC and CZE with ICP MS and electrospray MS-MS detection. *J Anal At Spectrom* 18:120–127
- Schiavon M, Pilon-Smits EA (2016) The fascinating facets of plant selenium accumulation—biochemistry, physiology, evolution and ecology. *New Phytol.* doi:10.1111/nph.14378
- Schiavon M, Pilon M, Malagoli M, Pilon-Smits EAH (2015) Exploring the importance of sulfate transporters and ATP sulphurylases for selenium hyperaccumulation—a comparison of *Stanleya pinnata* and *Brassica juncea* (Brassicaceae). *Front Plant Sci* 6:1–13
- Sessitsch A, Hardoim P, D  ring J, Weilharter A, Krause A, Woyke T, Mitter B, Hauberg-Lotte L, Friedrich F, Rahalkar M, Hurek T, Sarkar A, Bodrossy L, van Overbeek L, Br  r D, van Elsas JD, Reinhold-Hurek B (2012) Functional characteristics of an endophyte community colonizing rice roots as revealed by metagenomic analysis. *Mol Plant Microbe Interact* 25:28–36
- Shahzad Z, Gosti F, Fr  rot H, Lacombe E, Roosens N, Saumitou-Laprade P, Berthomieu P (2010) The five AhMTP1 zinc transporters undergo different evolutionary fates towards adaptive evolution to zinc tolerance in *Arabidopsis halleri*. *PLoS Genet* 6:e1000911
- Shahzad Z, Ranwez V, Fizames C, Marqu  s L, Le Martret B, Alassimone J, God   C, Lacombe E,

- Castillo T, Saumitou-Laprade P, Berthomieu P, Gosti F (2013) Plant defensin type 1 (PDF1): protein promiscuity and expression variation within the *Arabidopsis* genus shed light on zinc tolerance acquisition in *Arabidopsis halleri*. *New Phytol* 200: 820–833
- Shibagaki N, Rose A, McDermott JP, Fujiwara T, Hayashi H, Yoneyama T, Davies JP (2002) Selenate-resistant mutants of *Arabidopsis thaliana* identify Sultr1;2, a sulfate transporter required for efficient transport of sulfate into roots. *Plant J* 29:475–486
- Socha AL, Guerinot ML (2014) Mn-euvering manganese: the role of transporter gene family members in manganese uptake and mobilization in plants. *Front Plant Sci* 5:106. doi:10.3389/fpls.2014.00106
- Takahashi H, Watanabe-Takahashi A, Smith FW, Blake-Kalff M, Hawkesford MJ, Saito K (2000) The roles of three functional sulphate transporters involved in uptake and translocation of sulphate in *Arabidopsis thaliana*. *Plant J* 23:171–182
- Talke IN, Hanikenne M, Krämer U (2006) Zinc-dependent global transcriptional control, transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of the hyperaccumulator *Arabidopsis halleri*. *Plant Physiol* 142:148–167
- The Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408:796–815
- Thomine S, Vert G (2013) Iron transport in plants: better be safe than sorry. *Curr Opin Plant Biol* 16:322–327
- Tian S, Lu L, Labavitch J, Yang X, He Z, Hu H, Sarangi R, Newville M, Commisso J, Brown P (2011) Cellular sequestration of cadmium in the hyperaccumulator plant species *Sedum alfredii*. *Plant Physiol* 157:1914–1925
- Tsednee M, Yang S-C, Lee D-C, Yeh K-C (2014) Root-secreted nicotianamine from *Arabidopsis halleri* facilitates zinc hypertolerance by regulating zinc bio-availability. *Plant Physiol* 166:839–852
- Turner TL, Bourne EC, Von Wettberg EJ, TT H, Nuzhdin SV (2010) Population resequencing reveals local adaptation of *Arabidopsis lyrata* to serpentine soils. *Nat Genet* 42:260–263
- Ueno D, Milner MJ, Yamaji N, Yokosho K, Koyama E, Clemencia Zambrano M, Kaskie M, Ebbs S, Kochian LV, Ma JF (2011) Elevated expression of TcHMA3 plays a key role in the extreme Cd tolerance in a Cd-hyperaccumulating ecotype of *Thlaspi caerulescens*. *Plant J* 66:852–862
- van de Mortel JE, Almar Villanueva L, Schat H, Kwekkeboom J, Coughlan S, Moerland PD, Ver Loren van Themaat E, Koornneef M, Aarts MG (2006) Large expression differences in genes for iron and zinc homeostasis, stress response and lignin biosynthesis distinguish *Arabidopsis thaliana* and the related metal hyperaccumulator *Thlaspi caerulescens*. *Plant Physiol* 142:1127–1147
- van de Mortel JE, Schat H, Moerland PD, Ver Loren van Themaat E, van der Ent S, Blankestijn H, Ghandilyan A, Tsiatsiani S, Aarts MG (2008) Expression differences for genes involved in lignin, glutathione and sulphate metabolism in response to cadmium in *Arabidopsis thaliana* and the related Zn/Cd-hyperaccumulator *Thlaspi caerulescens*. *Plant Cell Environ* 31:301–324
- van der Ent A, Baker AJM, Reeves RD, Pollard AJ, Schat H (2013) Hyperaccumulators of metal and metalloid trace elements: facts and fiction. *Plant Soil* 362: 319–334
- van der Weerden NL, Anderson MA (2013) Plant defensins: common fold, multiple functions. *Fungal Biol Rev* 26:121–131
- Verbruggen N, Hermans C, Schat H (2009a) Mechanisms to cope with arsenic or cadmium excess in plants. *Curr Opin Plant Biol* 12:364–372
- Verbruggen N, Hermans C, Schat H (2009b) Molecular mechanisms of metal hyperaccumulation in plants. *New Phytol* 181:759–776
- Verbruggen N, Hanikenne M, Clemens S (2013) A more complete picture of metal hyperaccumulation through next-generation sequencing technologies. *Front Plant Sci* 4:388
- Vert G, Grotz N, Dedaldechamp F, Gaymard F, Guerinot ML, Briat JF, Curie C (2002) IRT1, an *Arabidopsis* transporter essential for iron uptake from the soil and for plant growth. *Plant Cell* 14:1223–1233
- Villafort Carvalho MT, Amaral DC, Guilherme LR, Aarts MG (2013) *Gomphrena clausenii*, the first South-American metallophyte species with indicator-like Zn and Cd accumulation and extreme metal tolerance. *Front Plant Sci* 4:180. doi:10.3389/fpls.2013.00180
- Villafort Carvalho MT, Pongrac P, Mumm R, van Arkel J, van Aelst A, Jeromel L, Vavpetic P, Pelicon P, Aarts MG (2015) *Gomphrena clausenii*, a novel metal-hypertolerant bioindicator species, sequesters cadmium, but not zinc, in vacuolar oxalate crystals. *New Phytol* 208:763–775
- Visioli G, D'Egidio S, Vamerali T, Mattarozzi M, Sanangelantoni AM (2014) Culturable endophytic bacteria enhance Ni translocation in the hyperaccumulator *Noccaea caerulescens*. *Chemosphere* 117:538–544
- Visioli G, Vamerali T, Mattarozzi M, Dramis L, Sanangelantoni AM (2015) Combined endophytic inoculants enhance nickel phytoextraction from serpentine soil in the hyperaccumulator *Noccaea caerulescens*. *Front Plant Sci* 6:1–12
- Weber M, Harada E, Vess C, Roepenack-Lahaye EV, Clemens S (2004) Comparative microarray analysis of *Arabidopsis thaliana* and *Arabidopsis halleri* roots identifies nicotianamine synthase, a ZIP transporter and other genes as potential metal hyperaccumulation factors. *Plant J* 37:269–281
- Weber M, Trampczynska A, Clemens S (2006) Comparative transcriptome analysis of toxic metal responses in *Arabidopsis thaliana* and the Cd²⁺-hypertolerant facultative metallophyte *Arabidopsis halleri*. *Plant Cell Environ* 29:950–963
- Wei W, Chai T, Zhang Y, Han L, Xu J, Guan Z (2009) The *Thlaspi caerulescens* NRAMP homologue TcNRAMP3 is capable of divalent cation transport. *Mol Biotechnol* 41:15–21

- White PJ (2016) Selenium accumulation by plants. *Ann Bot* 117:217–235
- Willems G, Dräger DB, Courbot M, Gode C, Verbruggen N, Saumitou-Laprade P (2007) The genetic basis of zinc tolerance in the metallophyte *Arabidopsis halleri* ssp. *halleri* (Brassicaceae): an analysis of quantitative trait loci. *Genetics* 176: 659–674
- Willems G, Frérot H, Gennen J, Salis P, Saumitou-Laprade P, Verbruggen N (2010) Quantitative trait loci analysis of mineral element concentrations in an *Arabidopsis halleri* x *Arabidopsis lyrata petraea* F₂ progeny grown on cadmium-contaminated soil. *New Phytol* 187:368–379
- Wong CKE, Cobbett CS (2009) HMA P-type ATPases are the major mechanism for root-to-shoot Cd translocation in *Arabidopsis thaliana*. *New Phytol* 181: 71–78
- Wu LH, Liu YJ, Zhou SB, Guo FG, Bi D, Guo XH, Baker AJM, Smith JAC, Luo YM (2013) *Sedum plumbizincicola* X.H. Guo et S.B. Zhou ex L.H. Wu (Crassulaceae): a new species from Zhejiang Province, China. *Plant Syst Evol* 299:487–498
- Wycisk K, Kim EJ, Schroeder JI, Krämer U (2004) Enhancing the first enzymatic step in the histidine biosynthesis pathway increases the free histidine pool and nickel tolerance in *Arabidopsis thaliana*. *FEBS Lett* 578:128–134
- Yang XE, Long XX, Ye HB, He ZL, Calvert DV, Stoffella PJ (2004) Cadmium tolerance and hyperaccumulation in a new Zn-hyperaccumulating plant species (*Sedum alfredii* Hance). *Plant Soil* 259: 181–189
- Yang X, Li T, Yang J, He Z, Lu L, Meng F (2006) Zinc compartmentation in root, transport into xylem, and absorption into leaf cells in the hyperaccumulating species of *Sedum alfredii* Hance. *Planta* 224:185–195
- Yogeeswaran K, Frary A, York TL, Amenta A, Lesser AH, Nasrallah JB, Tanksley SD, Nasrallah ME (2005) Comparative genome analyses of *Arabidopsis* spp.: inferring chromosomal rearrangement events in the evolutionary history of *A. thaliana*. *Genome Res* 15: 505–515
- Yokosho K, Yamaji N, Ueno D, Mitani N, Ma JF (2009) OsFRDL1 is a citrate transporter required for efficient translocation of iron in rice. *Plant Physiol* 149: 297–305
- Zhang M, Senoura T, Yang X, Nishizawa NK (2011) Functional analysis of metal tolerance proteins isolated from Zn/Cd hyperaccumulating ecotype and non-hyperaccumulating ecotype of *Sedum alfredii* Hance. *FEBS Lett* 585:2604–2609
- Zhang L, Hu B, Li W, Che R, Deng K, Li H, Yu F, Ling H, Li Y, Chu C (2014) OsPT2, a phosphate transporter, is involved in the active uptake of selenite in rice. *New Phytol* 201:1183–1191
- Zhang Z, Yu Q, Du H, Ai W, Yao X, Mendoza-Cózatl DG, Qiu B (2016) Enhanced cadmium efflux and root-to-shoot translocation are conserved in the hyperaccumulator *Sedum alfredii* (Crassulaceae family). *FEBS Lett* 590:1757–1764
- Zhao XQ, Mitani N, Yamaji N, Shen RF, Ma JF (2010) Involvement of silicon influx transporter OsNIP2;1 in selenite uptake in rice. *Plant Physiol* 153:187–1877
- Zhu YG, Pilon-Smits EA, Zhao FJ, Williams PN, Meharg AA (2009) Selenium in higher plants: understanding mechanisms for biofortification and phytoremediation. *Trends Plant Sci* 14:436–442