

Accepted Manuscript

Infusion of mesenchymal stromal cells after deceased liver transplantation: A phase I-II, open-label, clinical study

Olivier Detry, Morgan Vandermeulen, Marie-Hélène Delbouille, Joan Somja, Noella Bletard, Alexandra Briquet, Chantal Lechanteur, Olivier Giet, Etienne Baudoux, Muriel Hannon, Frederic Baron, Yves Beguin


PII: S0168-8278(17)30131-9
DOI: <http://dx.doi.org/10.1016/j.jhep.2017.03.001>
Reference: JHEPAT 6448

To appear in: *Journal of Hepatology*

Received Date: 12 October 2016
Revised Date: 24 February 2017
Accepted Date: 1 March 2017

Please cite this article as: Detry, O., Vandermeulen, M., Delbouille, M-H., Somja, J., Bletard, N., Briquet, A., Lechanteur, C., Giet, O., Baudoux, E., Hannon, M., Baron, F., Beguin, Y., Infusion of mesenchymal stromal cells after deceased liver transplantation: A phase I-II, open-label, clinical study, *Journal of Hepatology* (2017), doi: <http://dx.doi.org/10.1016/j.jhep.2017.03.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Infusion of mesenchymal stromal cells after deceased liver transplantation: A phase I-II, open-label, clinical study

Olivier Detry^{1,2}, Morgan Vandermeulen^{1,2}, Marie-Hélène Delbouille¹, Joan Somja³, Noella Bletard³, Alexandra Briquet⁴, Chantal Lechanteur⁴, Olivier Giet⁴, Etienne Baudoux⁴, Muriel Hannon⁵, Frederic Baron^{5,6}, Yves Beguin^{5,6}

1. Department of Abdominal Surgery and Transplantation, CHU Liege, University of Liege, (CHU ULg), Belgium

2. Mesenchymal stromal cell In Solid Organ Transplantation (MISOT) consortium

3. Department of Pathology, CHU Liege, University of Liege, (CHU ULg), Belgium

4. Laboratory of Cell and Gene Therapy (LTCG), CHU Liege, University of Liege, (CHU ULg), Belgium

5. Interdisciplinary Cluster for Applied Genoproteomics (GIGA)-I3-hematology, University of Liege, Belgium

6. Department of Haematology, CHU Liege, University of Liege, (CHU ULg), Belgium

Corresponding author:

Dr Olivier Detry, Department of Abdominal Surgery and Transplantation, CHU Liege, University of Liege, Belgium. Tel: +3243667645; Fax: +3243667069; Email: olivier.detry@transplantation.be

Electronic word count: 6230

Number of figures and tables: 8

Abbreviations: LT: liver transplantation; MSC: mesenchymal stromal cell; SOT: solid organ transplantation; Treg: regulatory T cell; ET-DRI: Eurotransplant donor risk index; BMI: body mass index; MELD: Model for End-Stage Liver Disease; MMF : mycophenolate mofetyl; CMV: cytomegalovirus; BM: bone marrow; HLA: human leucocyte antigen; LTCG: Laboratory of Cell and Gene Therapy; MLR: mixed-lymphocyte reaction; CRP: C reactive protein; INR: international normalized ratio; CDC: complement-dependent cytotoxicity; HCC: hepatocellular carcinoma; MP: mycophenolic acid.

Keywords: Stem cells; mesenchymal stem cells; cell therapy; liver failure; liver diseases; hepatic insufficiency; cirrhosis; tolerance; cancer; MISOT.

Conflicts of interest: The authors have no conflict of interest related to this study.

Financial support: This study was supported by the Fonds d'investissement de recherche scientifique (FIRS) du CHU de Liege, the Fonds Léon Frédéricq of the Medical School of the University of Liege, and through a Senior Research Grant from the European Society for Organ Transplantation (ESOT). The LTCG is supported in part by the FNRS and the “Federal cancer plan – Belgium”. MV is a PhD fellow sponsored by the FNRS. FB is a senior research associate of the FNRS, Belgium.

Author's contribution: OD designed the study, collected and analysed the data, and wrote the manuscript; MV and MD collected and analysed the data; JS and NB analysed the graft biopsies; AB, CL, OG, EB produced MSC in a GMP-compatible environment MH and FB designed and performed the Treg subphenotyping analyses. YB designed the study, supervised the MSC production and co-wrote the manuscript. All authors approved the final version of the manuscript.

Acknowledgements: The authors value and thank all the members (physicians, nurses, coordinators) to the liver transplantation program of the CHU de Liege, who greatly contributed to this study, and the technicians of the LTCG. The authors thank also Jacques Foguene and André Gothot for the lymphocyte phenotyping.

Clinical Trial number: Eudract: # 2011-001822-81, ClinicalTrials.gov: # NCT 01429038

Lay summary:

Therapy with mesenchymal stromal cells (MSCs) has been proposed as a mean to improve results of solid organ transplantation. One of the potential MSC role could be to induce tolerance after liver transplantation, i.e. allowing the cessation of several medications with severe side effects. This study is the first-in-man use of MSC therapy in 10 liver transplant recipients. This study did not show toxicity after a single MSC infusion but it was not sufficient to allow withdrawal of immunosuppression.

Abstract word count: 269

Abstract

Background & aim: Mesenchymal stromal cell (MSC) infusion could be a mean to establish tolerance in solid organ recipients. The aim of this prospective, controlled, phase-1 study was to evaluate the feasibility, safety and tolerability of a single infusion of MSCs in liver transplant recipients.

Methods: Ten liver transplant recipients under standard immunosuppression received $1.5-3 \times 10^6$ /kg third-party unrelated MSCs on post-operative day 3 ± 2 , and were prospectively compared to a control group of 10 liver transplant recipients. As primary end-points, MSC infusional toxicity was evaluated, and infectious and cancerous complications were prospectively recorded until month 12 in both groups. As secondary end-points, rejection rate, month-6 graft biopsies, and peripheral blood lymphocyte phenotyping were compared. Progressive immunosuppression weaning was attempted from month 6 to 12 in MSC recipients.

Results: No variation in vital parameters or cytokine release syndrome could be detected during and after MSC infusion. No patient developed impairment of organ functions (including liver graft function) following MSC infusion. No increased rate of opportunistic infection or *de novo* cancer was detected. As secondary end-points, there was no difference in overall rates of rejection or graft survival. Month-6 biopsies did not demonstrate a difference between groups in the evaluation of rejection according to the Banff criteria, in the fibrosis score or in immunohistochemistry (including Tregs). No difference in peripheral blood lymphocyte typing could be detected. The immunosuppression weaning in MSC recipients was not successful.

Conclusions: No side effect of MSC infusion at day 3 after liver transplant could be detected, but this infusion did not promote tolerance. This study opens the way for further MSC or Treg-based trials in liver transplant recipients.

Introduction

Liver transplantation (LT) has become the gold standard treatment of many hepatic end-stage diseases. Long-term graft and patient survivals are now common after LT, but recipients are still submitted to life-long immunosuppression, which impairs quality of life and might reduce survival by promoting cancer development or by increasing the risks for infection, kidney function impairment and cardiovascular diseases. There is therefore a need for improvement in the immunosuppressive protocols after LT. Finding a way to establish donor-specific immunological tolerance without the need for non-specific immunosuppression remains one of the major goals in transplantation medicine [1].

Mesenchymal stromal cells (MSCs) are multipotent progenitors within the bone marrow, capable of differentiating into various cells and tissues, such as chondrocytes, osteoblasts and adipocytes [2]. MSCs can be isolated after *ex vivo* culture of the adherent mononuclear bone marrow cell fraction. In addition to the bone marrow, MSCs reside in the connective tissues of many organs including the liver. After *ex vivo* expansion, human MSCs have a fibroblastic-like morphology, and are uniformly positive for SH2, SH3, CD29, CD44, CD71, CD90, CD105, CD106, CD120a, CD124, and CD166, but are negative for common hematopoietic markers such as CD14, CD45 or CD34 [2]. Human MSCs express HLA-class I and can be induced to express HLA-class II by IFN- γ . A large number of *in vitro* and *in vivo* studies have documented the anti-inflammatory and immunoregulatory properties of MSCs on both the adaptive and innate immune system [3], as well as a potential beneficial effect in ischemia-reperfusion injury [4, 5]. Specifically, MSCs have been shown to decrease effector T-cell response while promoting the emergence of regulatory T-cells (Treg) [6]. These MSC properties suggest that they could be particularly attractive in solid organ transplantation (SOT) [7, 8], and a consortium of European academic centres studying this subject has been

created (<http://www.misot.eu>). A first randomized controlled trial evaluating the effects of autologous MSCs in living related kidney transplantation has been performed in China [9]. In this study, MSCs significantly correlated with fewer acute rejections, a lower risk of opportunistic infections and a better renal function at 1 month. Furthermore, fewer adverse effects were seen in the MSC groups compared to the control group [9]. Compared to other transplanted organs, the liver graft is immunologically protected, and LT recipients are considered the ideal candidates for MSC therapy and for operative tolerance trials after SOT [10]. To date there has been no published trial evaluating MSC infusion in a series of LT patients [1].

Despite the absence of major adverse effects in the preliminary clinical trials evaluating MSC-based therapy to date [11], clinical infusion of MSCs might theoretically be complicated by impairment of pulmonary function due to MSC embolism in the lung vasculature [12] and by a cytokine-release syndrome [13]. In addition, as MSCs are potentially immunosuppressive, another concern is the potential emergence of higher rates of opportunistic infections and induced cancers after MSC infusion in SOT recipients under immunosuppression. In a small European clinical series, MSC infusion in kidney recipients was associated with transient renal dysfunction [14] and opportunistic infections [15]. It is also possible that MSC injection promotes liver fibrosis [16]. Finally, *in vitro* MSC expansion and culture might generate genomic instability and chromosomal aberrations with a potential risk of MSC neoplastic transformation [17, 18].

The aim of this study was to evaluate the feasibility, the safety and the tolerability of a single MSC infusion after LT in a first-in-man, prospective, controlled, phase-1 study. The primary endpoints were set to clinically detect potential side effects of MSC infusion, as well as the occurrence of infectious and malignant complications. As secondary end-points, the potential

immune-regulative effects of MSCs and the impact of MSCs on Treg counts and phenotype were analysed by comparison with a control group. In addition, in a second step, progressive immunosuppressive withdrawal was attempted as a phase 2 study in stable patients who received MSCs, to evaluate if a single infusion of MSCs might induce operative tolerance after LT.

Methods

Study Design

This study was a monocentric, prospective, non-randomized, controlled, open-label trial. Protocol inclusion and exclusion criteria are presented in Table 1. Between March 2012 and February 2014, 10 stable and low-risk LT recipients under standard immunosuppression received $1.5 \times 10^6/\text{kg}$ third-party MSCs on post-operative day 3 ± 2 (MSC group). These patients were prospectively compared to a control group of 10 LT recipients who fulfilled the study inclusion criteria, declined to receive MSCs, but accepted to be included in the trial as control patients during the same period (Control group). In addition, in patients from the MSC group who did not develop rejection and had normal graft function and month-6 graft biopsy, progressive weaning of immunosuppression was attempted (Fig 1). Weaning of immunosuppression was not considered in the control group as it is well established that early (<1 year) immunosuppression withdrawal is not possible and unethical in LT recipients under regular immunosuppression protocols. The study protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki and was approved by the local Ethics Committee and by the Belgian Federal Agency for Medicines and Health Products (Eudract # 2011-001822-81). The study was registered at ClinicalTrials.gov (protocol # NCT 01429038). Written informed consent was obtained from each MSC donor and LT patient.

Liver transplant procedures and postoperative management

The following deceased liver graft donor characteristics were prospectively collected: age, gender, donation after brain or circulatory death, Eurotransplant donor risk index (ET-DRI) [19], cause of brain damage, terminal blood sodium level, terminal liver function tests, need for vasopressors, length of intensive care unit stay, body mass index (BMI), last 24-hour diuresis, and past cardiopulmonary resuscitation.

The LT procedures were regular deceased LT as performed in the authors' centre [20, 21]. The following LT recipient characteristics were collected: age, gender, BMI, LT indication, and the laboratory Model for End-Stage Liver Disease (MELD) score at admission for transplantation. Cold and total graft ischemic times were recorded. The immunosuppressive regimen consisted of a triple therapy of tacrolimus, mycophenolate mofetyl (MMF) and steroids. The tacrolimus dose was adapted according to trough whole blood values (between 8 and 12 ng/ml the first 28 days and between 5 and 8 thereafter) until day 180 in both groups. In the MSC group, if a rejection episode had not been suspected based on the liver tests and month 6 biopsy, tacrolimus was progressively tapered from day 180 to be discontinued by day 270 in the absence of rejection (Fig 1). A graft biopsy was performed at day 270 \pm 15 in the MSC group. MMF was administered orally from day 1 through day 270 at the dose of 500 mg bid. In the MSC group, if the patient did not develop rejection during tacrolimus withdrawal and at day 270 graft biopsy, MMF was progressively tapered and definitely discontinued by day 365 in the absence of rejection (Fig 1). Steroid treatment consisted of administration of methylprednisolone 500 mg iv before liver graft reperfusion, followed by progressively decreasing daily doses until progressive withdrawal during month 1 (Fig 1). Liver graft rejection was assessed according to standard criteria, including clinical symptoms, blood liver enzymes, and liver graft biopsy if needed. Therapy for rejection included an increase in

tacrolimus administration, boluses of methylprednisolone 500 mg iv per day for 3 days, and anti-thymocyte globulins in steroid-resistant rejection, if needed.

Antibacterial and antiviral prophylaxis was standardized between groups including cefuroxime 3 x 1.5g or piperacillin-tazobactam 4 x 4 g/d for 5 days, prevention of pneumocystis (co-trimoxazole 500 mg po 1/d for three months) and of cytomegalovirus (CMV) infection if indicated (D+, R-, 100 days of valgancyclovir 2 x 450 mg/d po).

MSC donors

Inclusion criteria for MSC donors included: unrelated to the recipient; aged >18 years; no human leucocyte antigen (HLA) matching required; fulfilling generally accepted criteria for allogeneic hematopoietic stem cell donation; and informed consent given. Exclusion criteria were: known allergy to lidocaine; any risk factor for transmissible infectious diseases; meeting generally accepted exclusion criteria for allogeneic hematopoietic stem cell donation [22].

MSC production

MSC expansion cultures were performed and evaluated at the Laboratory of Cell and Gene Therapy (LTCG) of the University Hospital of Liege, CHU of Liege, as previously described [22, 23]. Briefly, bone marrow (BM) (30-50 mL) was collected under local anaesthesia in sterile conditions, and put in sterile heparin-containing syringes. Mononuclear BM cells were isolated by Ficoll (GE Healthcare-Amersham Biosciences AB, Upsala, Sweden), seeded in sterile tissue culture flasks (BD Falcon, Bedford, MA), and cultured in Dulbecco's Modified Eagles Medium-Low Glucose (Invitrogen, Merelbeke, Belgium) with glutamate supplemented with 10% irradiated fetal bovine serum (Hyclone- Perbio Science, Merelbeke, Belgium) and antibiotics (penicillin/streptomycin, Lonza Bio Science, Verviers, Belgium).

Cultures were maintained at 37 °C in humidified atmosphere containing 5% CO₂ for a total of about 4 weeks. The medium was replaced twice a week and, after approximately 2 weeks, the cultures were near confluence (>70%). Cells were then detached by treatment with irradiated trypsin-EDTA (Invitrogen, Merelbeke, Belgium) and replated (passaged) at a lower density to allow further expansion. A second passage was performed when the cells reached confluence again (>70%). At confluence, the cells were harvested, washed, and re-suspended using phosphate-buffered saline (PBS)-EDTA (Miltenyi Biotec, Utrecht, The Netherlands) and Human Serum Albumin (HSA) (CDF-CAF, Brussels, Belgium). The MSCs were then frozen in a medium containing 70% PBS, 20% HSA, and 10% DMSO (WAK-Chemie, Steinbach, Germany) using standard techniques. Before infusion, the MSCs were thawed and diluted in PBS, and then injected into the patients within 60 minutes. All reagents were certified sterile, and endotoxin-free, and had been used in other clinical trials in Europe. In addition, the batch of fetal bovine serum used was selected after extensive testing, and was irradiated to ensure removal of all potential viruses. The following analyses were performed as quality controls for each MSC expansion culture: nucleated cell count on a manual cell counter, flow cytometry analysis with determination of the % cells (out of total cells) positive for CD73, CD90, and CD105, and negative for HLA-DR, CD31, CD80, CD14, CD45, CD3, and CD34; cell viability using trypan blue exclusion; microbiology testing, including standard virology, bacterial culture, and search for mycoplasma; endotoxin detection using the limulus test; and cytogenetics. MSC potency was evaluated by determining the percentage inhibition of T-cell proliferation in Mixed-Lymphocyte Reaction (MLR) assay. Finally, MSC differentiation into adipocytes, osteocytes, and chondrocytes was validated in preliminary experiments [22].

MSC infusion

Third-party unrelated MSC infusion was performed on post-transplant day 3±2 through a

central intravenous line in fully monitored, stable, conscious and extubated patients who were receiving standard LT recipient care, after liver Doppler ultrasonography confirming arterial and portal flows. MSC infusion had to be performed within 60 min of thawing, with two investigators at the patients' bedside.

Primary endpoints

- MSC infusional toxicity: the duration and volume of the MSC infusion were noted. To assess pulmonary and systemic toxicity of MSC infusion, tympanic body temperature, heart rate, mean arterial blood pressure and peripheral capillary oxygen saturation (SpO₂) were recorded 5 min before infusion, after 15 minutes and at the end of the MSC infusion. Clinical signs of allergy, such as skin reaction or anaphylactic shock, were also recorded.
- MSC infectious and cancerous complications: the incidence, timing and severity of any infections (bacterial, viral, fungal) and any malignant diseases were prospectively recorded until month 12 in both groups.

Secondary endpoints

Patient and graft survivals and biopsy-proven graft rejection rates were prospectively recorded in both groups until month 12. Liver graft function (bilirubin, liver enzymes, international normalized ratio (INR)), kidney function (creatinine), C-reactive protein (CRP) and tacrolimus levels were compared using standard clinical blood tests at day 7 and months 1, 3, and 6. Blood immunoglobulin levels were compared at months 1 and 6.

Liver graft biopsy and immunohistochemistry

Month-6 formalin-fixed, paraffin-embedded graft biopsies were blindly analysed by two gastrointestinal pathologists (N.B., J.S), who described fibrosis and signs of graft rejection

according to the Banff criteria [24]. Paraffin-embedded sections of liver biopsy specimens (4 μ m thick) underwent immunostaining using an automated immunostainer (Ventana Medical Systems, Tucson, AZ) with antibodies directed against human CD3, CD4, CD8, CD20, CD138, CD68, CD1a and FoxP3. An amplification kit (Ventana Medical Systems) and a detection system including diaminobenzidine (Ventana Medical Systems) as a chromogen were used during the automated procedure. Archival lymph node sections were used as positive controls. For negative controls, the primary antibody was omitted. The mean number of positive cells in each patient was calculated by counting these cells (original magnification, x400) in the three most cellular microscopic fields, also called hot spots.

Peripheral blood lymphocyte immunophenotyping and CD4 phenotyping

Peripheral blood mononuclear cells were phenotyped on days 30, 90 and 180 using 4-color flow cytometry after treatment with a red blood cell lysing solution as described [25]. The analyzed cell subsets were T cells (CD3⁺), CD4⁺ T cells (CD3⁺CD4⁺lymphocytes), CD8⁺ T cells (CD3⁺CD8⁺ lymphocytes), naïve CD4⁺ T cells (CD4⁺CD45RA^{high} lymphocytes), memory CD4⁺ T cells (CD4⁺CD45RO⁺ lymphocytes), NK cells (CD3⁻CD56⁺ lymphocytes), as well as B cells (CD19⁺ lymphocytes). The percentage of positive cells was measured relative to total nucleated cells, after subtraction of nonspecific staining. Absolute counts were obtained by multiplying the percentages of positive cells by the white blood cell counts (Advia 120 hematology analyzer, Bayer Technicon).

More detailed CD4⁺ T cell phenotyping was performed on days 0 (before LT), 30 and 90 as previously reported [19]. Tregs were defined as CD4⁺CD25⁺CD127^{dim}FOXP3⁺ lymphocytes while remaining CD4⁺ T cells were considered as conventional T cells (Tconvs). Naïve Tregs were defined as CD45RA⁺HLA-DR^{neg} Tregs, and activated effector Tregs were defined as CD45RA^{neg}HLA-DR⁺ Tregs as previously reported [26]. T cell proliferation was assessed by

KI67 expression, and IL-2 signaling was estimated by quantifying the expression of phosphorylated STAT5 (phosphoSTAT5) [27]. The following antibodies specific for human epitopes were used: CD4-APC (RPA-T4), CD25-PeCy7 (BC96, Sony), CD127-biotin (eBioRDR5), CD45RA-BV510 (HI100,BD), HLA-DR-PE (L243), FOXP3-AlexaFluor488 (259D, Biolegend, ImTech Antwerp, Belgium), phosphoSTAT5-BV421 (pY694,BD), Ki67-PercPCy5.5 (B56, BD) and anti-streptavidin APCCy7 (all from eBioscience, unless otherwise indicated). Samples from patients were thawed and washed with staining buffer. One million cells of each sample were then incubated with surface antibodies for 20 min at 4°C in the dark and washed with staining buffer. This process was repeated for a 15-minute period for the streptavidin staining step. Then, samples were permeabilized using the PerFix EXPOSE (Beckman Coulter, Brea, CA) according to the manufacturer's instructions and as previously reported [26]. Data were acquired using a FACS Canto II (Becton Dickinson) and were analyzed with FlowJo v7.6.5 (Treestar Inc., San Carlos, CA).

Statistical analysis

Data are presented as median values and ranges, and the difference between groups was evaluated by the Mann-Whitney test. Proportions were analysed using Fischer's test. Differences between repeated measures were evaluated by one-way ANOVA using the Friedman test as a post-hoc test. Survival rates were calculated with the Kaplan-Meier method and compared with the log-rank (Mantel-Cox) test. A value of $P < 0.05$ was considered significant. Data were analysed using Prism 6.0c software for Macintosh OSX (GraphPad Software, San Diego, CA).

Results

Liver transplantation donor and recipient characteristics

No statistical difference could be detected between the MSC and control groups concerning the characteristics of both liver graft donors and recipients (supplementary material).

Primary endpoints

a) MSC infusional toxicity.

On day 3 (2-5), the 10 MSC patients received $2.1 \times 10^6/\text{kg}$ (1.9-2.7) MSC, representing a perfusion volume of 341 mL (302-614). Median duration of infusion was 25 min (11-60). No variation in vital parameters or cytokine release syndrome could be clinically detected during and after MSC infusion (Table 2). No MSC patient developed clinical signs of allergy or impairment of vital functions (including liver graft function) within the week following MSC infusion.

b) Infectious and cancerous complications.

No patient in either group developed life-threatening opportunistic infection or *de novo* cancer (including post-transplant lymphoproliferative disease) during follow-up. There was no difference in overall rates of infection between the two groups (Table 3). In the MSC group, two patients developed labial herpetic infections successfully treated by oral acyclovir. In addition, two MSC patients at high risk of CMV (D+, R-) developed asymptomatic CMV seroconversion under valganciclovir therapy. No patients developed CMV disease. Two patients transplanted for hepatocellular carcinoma (HCC) complicating cirrhosis had a pejorative pathology report and developed HCC recurrence: one MSC patient had a R1 LT with a HCC nodule invading the diaphragm (he died from HCC recurrence at month 10) and one control patient had an unsuspected neoplastic thrombus in a supra-hepatic vein at liver pathology (still alive at 5-year follow-up after HCC recurrence at month 23 and resection of pulmonary metastases).

Secondary Endpoints

No patient required retransplantation during the first year of follow-up. One patient from the control group died at day 16 from a hypovolemic shock induced by a fistula between the hepatic artery and the bile duct, probably due to an infected pseudoaneurysm. Six-month graft and patient survivals were 100% and 90% in the MSC and control group, respectively (NS). One-year graft and patient survivals were 90% in both groups (NS). No patient in either group developed biopsy-proven rejection during the first 6 months of follow-up. Protocol month-6 biopsies did not demonstrate a difference between groups in the evaluation of the Banff criteria, the fibrosis score or the immunohistochemistry (Table 4; Fig S1 and Fig S2 in supplementary material). No difference could be detected in liver graft or kidney function between the two groups during the 6 months of comparison (Table 5; Fig S3 in supplementary material). No difference in peripheral blood lymphocyte phenotyping could be detected on day 30, 90 and 180 (Table 6; Fig S4 in supplementary material).

Impact of MSCs on peripheral blood CD4⁺ T cells (including Tregs)

The 2 groups of patients had similar counts of peripheral blood CD4⁺ T cells and Tconvs on days 0, 30 and 90 after transplantation (Fig 2 A-B). As shown in Fig 2 C-F, Treg counts and phenotype (naïve versus resting versus activated) were comparable in the 2 groups of patients at each time point. Furthermore, Treg as well as Tconv proliferation (assessed by KI67 expression) was also similar in the 2 groups of patients, as were the levels of phosphoSTAT5 in Tregs (the latter translating similar IL-2 signalling in Tregs). These combined observations suggest that a single MSC infusion had no impact on Treg count or phenotype in this study.

Immunosuppression withdrawal in the MSC group

One patient from the MSC group was excluded from immunosuppression withdrawal attempt due to HCC recurrence, but the nine others met the necessary criteria. In one patient, tacrolimus and MMF withdrawal was performed without rejection and she remained off immunosuppression for 12 months. In two patients, MMF monotherapy was achieved at month 9, but graft rejection occurred during MMF withdrawal and was successfully treated by tacrolimus reintroduction. In 6 patients, the transaminases significantly increased during tacrolimus withdrawal. In these cases, withdrawal was cancelled and liver tests normalised after increase of the tacrolimus dose.

Discussion

This phase 1, prospective, controlled study is the first to evaluate the feasibility, safety and tolerability of MSC infusion in a series of 10 LT patients under classical tacrolimus-based immunosuppression. In these patients, a post-transplantation intravenous $1.5\text{--}3 \times 10^6/\text{kg}$ MSC infusion was well tolerated, without evidence of pulmonary dysfunction or of cytokine-release syndrome. This dosing was chosen according to the authors' experiences with MSC infusion after HSC transplantation [23, 28]. These LT patients receiving MSC did not develop any evidence of impairment in vital organ functions, including the liver graft and the kidneys. In addition, they did not suffer from an increased susceptibility to infections. No *de novo* cancer was detected after one year of follow-up, and a HCC recurrence was observed in a patient with a very poor prognosis due to unexpected extra-hepatic HCC spread discovered during LT. For all these primary endpoints, the LT recipients who received MSCs did not react differently compared to patients in the control group. This finding is an important step in the evaluation of the potential role of MSCs in SOT recipients, and particularly after LT.

In the last decade, MSCs have been extensively studied both *in vitro* and *in vivo*. Their anti-inflammatory and immunoregulatory properties [29] [3], added to potential beneficial effects on ischemia/reperfusion injury [5], might select MSCs as a potential future therapy for SOT

recipients in whom life-long immunosuppression and chronic allograft dysfunction still impair quality of life and graft survival. However, as the clinical use of MSCs is still under evaluation in preliminary trials in non-transplant patients, their potential secondary effects need to be carefully assessed in SOT recipients. Due to their size, MSCs are known to embolize within the pulmonary circulatory bed when they are infused in the peripheral or central venous circulation of mice [12]. There is therefore a theoretical risk of decreased pulmonary exchange after MSC infusion, but this complication has not been reported so far in the early phase clinical trials nor in the randomized study in living-related kidney transplantation performed in China [9]. As reported previously by our group, MSC infusion in hematopoietic stem cell transplant recipients has not been associated with any infusional toxicity [23], nor with long-term impairment of lung function [28]. This was confirmed in the current trial, as our 10 patients receiving MSCs did not develop any sign of pulmonary dysfunction. In addition, this study did not observe any suspicion of allergy or cytokine-release syndrome, or of any other possible complications concerning the liver graft or extra-hepatic organ function. In a preliminary evaluation in 2 kidney recipients, possible toxicity of MSC infusion on kidney graft function was suggested [14], but this “engraftment syndrome” was not detected in our cohort of LT recipients or in any other MSC clinical trial to date.

As MSCs are immunosuppressive, SOT recipients who receive MSCs in addition to standard immunosuppression could be over-immunosuppressed and develop higher rates of opportunistic infections [26]. Again, in a small series of kidney recipients, opportunistic infections were observed after MSC treatment [15]. On the contrary, in the largest experience reported so far of MSC infusion after living-related kidney transplantation, MSC recipients developed fewer infectious complications than controls [9]. In our series, the MSC patients did not develop life-threatening infections, and no difference could be detected in comparison with the control group.

It has been suggested in *in vitro* experiments that MSCs might carry a potential for cancerous degeneration [17]. This potential risk has so far not been demonstrated in the preliminary MSC clinical experiences in either SOT or in non-SOT patients, and no patient in our series had developed *de novo* cancer after one-year follow-up. This important issue needs to be confirmed by further follow-ups of this series and by further experience in larger series. Furthermore, in the series described here, one patient died from early HCC recurrence after a R1 LT with a very bad prognosis. The authors do not consider that HCC within Milan criteria should be excluded for further MSC trials in LT, but the possibility of an increased risk of HCC recurrence after MSC infusion cannot be excluded by this preliminary phase 1 study.

As secondary end-points, this study prospectively evaluated the possible effects of a single infusion of MSCs on LT recipient immunity by comparison with a control group. No difference could be detected between the MSC and control groups on graft rejection episodes, opportunistic infection rates, graft histology and immunohistochemistry at day 180 and on peripheral blood CD4+ T-cell subsets. Particularly, no impact of MSC infusion on Tconv counts/proliferation was demonstrated, suggesting that MSCs did not impact T-cell immunity while, in contrast to what has been observed in mice [30], MSC infusion did not influence Treg number, proliferation or phenotype in this cohort of patients. This finding might indicate that a single infusion of MSCs in LT patients receiving tacrolimus and MMF will not modify their immunity status. As MSCs and immunosuppressive drugs inhibit the same targets (essentially T cells), it is reasonable to consider that interactions between them can occur. The current standard of immunosuppression after LT is a triple therapy associating low-dose steroids, MMF and tacrolimus, with rapid steroid weaning. *In vitro*, some authors have shown that tacrolimus and rapamycin might decrease MSC immunosuppressive properties [31], and conversely, that MSCs might reduce the immunosuppressive capacities of tacrolimus and rapamycin. Such an effect has not been found with mycophenolic acid (MPA), an MMF

metabolite. Moreover, a high dose of tacrolimus seems to be toxic for MSCs, while MPA and rapamycin at a therapeutic dose just inhibit MSC proliferation [32]. Furthermore, it has been demonstrated that MPA and MSCs have a synergistic immunosuppressive effect [32]. *In vivo*, MPA and MSCs also synergize to promote long-term allograft tolerance in rat heart transplantation [33]. As Tregs probably play an important role in MSC-mediated immunomodulatory effects, it is important to confirm that such a combination therapy is also favorable for Treg expansion. Hence, a recent study supported that mTOR inhibitor-based immunosuppression favours survival of Tregs after administration in a nonhuman primate model, whereas tacrolimus does not [34].

In addition, in a phase 2 part of this study, patients from the MSC group underwent unsuccessful progressive immunosuppression weaning. Induction of operational tolerance is a major goal in SOT and particularly in LT patients [1]. Operational tolerance is a rare phenomenon after LT [18]. Tregs have been proposed to be key in strategies aiming for tolerance and immunomodulation after SOT [35]. In a recent paper, Todo et al. demonstrated that a single Treg injection might promote operational tolerance after living-related LT [36]. Recently, it has been demonstrated both *in vitro* and *in vivo* that MSCs could promote Treg expansion by their effects on immature dendritic cells [37]. In this study, the authors were not able to show that a single infusion of MSCs at day 3 after deceased LT could promote Treg expansion, Treg infiltration of the liver graft at biopsy at day 180, or operational tolerance.

There are many shortcomings to this study. First, it is clear that this first study in 10 LT recipients does not prove the safety of MSC infusion in this setting. These results will have to be confirmed by further studies in larger groups of SOT recipients. The absence of detectable effects of MSCs might be due to an insufficient sample size, to the tacrolimus-based immunosuppressive regimen or to an insufficient MSC dosing, which should possibly be increased or repeated. The timing (pre-, intra- or post-operative) and the infusion routes

(peripheral vein, portal vein or hepatic artery) of MSC infusion should also be evaluated. Different sources (BM, fat tissue, liver) or donors (organ donor, organ recipient) of MSCs might also be tested in further studies.

In summary, this study reported the first prospective controlled phase 1 clinical trial evaluating the toxicity of a MSC-based immune-regulating regimen in a series of deceased LT recipients receiving classical tacrolimus-based immunosuppression. In this trial, no side effects of MSC infusion at day 3 after transplantation could be detected. Even if no modification of the patient immunity and Treg expansion could be demonstrated, and even if immunosuppression weaning was not successful in this first series of 10 patients, this study opens the way for further MSC or Treg-based trials in LT recipients.

LEGENDS OF FIGURES AND TABLES

Figure 1: Scheme of the study

MSC: Mesenchymal stromal cell; MMF: mycophenolate mofetil; MP: Methylprednisolone

Figure 2: Evolution of on peripheral blood CD4+ T cells (including Tregs) on days 0, 30 and 90 in the control (white rectangles) or MSC (grey rectangles) groups

A. Total CD4+ T cells,

B. Conventional CD4+ T cells (Tconv)

C. Regulatory T cells (Tregs)

D. Treg subsets

E. Treg proliferation (assessed by KI67 expression)

F. Treg IL-2 signaling (assessed through phosphoSTAT5 expression)

HLA-DR^{pos}CD45RA^{neg} (DR+RA-) Tregs refer to activated Tregs, HLA-DR^{neg}CD45RA^{neg} (DR+RA-) Tregs refer to resting Tregs and HLA-DR^{neg}CD45RA^{pos} (DR-RA+) Tregs refer to naive Tregs. Plots display the median, 25th and 75th percentiles of the distribution (boxes), and whiskers extend to the 10th and 90th percentiles. Grey zones show normal ranges (from 5th to 95th percentiles) and horizontal lines the medians in 45 age-matched healthy controls. No statistical difference could be detected between the two. Mann-Whitney test

Table 1: LT recipient inclusion and exclusion criteria

LT: Liver transplantation; DBD: donation after brain death; DCD: donation after circulatory death; MSC: Mesenchymal Stromal Cells; HIV: Human Immunodeficiency virus; HCV: Hepatitis C virus; EBV: Epstein-Barr virus

Table 2: Comparison of vital parameters before, during and after MSC infusion

MSC: Mesenchymal stromal cells; SpO₂: peripheral capillary oxygen saturation
One-way ANOVA using Friedman test as a post-hoc test

Table 3: Cancerous and infectious complications (1-year follow-up)

MSC: Mesenchymal stromal cell; HCC: hepatocellular carcinoma; CMV: cytomegalovirus; HSV: herpes simplex virus; VZV: Varicella-zoster virus
Fischer's test

Table 4: Histology and immunohistochemistry of D180 liver graft biopsies

MSC: mesenchymal stromal cell
Data are presented as median and ranges; Mann-Whitney test

Table 5: Postoperative laboratory tests

MSC: mesenchymal stromal cells; D: day; AST: aspartate aminotransferase; Alkaline Ph : alkaline phosphatase ; GGT: gamma glutamil transferase; INR: international normalized ratio
CRP: C reactive protein
Data are presented as median and ranges; Mann-Whitney test

Table 6: Peripheral blood lymphocyte counts

Data are presented as median and ranges; Mann-Whitney test

References

- [1] Adams DH, Sanchez-Fueyo A, Samuel D. From immunosuppression to tolerance. *J Hepatol* 2015;62:S170-185.
- [2] Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 2006;8:315-317.
- [3] Crop M, Baan C, Weimar W, Hoogduijn M. Potential of mesenchymal stem cells as immune therapy in solid-organ transplantation. *Transpl Int* 2009;22:365-376.
- [4] Erpicum P, Detry O, Weekers L, Bonvoisin C, Lechanteur C, Briquet A, et al. Mesenchymal stromal cell therapy in conditions of renal ischaemia/reperfusion. *Nephrol Dial Transplant* 2014;29:1487-1493.
- [5] Rowart P, Erpicum P, Detry O, Weekers L, Gregoire C, Lechanteur C, et al. Mesenchymal Stromal Cell Therapy in Ischemia/Reperfusion Injury. *J Immunol Res* 2015;2015:602597.
- [6] Casiraghi F, Perico N, Remuzzi G. Mesenchymal stromal cells to promote solid organ transplantation tolerance. *Curr Opin Organ Transplant* 2013;18:51-58.
- [7] Franquesa M, Hoogduijn MJ, Reinders ME, Eggenhofer E, Engela AU, Mensah FK, et al. Mesenchymal Stem Cells in Solid Organ Transplantation (MiSOT) Fourth Meeting: lessons learned from first clinical trials. *Transplantation* 2013;96:234-238.
- [8] English K, Wood KJ. Mesenchymal stromal cells in transplantation rejection and tolerance. *Cold Spring Harb Perspect Med* 2013;3:a015560.
- [9] Tan J, Wu W, Xu X, Liao L, Zheng F, Messinger S, et al. Induction therapy with autologous mesenchymal stem cells in living-related kidney transplants: a randomized controlled trial. *JAMA* 2012;307:1169-1177.

- [10] Vandermeulen M, Gregoire C, Briquet A, Lechanteur C, Beguin Y, Detry O. Rationale for the potential use of mesenchymal stromal cells in liver transplantation. *World J Gastroenterol* 2014;20:16418-16432.
- [11] Lalu MM, McIntyre L, Pugliese C, Fergusson D, Winston BW, Marshall JC, et al. Safety of cell therapy with mesenchymal stromal cells (SafeCell): a systematic review and meta-analysis of clinical trials. *PLoS One* 2012;7:e47559.
- [12] Furlani D, Ugurlucan M, Ong L, Bieback K, Pittermann E, Westien I, et al. Is the intravascular administration of mesenchymal stem cells safe? Mesenchymal stem cells and intravital microscopy. *Microvasc Res* 2009;77:370-376.
- [13] Dillmann J, Popp FC, Fillenberg B, Zeman F, Eggenhofer E, Farkas S, et al. Treatment-emergent adverse events after infusion of adherent stem cells: the MiSOT-I score for solid organ transplantation. *Trials* 2012;13:211.
- [14] **Perico N, Casiraghi F, Introna M**, Gotti E, Todeschini M, Cavinato RA, et al. Autologous mesenchymal stromal cells and kidney transplantation: a pilot study of safety and clinical feasibility. *Clin J Am Soc Nephrol* 2011;6:412-422.
- [15] Reinders ME, de Fijter JW, Roelofs H, Bajema IM, de Vries DK, Schaapherder AF, et al. Autologous bone marrow-derived mesenchymal stromal cells for the treatment of allograft rejection after renal transplantation: results of a phase I study. *Stem Cells Trans Med* 2013;2:107-111.
- [16] Baerschiger RM, Serre-Beinier V, Morel P, Bosco D, Peyrou M, Clement S, et al. Fibrogenic potential of human multipotent mesenchymal stromal cells in injured liver. *PLoS One* 2009;4:e6657.
- [17] Haarer J, Johnson CL, Soeder Y, Dahlke MH. Caveats of mesenchymal stem cell therapy in solid organ transplantation. *Transpl Int* 2015;28:1-9.

- [18] Barkholt L, Flory E, Jekerle V, Lucas-Samuel S, Ahnert P, Bisset L, et al. Risk of tumorigenicity in mesenchymal stromal cell-based therapies--bridging scientific observations and regulatory viewpoints. *Cytotherapy* 2013;15:753-759.
- [19] **Braat AE, Blok JJ**, Putter H, Adam R, Burroughs AK, Rahmel AO, et al. The Eurotransplant donor risk index in liver transplantation: ET-DRI. *Am J Transplant* 2012;12:2789-2796.
- [20] Detry O, Deroover A, Meurisse N, Hans MF, Delwaide J, Lauwick S, et al. Donor age as a risk factor in donation after circulatory death liver transplantation in a controlled withdrawal protocol programme. *Br J Surg* 2014;101:784-792.
- [21] Belghiti J, Noun R, Sauvanet A. Temporary portocaval anastomosis with preservation of caval flow during orthotopic liver transplantation. *Am J Surg* 1995;169:277-279.
- [22] Lechanteur C, Briquet A, Giet O, Delloye O, Baudoux E, Beguin Y. Clinical-scale expansion of mesenchymal stromal cells: a large banking experience. *J Transl Med* 2016;14:145.
- [23] **Baron F, Lechanteur C**, Willems E, Bruck F, Baudoux E, Seidel L, et al. Cotransplantation of mesenchymal stem cells might prevent death from graft-versus-host disease (GVHD) without abrogating graft-versus-tumor effects after HLA-mismatched allogeneic transplantation following nonmyeloablative conditioning. *Biol Blood Marrow Transplant* 2010;16:838-847.
- [24] Banff schema for grading liver allograft rejection: an international consensus document. *Hepatology* 1997;25:658-663.
- [25] Hannon M, Beguin Y, Ehx G, Servais S, Seidel L, Graux C, et al. Immune Recovery after Allogeneic Hematopoietic Stem Cell Transplantation Following Flu-TBI versus TLI-ATG Conditioning. *Clin Cancer Res* 2015;21:3131-3139.

- [26] Balan A, Lucchini G, Schmidt S, Schneider A, Tramsen L, Kuci S, et al. Mesenchymal stromal cells in the antimicrobial host response of hematopoietic stem cell recipients with graft-versus-host disease--friends or foes? *Leukemia* 2014;28:1941-1948.
- [27] **Ehx G, Hannon M**, Beguin Y, Humblet-Baron S, Baron F. Validation of a multicolor staining to monitor phosphoSTAT5 levels in regulatory T-cell subsets. *Oncotarget* 2015;6:43255-43266.
- [28] Moermans C, Lechanteur C, Baudoux E, Giet O, Henket M, Seidel L, et al. Impact of cotransplantation of mesenchymal stem cells on lung function after unrelated allogeneic hematopoietic stem cell transplantation following non-myeloablative conditioning. *Transplantation* 2014;98:348-353.
- [29] Eggenhofer E, Luk F, Dahlke MH, Hoogduijn MJ. The life and fate of mesenchymal stem cells. *Front Immunol* 2014;5:148.
- [30] Casiraghi F, Azzollini N, Cassis P, Imberti B, Morigi M, Cugini D, et al. Pretransplant infusion of mesenchymal stem cells prolongs the survival of a semiallogeneic heart transplant through the generation of regulatory T cells. *J Immunol* 2008;181:3933-3946.
- [31] Buron F, Perrin H, Malcus C, Hequet O, Thaumat O, Kholopp-Sarda MN, et al. Human mesenchymal stem cells and immunosuppressive drug interactions in allogeneic responses: an in vitro study using human cells. *Transplant Proc* 2009;41:3347-3352.
- [32] Hoogduijn MJ, Crop MJ, Korevaar SS, Peeters AM, Eijken M, Maat LP, et al. Susceptibility of human mesenchymal stem cells to tacrolimus, mycophenolic acid, and rapamycin. *Transplantation* 2008;86:1283-1291.
- [33] **Popp FC, Eggenhofer E**, Renner P, Slowik P, Lang SA, Kaspar H, et al. Mesenchymal stem cells can induce long-term acceptance of solid organ allografts in synergy with low-dose mycophenolate. *Transpl Immunol* 2008;20:55-60.

- [34] Singh K, Stempora L, Harvey RD, Kirk AD, Larsen CP, Blazar BR, et al. Superiority of rapamycin over tacrolimus in preserving nonhuman primate Treg half-life and phenotype after adoptive transfer. *Am J Transplant* 2014;14:2691-2703.
- [35] **Taubert R, Danger R**, Londono MC, Christakoudi S, Martinez-Picola M, Rimola A, et al. Hepatic Infiltrates in Operational Tolerant Patients After Liver Transplantation Show Enrichment of Regulatory T Cells Before Proinflammatory Genes Are Downregulated. *Am J Transplant* 2016;16:1285-1293.
- [36] **Todo S, Yamashita K**, Goto R, Zaitzu M, Nagatsu A, Oura T, et al. A Pilot Study of Operational Tolerance with a Regulatory T Cell-Based Cell Therapy in Living Donor Liver Transplantation. *Hepatology* 2016;64:632-643.
- [37] Choi YS, Jeong JA, Lim DS. Mesenchymal stem cell-mediated immature dendritic cells induce regulatory T cell-based immunosuppressive effect. *Immunol Invest* 2012;41:214-229.

Author names in bold designate shared co-first authorship

Table 1: LT recipient inclusion and exclusion criteria

Inclusion criteria	
Surgery	First whole liver deceased LT (DBD or DCD)
Age	Between 18 and 75 years
Graft	Functioning graft at time of MSC infusion
	Graft doppler ultrasonography confirming arterial and portal flows
Exclusion criteria	
Surgery	Re LT, partial LT, combined LT
Cancer	Past history of cancer in the donor or recipient, with the exception of hepatocarcinoma within Milan criteria
Infection	Active infection in the donor or recipient, including HIV and HCV
	EBV negative (recipient)
Miscellaneous	Auto-immune liver disease (recipient)
	Endotracheal intubation (recipient)
	Severe postoperative complications (recipient)

LT: Liver transplantation; DBD: donation after brain death; DCD: donation after circulatory death; MSC: Mesenchymal Stromal Cells; HIV: Human Immunodeficiency virus; HCV: Hepatitis C virus; EBV: Epstein-Barr virus

Table 2: Comparison of vital parameters before, during and after MSC infusion

	Pre MSC infusion	After 15 min	End of MSC infusion	P
Temperature (°C)	36.1 (35.4-37.7)	36.4 (35-36.9)	36.2 (35.5-37)	0.87
Mean arterial pressure (mmHg)	103 (87-124)	107 (84-120)	106 (94-115)	0.83
Heart rate (per min)	81 (65-102)	83 (65-102)	81 (68-101)	0.17
SpO ₂ (%)	99 (93-100)	100 (92-100)	98 (93-100)	0.67

MSC: Mesenchymal stromal cells; SpO₂: peripheral capillary oxygen saturation

ANOVA using the Friedman test as a post-hoc test

Table 3: Cancerous and infectious complications (1-year follow-up)

			MSC group (n=10)	Control group (n=9)	P
Cancer					
	Overall		1	0	> 0.99
	De novo		0	0	
	HCC recurrence		1	0	
Infection					
	Overall		2	6	0.06
	Fungal		0	0	
	Viral	CMV disease	0	0	
		HSV	2	0	
		VZV	0	1	
	Other	Wound	0	1	
		Urinary	0	2	
		Sinusitis	0	1	
		Pulmonary	0	1	

MSC: Mesenchymal stromal cell; HCC: hepatocellular carcinoma; CMV: cytomegalovirus;

HSV: herpes simplex virus; VZV: Varicella-zoster virus

Fischer's test

Table 4: Histology and immunohistochemistry of D180 liver graft biopsies

	MSC group (n=10)	Control group (n=9)	P
Banff score	3 (0-6)	4 (0-7)	0.21
Fibrosis score	1 (0-2)	1 (0-3)	0.48
CD3	196 (95-334)	162 (93-590)	0.86
CD4	101 (54-212)	103 (17-496)	> 0.99
CD8	69 (15-196)	85 (12-300)	0.49
CD68	28.5 (12-75)	40 (15-104)	0.58
CD1a	1 (0-3)	1 (0-3)	0.83
CD138	7.5 (4-38)	6 (2-44)	0.50
CD20	27 (3-95)	28 (10-163)	0.66
FoxP3	2 (0-16)	4 (0-33)	0.49

MSC: mesenchymal stromal cell

Data are presented as median and ranges; Mann-Whitney test

Table 5: Postoperative laboratory tests

	MSC group (n=10)	Control group (n=9)	P
D7			
total bilirubin (mg/L)	10.2 (4.6-26.8)	8.3 (3.7-20.7)	0.21
AST (U/L)	28.5 (19-101)	46 (30-105)	0.16
Alcaline Ph (U/L)	140 (43-475)	256 (172-590)	0.04
GGT (U/L)	218 (29-626)	368 (172-760)	0.24
INR	1.14 (1-1.21)	1.06 (1-1.26)	0.16
creatinine (mg/L)	11.55 (5.7-36)	8.9 (5.9 – 16.9)	0.32
CRP (mg/L)	32.8 (8.4-50.1)	24.6 (12.8-144.3)	0.82
tacrolimus (µg/L)	7.1 (3.1-9)	9 (2.1-11.7)	0.12
D30			
total bilirubin (mg/L)	5.6 (3.4-11.6)	4.6 (1.3-7.5)	0.34
AST (U/L)	18 (11-51)	16 (9-61)	0.48
Alcaline Ph (U/L)	137.5 (53-554)	144 (103-857)	0.43
GGT (U/L)	101 (26-596)	112 (42-690)	0.82
INR	1.15 (0.97-1.26)	1.08 (1-1.19)	0.53
creatinine (mg/L)	16.2 (5.3-24.4)	14.1 (8.2-27.6)	0.45
CRP (mg/L)	12.9 (4.8-62.2)	17.2 (3.5-73)	0.94
tacrolimus (µg/L)	8.1 (2.4-10)	8 (5-16.3)	0.51
D90			
total bilirubin (mg/L)	4.8 (3-19.8)	4.3 (2.3-7.5)	0.34
AST (U/L)	20 (14-31)	20 (11-58)	0.79
Alcaline Ph (U/L)	101.5 (56-1461)	119 (86-570)	0.54
GGT (U/L)	58.5 (15-695)	49 (14-332)	0.76
INR	1.1 (0.95-1.29)	1.13 (1.01-1.56)	0.65
creatinine (mg/L)	12.05 (5-25.7)	13.4 (7-21.7)	0.92
CRP (mg/L)	3.1 (1-27.6)	6.8 (1.3-23.5)	0.20
tacrolimus (µg/L)	7.7 (3.7-13)	6.4 (5.2-13.2)	0.61
D180			
total bilirubin (mg/L)	6.6 (3.7-25.7)	4.6 (0.43-27)	0.27
AST (U/L)	25 (15-44)	24 (14-136)	0.64
Alcaline Ph (U/L)	143.5 (67-1,166)	186 (82-554)	0.26
GGT (U/L)	81 (22-978)	53 (12-2,064)	0.43
INR	1.1 (1-1.26)	1.07 (1-1.17)	0.23
creatinine (mg/L)	11.6 (7.1-18.9)	10.1 (1.28-15.8)	0.30
CRP (mg/L)	3.5 (0.7-36.5)	5.6 (0.9-151)	0.23
tacrolimus (µg/L)	4.9 (2.3-9.3)	7.4 (4.9-13)	0.02

MSC: mesenchymal stromal cells; D: day; AST: aspartate aminotransferase; Alkaline Ph : alkaline phosphatase ; GGT: gamma glutamil transferase; INR: international normalized ratio
CRP: C reactive protein

Data are presented as median and ranges; Mann-Whitney test

Table 6: Peripheral blood lymphocyte counts

	MSC group (n=10)	Control group (n=9)	P
D30			
White blood cells (/μL)	6,630 (3,280-9,700)	5,190 (4,150-10,030)	0.67
Lymphocytes (/μL)	855 (380-1,690)	940 (300-1,550)	0.92
CD3 (/μL)	687 (288-1,406)	620 (200-1,336)	0.48
CD45RA (/μL)	119 (50-557)	147 (48-234)	0.70
CD45RO (/μL)	373 (179-516)	201 (79 -609)	0.23
CD3+CD4+ (/μL)	535 (230-978)	349 (128-786)	0.30
CD3+CD56+ (/μL)	27 (1-87)	42 (4-154)	0.35
CD3+CD8+ (/μL)	115 (49-418)	142 (57-336)	0.76
CD19 (/μL)	144 (30-286)	99 (38-369)	0.70
CD56 (/μL)	109 (45-365)	188 (58-618)	0.27
D90			
White blood cells (/μL)	5,265 (970-8,160)	5,200 (2,470-7,030)	0.39
Lymphocytes (/μL)	875 (420-1,880)	760 (490-1,760)	0.82
CD3 (/μL)	767 (352-1,225)	553 (274-1,419)	0.30
CD45RA (/μL)	123 (51-389)	82 (54-259)	0.58
CD45RO (/μL)	381 (171-680)	179 (135-765)	0.23
CD3+CD4+ (/μL)	516 (292-923)	285 (202-976)	0.27
CD3+CD56+ (/μL)	21 (1-99)	34 (2-197)	0.76
CD3+CD8+ (/μL)	202 (41-496)	228 (56-362)	0.94
CD19 (/μL)	93 (34-354)	100 (21-321)	0.76
CD56 (/μL)	154 (66-331)	119 (59-550)	0.82
D180			
White blood cells (/μL)	4,815 (4,200-8,150)	5,440 (2,680-11,430)	0.99
Lymphocytes (/μL)	1,250 (660-2,260)	1,000 (540-1,340)	0.23
CD3 (/μL)	880 (395-2,098)	592 (342-1,366)	0.27
CD45RA (/μL)	127 (76-364)	108 (61-298)	0.58
CD45RO (/μL)	396 (214-615)	267 (156-864)	0.20
CD3+CD4+ (/μL)	623 (348-728)	359 (224-1,163)	0.20
CD3+CD56+ (/μL)	31 (1-91)	36 (3-117)	0.54
CD3+CD8+ (/μL)	238 (38-1,471)	210 (73-345)	0.70

CD19 (/μL)	99 (25-256)	192 (52-258)	0.27
CD56 (/μL)	191 (66-386)	210 (55-490)	>0.99

Data are presented as median and ranges; Mann-Whitney test

