

COMMISSION 27 OF THE I. A. U.  
INFORMATION BULLETIN ON VARIABLE STARS

Number 2625

Konkoly Observatory  
Budapest  
13 November 1984  
HU ISSN 0374 - 0676

ON THE  $\delta$  WEAK VARIABLE HD 28843

Photometric observations of HD 28843 (= HR 1441) in the  $uvby$  system have led to the derivation of a period of  $1.37390 \pm 0.00015$  d (Manfroid and Mathys, 1984). The observations covered a time span of less than two years, from January 1976 to November 1977.

Contrarily to what was stated by Manfroid and Mathys, the star was already known as a variable as early as 1966 (Cousins, 1966). In fact, eight observations in the  $V$  band are available. These  $V$  data can be used to refine the period determination.  $V$  and  $y$  magnitude can be compared directly. Several frequency peaks are obtained when performing the analysis of the complete set of data, both by a Fourier-Deeming technique and by the method of minimization of the standard deviation of the observations from a least-square fit by a sine wave and its first harmonic (Mathys and Manfroid, 1984). Unlike the former, the latter method allows in the present case to determine unambiguously the best value of the frequency ( $f = 0.727901 \pm 0.000006$  d $^{-1}$ ; the relevant part of the periodogram is shown on Fig. 1).




FIG. 1 — Part of the periodogram of HD 28843 showing the most probable values of the frequency




FIG. 2 — Phase diagram for HD 28843 with the period  $P = 1.373813$  d. Phase origin is JD 2,437,847.336. The solid curve is a least-square fit of the observations by a sine wave and its first harmonic. Z: AWJC's 1966 V data. Y: Pedersen and Thomsen's y observations, January–February 1976 (Pedersen and Thomsen, 1977). X: JM and GM's y measurements, November 1977

A study of the phase diagrams confirms that the alternative values 0.726347 and  $0.729453 \text{ d}^{-1}$  are much less good. Hence we adopt

$$P = 1.373813 \pm 0.000012 \text{ d}$$

as the most probable value of the period of HD 28843. The phase diagram obtained with that period is shown on Fig. 2, together with a least-square fit of the data by a sine wave and its first harmonic.

J. MANFROID  
Institut d'Astrophysique  
5, avenue de Cointe  
B-4200 Cointe-Ougrée  
Belgium

G. MATHYS  
Institut für Astronomie  
ETH-Zentrum  
CH-8092 Zurich  
Switzerland

A.W.J. COUSINS  
South African Astronomical  
Observatory  
PO Box 9  
Observatory, 7935  
South Africa

#### References:

Cousins, A.W.J. and Stoy, R.H., 1966, *R. Obs. Bull.* No. 121.  
 Manfroid, J. and Mathys, G., 1984, *Inf. Bull. Var. Stars* 2551.  
 Mathys, G. and Manfroid, J., 1984, submitted to *Astron. Astrophys. Suppl. Ser.*  
 Pedersen, H. and Thomsen, B., 1977, *Astron. Astrophys. Suppl. Ser.* 30, 11.