



A way of extending Pascal and Sierpiński triangles to finite words

Joint work with Julien Leroy and Michel Rigo (ULiège)

Manon Stipulanti (ULiège)

FRIA grantee

Young Mathematicians Symposium of the Greater Region
Nancy (France)
September 24-25, 2018

Classical Pascal triangle

$\binom{m}{k}$	k								
	0	1	2	3	4	5	6	7	...
0	1	0	0	0	0	0	0	0	
1	1	1	0	0	0	0	0	0	
2	1	2	1	0	0	0	0	0	
m	3	1	3	3	1	0	0	0	0
	4	1	4	6	4	1	0	0	0
	5	1	5	10	10	5	1	0	0
	6	1	6	15	20	15	6	1	0
	7	1	7	21	35	35	21	7	1
	\vdots								\ddots

Usual binomial coefficients
of integers:


$$\binom{m}{k} = \frac{m!}{(m-k)!k!}$$

Pascal's rule:

$$\binom{m}{k} = \binom{m-1}{k} + \binom{m-1}{k-1}$$

A specific construction

- Grid: first 2^n rows and columns



- Color the grid:
Color the first 2^n rows and columns of the Pascal triangle

$$\left(\binom{m}{k} \bmod 2 \right)_{0 \leq m, k < 2^n}$$

in

- white if $\binom{m}{k} \equiv 0 \pmod{2}$
- black if $\binom{m}{k} \equiv 1 \pmod{2}$

- Color the grid:
Color the first 2^n rows and columns of the Pascal triangle

$$\left(\binom{m}{k} \bmod 2 \right)_{0 \leq m, k < 2^n}$$

in


- white if $\binom{m}{k} \equiv 0 \pmod{2}$
- black if $\binom{m}{k} \equiv 1 \pmod{2}$
- Normalize by a homothety of ratio $1/2^n$
(bring into $[0, 1] \times [0, 1]$)
 \rightsquigarrow sequence belonging to $[0, 1] \times [0, 1]$

What happens for $n \in \{0, 1\}$

$$n = 0$$



What happens for $n \in \{0, 1\}$

$$n = 0$$






What happens for $n \in \{0, 1\}$

$n = 0$






What happens for $n \in \{0, 1\}$

$n = 0$



What happens for $n \in \{0, 1\}$




$n = 0$




$n = 1$

What happens for $n \in \{0, 1\}$

$n = 0$






$n = 1$





What happens for $n \in \{0, 1\}$

$n = 0$






$n = 1$






What happens for $n \in \{0, 1\}$







$n = 0$




$n = 1$



The first six elements of the sequence



The tenth element of the sequence




The Sierpiński gasket



The Sierpiński gasket



The Sierpiński gasket



Folklore fact

The latter sequence converges to the Sierpiński gasket when n tends to infinity (for the Hausdorff distance).

Folklore fact

The latter sequence converges to the Sierpiński gasket when n tends to infinity (for the Hausdorff distance).

Definitions:

- ϵ -fattening of a subset $S \subset \mathbb{R}^2$

$$[S]_\epsilon = \bigcup_{x \in S} B(x, \epsilon)$$

- $(\mathcal{H}(\mathbb{R}^2), d_h)$ complete space of the non-empty compact subsets of \mathbb{R}^2 equipped with the *Hausdorff distance* d_h

$$d_h(S, S') = \min\{\epsilon \in \mathbb{R}_{\geq 0} \mid S \subset [S']_\epsilon \quad \text{and} \quad S' \subset [S]_\epsilon\}$$

Replace usual binomial coefficients of integers by
binomial coefficients of **finite words**

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = 101001$ $v = 101$

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = \mathbf{101}001$ $v = 101$ 1 occurrence

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = \mathbf{101001}$ $v = 101$ 2 occurrences

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = 101001$ $v = 101$ 3 occurrences

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = 101001$ $v = 101$ 4 occurrences

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = 101001$ $v = 101$ 5 occurrences

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = 101001$ $v = 101$ 6 occurrences

Definition: A *finite word* is a finite sequence of letters belonging to a finite set called *alphabet*.

Example: $101, 101001 \in \{0, 1\}^*$

Binomial coefficient of words

Let u, v be two finite words.

The *binomial coefficient* $\binom{u}{v}$ of u and v is the number of times v occurs as a subsequence of u (meaning as a “scattered” subword, non-consecutive letters).

Example: $u = 101001$ $v = 101$

$$\Rightarrow \binom{101001}{101} = 6$$

Remark:

Natural generalization of binomial coefficients of integers

With a one-letter alphabet $\{a\}$

$$\binom{a^m}{a^k} = \binom{\overbrace{a \cdots a}^{m \text{ times}}}{\underbrace{a \cdots a}_{k \text{ times}}} = \binom{m}{k} \quad \forall m, k \in \mathbb{N}$$

Definitions:

- $\text{rep}_2(n)$ greedy base-2 expansion of $n \in \mathbb{N}_{>0}$ starting with 1
- $\text{rep}_2(0) = \varepsilon$ where ε is the empty word

n		$\text{rep}_2(n)$
0		ε
1	1×2^0	1
2	$1 \times 2^1 + 0 \times 2^0$	10
3	$1 \times 2^1 + 1 \times 2^0$	11
4	$1 \times 2^2 + 0 \times 2^1 + 0 \times 2^0$	100
5	$1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0$	101
6	$1 \times 2^2 + 1 \times 2^1 + 0 \times 2^0$	110
\vdots	\vdots	\vdots
		$\{\varepsilon\} \cup 1\{0, 1\}^*$

Generalized Pascal triangle P_2 in base 2

		$\text{rep}_2(k)$								
		ε	1	10	11	100	101	110	111	\dots
$\text{rep}_2(m)$	ε	1	0	0	0	0	0	0	0	
	1	1	1	0	0	0	0	0	0	
	10	1	1	1	0	0	0	0	0	
	11	1	2	0	1	0	0	0	0	
	100	1	1	2	0	1	0	0	0	
	101	1	2	1	1	0	1	0	0	
	110	1	2	2	1	0	0	1	0	
	111	1	3	0	3	0	0	0	1	
	\vdots									\ddots

Binomial coefficient
of finite words:

$$\binom{\text{rep}_2(m)}{\text{rep}_2(k)}$$

Rule (not local):

$$\binom{ua}{vb} = \binom{u}{vb} + \delta_{a,b} \binom{u}{v}$$


$\binom{\text{rep}_2(m)}{\text{rep}_2(k)}$		$\text{rep}_2(k)$								
		ϵ	1	10	11	100	101	110	111	\dots
$\text{rep}_2(m)$	ϵ	1	0	0	0	0	0	0	0	
	1	1	1	0	0	0	0	0	0	
	10	1	1	1	0	0	0	0	0	
	11	1	2	0	1	0	0	0	0	
	100	1	1	2	0	1	0	0	0	
	101	1	2	1	1	0	1	0	0	
	110	1	2	2	1	0	0	1	0	
	111	1	3	0	3	0	0	0	1	
\vdots									\ddots	

The classical Pascal triangle

Questions:

- After coloring and normalization can we expect the convergence to an analogue of the Sierpiński gasket?
- Could we describe this limit object?

- Grid: first 2^n rows and columns of P_2



- Color the grid:
Color the first 2^n rows and columns of the generalized Pascal triangle P_2

$$\left(\binom{\text{rep}_2(m)}{\text{rep}_2(k)} \bmod 2 \right)_{0 \leq m, k < 2^n}$$







in

- white if $\binom{\text{rep}_2(m)}{\text{rep}_2(k)} \equiv 0 \pmod{2}$
- black if $\binom{\text{rep}_2(m)}{\text{rep}_2(k)} \equiv 1 \pmod{2}$
- Normalize by a homothety of ratio $1/2^n$
(bring into $[0, 1] \times [0, 1]$)
 \rightsquigarrow sequence $(U_n)_{n \geq 0}$ belonging to $[0, 1] \times [0, 1]$

$$U_n = \frac{1}{2^n} \bigcup_{\substack{0 \leq m, k < 2^n \text{ s.t.} \\ \binom{\text{rep}_2(m)}{\text{rep}_2(k)} \equiv 1 \pmod{2}}} \{(k, m) + Q\}$$

$$Q = [0, 1] \times [0, 1]$$

The elements U_0, \dots, U_5



The element U_2

		0	1/4	2/4	3/4	1
0		ε	1	10	11	
1/4	ε					
2/4	1					
3/4	10					
1	11					

The element U_2

		0	1/4	2/4	3/4	1
0	ε					
1/4	ε					
2/4	1					
3/4	10					
1	11					

$$\varepsilon \rightsquigarrow 0, \quad 1 \rightsquigarrow 1/4, \quad 10 \rightsquigarrow 2/4 = 1/2, \quad 11 \rightsquigarrow 3/4$$

$$0, 1, 2, 3 \rightsquigarrow w \in \{\varepsilon\} \cup 1\{0, 1\}^* \text{ with } |w| \leq 2$$

(\star)

$$(u, v) \text{ satisfies } (\star) \text{ iff } \begin{cases} u, v \neq \varepsilon \\ \binom{u}{v} \equiv 1 \pmod{2} \\ \binom{u}{v0} = 0 = \binom{u}{v1} \end{cases}$$

Example: $(u, v) = (101, 11)$ satisfies (\star)

$$\binom{101}{11} = 1$$

$$\binom{101}{110} = 0$$

$$\binom{101}{111} = 0$$

Lemma: Completion

(u, v) satisfies $(\star) \Rightarrow (u_0, v_0), (u_1, v_1)$ satisfy (\star)

Proof: Since (u, v) satisfies (\star)

$$\binom{u}{v} \equiv 1 \pmod{2}, \quad \binom{u}{v_0} = 0 = \binom{u}{v_1}$$

Proof for (u_0, v_0) :


$$\binom{u_0}{v_0} = \underbrace{\binom{u}{v_0}}_{=0 \text{ since } (\star)} + \underbrace{\binom{u}{v}}_{\equiv 1 \pmod{2}} \equiv 1 \pmod{2}$$

If $\binom{u_0}{v_{00}} > 0$ or $\binom{u_0}{v_{01}} > 0$, then v_0 is a subsequence of u .
This contradicts (\star) .


Same proof for (u_1, v_1) .

□


Example: $(u, v) = (101, 11)$ satisfies $(\star) \Rightarrow \binom{u}{v} \equiv 1 \pmod 2$



U_3



U_4



U_5

\rightsquigarrow Creation of segments of slope 1

Endpoint $(3/8, 5/8) = (\text{val}_2(11)/2^3, \text{val}_2(101)/2^3)$


Length $\sqrt{2} \cdot 2^{-3}$

$S_{u,v} \subset [0, 1] \times [1/2, 1]$ endpoint $(\text{val}_2(v)/2^{|u|}, \text{val}_2(u)/2^{|u|})$

length $\sqrt{2} \cdot 2^{-|u|}$

Definition: Set of segments of slope 1

$$\mathcal{A}_0 = \overline{\bigcup_{\substack{(u,v) \\ \text{satisfying}(\star)}} S_{u,v}} \subset [0, 1] \times [1/2, 1]$$




Modifying the slope

Example: $(1, 1)$ satisfies (\star)

Segment $S_{1,1}$ endpoint $(1/2, 1/2)$ length $\sqrt{2} \cdot 2^{-1}$

$c : (x, y) \mapsto (x/2, y/2)$ $h : (x, y) \mapsto (x, 2y)$




Modifying the slope

Example: $(1, 1)$ satisfies (\star)

Segment $S_{1,1}$ endpoint $(1/2, 1/2)$ length $\sqrt{2} \cdot 2^{-1}$

$c : (x, y) \mapsto (x/2, y/2)$ $h : (x, y) \mapsto (x, 2y)$




Modifying the slope

Example: $(1, 1)$ satisfies (\star)

Segment $S_{1,1}$ endpoint $(1/2, 1/2)$ length $\sqrt{2} \cdot 2^{-1}$

$c : (x, y) \mapsto (x/2, y/2)$ $h : (x, y) \mapsto (x, 2y)$




Modifying the slope

Example: $(1, 1)$ satisfies (\star)

Segment $S_{1,1}$ endpoint $(1/2, 1/2)$ length $\sqrt{2} \cdot 2^{-1}$

$c : (x, y) \mapsto (x/2, y/2)$ $h : (x, y) \mapsto (x, 2y)$




Modifying the slope

Example: $(1, 1)$ satisfies (\star)

Segment $S_{1,1}$ endpoint $(1/2, 1/2)$ length $\sqrt{2} \cdot 2^{-1}$

$c : (x, y) \mapsto (x/2, y/2)$ $h : (x, y) \mapsto (x, 2y)$




Definition: Set of segments of different slopes

$$c : (x, y) \mapsto (x/2, y/2)$$

$$h : (x, y) \mapsto (x, 2y)$$

$$\mathcal{A}_n = \bigcup_{0 \leq j \leq i \leq n} h^j(c^i(\mathcal{A}_0))$$




Definition: Set of segments of different slopes

$$c : (x, y) \mapsto (x/2, y/2)$$

$$h : (x, y) \mapsto (x, 2y)$$

$$\mathcal{A}_n = \bigcup_{0 \leq j \leq i \leq n} h^j(c^i(\mathcal{A}_0))$$




Lemma: $(\mathcal{A}_n)_{n \geq 0}$ is a Cauchy sequence

Definition: Set of segments of different slopes

$$c : (x, y) \mapsto (x/2, y/2)$$

$$h : (x, y) \mapsto (x, 2y)$$

$$\mathcal{A}_n = \bigcup_{0 \leq j \leq i \leq n} h^j(c^i(\mathcal{A}_0))$$




Lemma: $(\mathcal{A}_n)_{n \geq 0}$ is a Cauchy sequence

Definition: Limit object \mathcal{L}

Theorem (Leroy, Rigo, S., 2016)

The sequence $(U_n)_{n \geq 0}$ of compact sets converges to the compact set \mathcal{L} when n tends to infinity (for the Hausdorff distance).



“Simple” characterization of \mathcal{L} : (\star) condition

First step: coloring the cells of the grids regarding the parity

Extension using Lucas' theorem

Everything still holds for binomial coefficients $\equiv r \pmod{p}$ with

- base-2 expansions of integers
- p a prime
- $r \in \{1, \dots, p-1\}$

Theorem (Lucas, 1878)

Let p be a prime number.



If $m = m_k p^k + \dots + m_1 p + m_0$ and $n = n_k p^k + \dots + n_1 p + n_0$ then

$$\binom{m}{n} \equiv \prod_{i=0}^k \binom{m_i}{n_i} \pmod{p}.$$

Example with $p = 3$, $r = 2$

Left: binomial coefficients $\equiv 2 \pmod{3}$

Right: estimate of the corresponding limit object




Extension to any integer base

Everything still holds for binomial coefficients $\equiv r \pmod{p}$ with

- base- b expansions of integers with $b \geq 2$
- p a prime
- $r \in \{1, \dots, p-1\}$

Example: base 3, $\equiv 1 \pmod{2}$



Definitions:

- Fibonacci numbers $(F(n))_{n \geq 0}$:
 $F(0) = 1, F(1) = 2, F(n+2) = F(n+1) + F(n) \quad \forall n \geq 0$
 $1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, 377, 610, 987, 1597, \dots$
- $\text{rep}_F(n)$ greedy Fibonacci representation of $n \in \mathbb{N}_{>0}$ starting with 1
- $\text{rep}_F(0) = \varepsilon$ where ε is the empty word

n		$\text{rep}_F(n)$
0		ε
1	$1 \times F(0)$	1
2	$1 \times F(1) + 0 \times F(0)$	10
3	$1 \times F(2) + 0 \times F(1) + 0 \times F(0)$	100
4	$1 \times F(2) + 0 \times F(1) + 1 \times F(0)$	101
5	$1 \times F(3) + 0 \times F(2) + 0 \times F(1) + 0 \times F(0)$	1000
6	$1 \times F(3) + 0 \times F(2) + 0 \times F(1) + 1 \times F(0)$	1001
\vdots	\vdots	\vdots
		$\{\varepsilon\} \cup 1\{0, 01\}^*$

Generalized Pascal triangle P_F in Fibonacci base

$\binom{\text{rep}_F(m)}{\text{rep}_F(k)}$		$\text{rep}_F(k)$								
		ε	1	10	100	101	1000	1001	1010	...
$\text{rep}_F(m)$	ε	1	0	0	0	0	0	0	0	
	1	1	1	0	0	0	0	0	0	
	10	1	1	1	0	0	0	0	0	
	100	1	1	2	1	0	0	0	0	
	101	1	2	1	0	1	0	0	0	
	1000	1	1	3	3	0	1	0	0	
	1001	1	2	2	1	2	0	1	0	
	1010	1	2	3	1	1	0	0	1	
	\vdots									\ddots







Binomial coefficient
of finite words:


$$\binom{\text{rep}_F(m)}{\text{rep}_F(k)}$$

Rule (not local):

$$\binom{ua}{vb} = \binom{u}{vb} + \delta_{a,b} \binom{u}{v}$$


The first six elements of the sequence $(U'_n)_{n \geq 0}$





Lines of different slopes:


The tenth element



Lines of different slopes: φ^n , $n \geq 0$, with $\varphi = \frac{1+\sqrt{5}}{2}$ Golden Ratio

Theorem (S., 2018)

The sequence $(U'_n)_{n \geq 0}$ of compact sets converges to a limit compact set \mathcal{L}' when n tends to infinity (for the Hausdorff distance).



“Simple” characterization of \mathcal{L}' : (\star') condition

$$\beta \in \mathbb{R}_{>1} \quad A_\beta = \{0, 1, \dots, \lceil \beta \rceil - 1\}$$

$$x \in [0, 1) \rightsquigarrow x = \sum_{j=1}^{+\infty} c_j \beta^{-j}, \quad c_j \in A_\beta$$

Greedy way: $d_\beta(x) = c_1 c_2 c_3 \dots$

$$d_\beta(1) = \begin{cases} (\beta - 1)^\omega & \text{if } \beta \in \mathbb{N} \\ (\lceil \beta \rceil - 1) d_\beta(1 - (\lceil \beta \rceil - 1)/\beta) & \text{if } \beta \notin \mathbb{N} \end{cases}$$

Definition

$\beta \in \mathbb{R}_{>1}$ is a *Parry number* if $d_\beta(1)$ is ultimately periodic.

Example: $b \in \mathbb{N}_{>1}$: $d_b(1) = (b - 1)^\omega$

φ : $d_\varphi(1) = 110^\omega$

Parry number \rightsquigarrow algebraic integer whose conjugates have modulus less than β (Perron number)

Definition

A *linear numeration system* is a sequence $(U(n))_{n \geq 0}$ such that

- U increasing
- $U(0) = 1$
- $\sup_{n \geq 0} \frac{U(n+1)}{U(n)}$ bounded by a constant
- U linear recurrence relation
 $\exists k \geq 1, \exists a_0, \dots, a_{k-1} \in \mathbb{Z}$ such that

$$U(n+k) = a_{k-1}U(n+k-1) + \dots + a_0U(n) \quad \forall n \geq 0$$

Greedy representation in $(U(n))_{n \geq 0}$:

$$n = \sum_{j=0}^{\ell} c_j U(j) \quad \text{rep}_U(n) = c_\ell \cdots c_0 \in L_U = \text{rep}_U(\mathbb{N})$$

Example: integer base $(b^n)_{n \geq 0}$ with $b \in \mathbb{N}_{>1}$,
Fibonacci numeration system $(F(n))_{n \geq 0}$

Parry number $\beta \in \mathbb{R}_{>1} \rightsquigarrow$ linear numeration system $(U_\beta(n))_{n \geq 0}$

- $d_\beta(1) = t_1 \cdots t_m 0^\omega$

$$\begin{aligned}U_\beta(0) &= 1 \\U_\beta(i) &= t_1 U_\beta(i-1) + \cdots + t_i U_\beta(0) + 1 \quad \forall 1 \leq i \leq m-1 \\U_\beta(n) &= t_1 U_\beta(n-1) + \cdots + t_m U_\beta(n-m) \quad \forall n \geq m\end{aligned}$$

- $d_\beta(1) = t_1 \cdots t_m (t_{m+1} \cdots t_{m+k})^\omega$

$$\begin{aligned}U_\beta(0) &= 1 \\U_\beta(i) &= t_1 U_\beta(i-1) + \cdots + t_i U_\beta(0) + 1 \quad \forall 1 \leq i \leq m+k-1 \\U_\beta(n) &= t_1 U_\beta(n-1) + \cdots + t_{m+k} U_\beta(n-m-k) \quad \forall n \geq m+k \\&\quad + U_\beta(n-k) - t_1 U_\beta(n-k-1) - \cdots \\&\quad - t_m U_\beta(n-m-k)\end{aligned}$$

Examples:

$b \in \mathbb{N}_{>1} \rightsquigarrow (b^n)_{n \geq 0}$ base b

$\varphi \rightsquigarrow (F(n))_{n \geq 0}$ Fibonacci numeration system

Definition

A linear numeration system $(U(n))_{n \geq 0}$ is a *Bertrand numeration system* if


$$w \in L_U \Leftrightarrow w0 \in L_U \quad \forall w \neq \varepsilon.$$

Proposition (Bertrand-Mathis, 1989)

If $\beta \in \mathbb{R}_{>1}$ is a Parry number, then $(U_\beta(n))_{n \geq 0}$ is a Bertrand numeration system.


- $\beta \in \mathbb{R}_{>1}$ Parry number
- $(U_\beta(n))_{n \geq 0}$ Parry–Bertrand numeration system
- Generalized Pascal triangle P_β in $(U_\beta(n))_{n \geq 0}$ indexed by words of L_{U_β}
- Sequence of compact sets extracted from P_β (first $U_\beta(n)$ rows and columns of P_β)
- Convergence to a limit object (same technique)
 - Lines of different slopes: $\beta^n, n \geq 0$
 - (\star') condition
- Works modulo any prime number

Example 1

 φ^2 


Example 2

$\beta_1 \approx 2.47098$ dominant root of $P(X) = X^4 - 2X^3 - X^2 - 1$




Example 3

$\beta_2 \approx 1.38028$ dominant root of $P(X) = X^4 - X^3 - 1$




Example 4

$\beta_3 \approx 2.80399$ dominant root of $P(X) = X^4 - 2X^3 - 2X^2 - 2$



Example 5

$\beta_4 \approx 1.32472$ dominant root of polynomial $P(X) = X^5 - X^4 - 1$



In this talk:

	Generalized Pascal triangle	Convergence mod p
base 2	✓	✓
integer base	✓	✓
Fibonacci	✓	✓
Parry–Bertrand	✓	✓

- Regularity of the sequence counting subword occurrences: result for any integer base b and the Fibonacci numeration system
- Behavior of the summatory function: result for any integer base b (exact behavior) and the Fibonacci numeration system (asymptotics)

Conus textile or Cloth of gold cone



Color pattern of its shell \leftrightarrow Sierpiński gasket

Generalized Pascal triangles

Manon Stipulanti (ULiège)

49

- A. Bertrand-Mathis, Comment écrire les nombres entiers dans une base qui n'est pas entière, *Acta Math. Hungar* 54 (1989), 237–241.
- J. Leroy, M. Rigo, M. Stipulanti, Generalized Pascal triangle for binomial coefficients of words, *Adv. in Appl. Math.* 80 (2016), 24–47.
- J. Leroy, M. Rigo, M. Stipulanti, Counting the number of non-zero coefficients in rows of generalized Pascal triangles, *Discrete Math.* 340 (2017) 862–881.
- J. Leroy, M. Rigo, M. Stipulanti, Behavior of digital sequences through exotic numeration systems, *Electron. J. Combin.* 24 (2017), no. 1, Paper 1.44, 36 pp.
- J. Leroy, M. Rigo, M. Stipulanti, Counting Subword Occurrences in Base- b Expansions, *Integers* 18A (2018), Paper no. A13, 32 pp.
- M. Lothaire, *Combinatorics On Words*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1997.
- É. Lucas, Théorie des fonctions numériques simplement périodiques, *Amer. J. Math.* 1 (1878) 197–240.
- M. Stipulanti, Convergence of Pascal-Like Triangles in Parry–Bertrand Numeration Systems, *in press*.