

Developing a zebrafish-based pipeline to investigate the biological properties of medicinal plant extracts from Northern Vietnam

M. H. Tran^{1,2}, T. K. T. Nguyen², H. D. Le², L. T. Nguyen², D. T. Dinh¹, M. Muller¹

¹ University of Liege, Laboratory for Organogenesis and Regeneration, GIGA Institute, Liege, Liege, Belgium

² VNU, University of Science, Faculty of Biology, Hanoi, Hanoi, Viet Nam



Traditional medicines have been used in Vietnam for thousands of years, however, there is no clear definition of circumstances in which plants can be collected, conserved, and extracted; viable models for a productive and sustainable exploitation of natural extracts are missing, sometimes leading to severe and unfavorable consequences on health. In this study, we aim to develop a cost-effective pipeline to assess the pharmaco-/toxicological properties of these medicinal materials using the zebrafish embryo as an *in vivo* model. Proof-of-concept experiments were done by passing samples from two plants, *Caesalpinia sappan* L. and *Glochidion eriocarpum* which were collected from Northern Vietnam, to evaluate its efficiency.






Figure 1. A. Effects of *C. sappan* L. extract on zebrafish embryonic morphology and lethality were determined in the corresponding concentration-response curves (LC_{50} =500 mg/L and EC_{50} =40 mg/L). B. *C. sappan* L. extract retards or inhibits zebrafish embryos dechorination by 2mg/ml pronase.




Figure 2. Anti-angiogenic properties in a dose dependent manner, (a) hemostasis phenotype under treatment, (b) control: blood vessels of transgenic *fli1a:gfp* zebrafish embryos can be easily observed; Sub intestinal vessels (SIVs) are marked by white arrows, (c,d) 50 mg/L and 100 mg/L *C. sappan* L. extract treatment triggers ectopic angiogenesis of SIVs.




Figure 3. A. *G. eriocarpum* extract exposure led to a biphasic response from the zebrafish embryos, B. non-hatching rate was correlated with the second phase response in graph A, suggesting a protective role of the embryonic chorion



Our results may contribute to the design of a rational high throughput pharmacological screening approach using the zebrafish model

