

Trophic strategies of reef-building corals under contrasted environmental conditions of East Asian Seas

Nicolas STURARO^{1,2*}, Yun Li Eric HSIEH¹, Ling-Wen LIU¹, Pei-Ling WANG¹, Vianney DENIS¹

KEYWORDS: — Oceanography; Corals; Trophic plasticity; Stable isotopes; Taiwan.

ABSTRACT: — Most reef-building corals can derive nutrition either autotrophically or heterotrophically, which allows them to use diverse trophic pathways (HOULBRÈQUE & FERRIER-PAGÈS, 2009; MUSCATINE, 1980). Therefore, when facing environmental changes, these organisms are expected to demonstrate an intrinsic ability to acclimatise through trophic plasticity. Despite the ecological importance of these corals, our understanding of their trophic strategies is currently impaired by a lack of rigorous research approaches; a failure to consider the intraspecific variability of coral species and an oversimplification of the proxies of heterotrophic habits (e.g. corallite diameter) (MADIN et al., 2016). In order to understand how trophic plasticity could allow them to acclimatise, this study proposed to assess the trophic plasticity of morphologically contrasted coral species (e.g. *Stylophora pistillata*, *Porites* sp, *Isopora palifera*). We determined the stable isotope ratios of carbon and nitrogen in the coral host tissues and algal symbionts and compared these in corals inhabiting areas around Taiwan characterised by contrasted temperature (from high to low latitudes) and light levels (from shallow to mesophotic waters). For each area, we evaluated the intraspecific trophic variability by estimating and comparing coral isotopic niches as a proxy for trophic niches. Our results on *Stylophora pistillata* revealed no overlap of the isotopic niches for the host and symbiont from different locations (unpublished data), suggesting that these coral colonies are supported by different core resources. Moreover, the isotopic niche of higher latitude coral colonies was larger than those from the lower latitudes (unpublished data), highlighting a certain trophic plasticity that may be related to more variable environmental conditions in the higher latitudes. Analyses of additional species and locations will provide essential insights into the trophic strategies of reef-building corals and how these species might adjust their nutrition in response to environmental changes.

This research was supported by the Ministry Of Science and Technology (Taiwan, no. 106-2611-M-002-008). NS was funded by a National Taiwan University postdoctoral fellowship (Taiwan, no. 106R4000) and is currently funded by a Research Project of the Fund for Scientific Research (Belgium, PDR 31240620).

REFERENCES

- HOULBRÈQUE, F. & FERRIER-PAGÈS, C. 2009. Heterotrophy in tropical scleractinian corals. — *Biological Reviews*, **84**: 1-17.
- MADIN, J. S., HOOGENBOOM, M. O., CONNOLLY, S. R., DARLING, E. S., FALSTER, D. S., HUANG, D., KEITH, S. A., MIZEREK, T., PANDOLFI, J. M., PUTNAM, H. M., BAIRD, A. H. 2016. A trait-based approach to advance coral reef science. — *Trends in Ecology & Evolution*, **31**: 419-428.
- MUSCATINE, L. 1980. Productivity of zooxanthellae. — *In*: FALKOWSKI, P.G. (Ed.), Primary productivity in the sea. New York, Plenum Press, pp. 381-402.

¹National Taiwan University - NTU, Institute of Oceanography, Taipei, Taiwan

²University of Liège - ULiège, Laboratory of Oceanology, FOCUS, Liège, Belgium

*Corresponding Author. Email: nicolas.sturaro@uliege.be