





Fundamental parameters of massive stars: the super-computer era

Laurent Mahy
FNRS - ULiège - KU Leuven


Massive stars :



Massive stars :




Massive stars :



Massive stars :

- Luminosity $> 10^{4-5} L_{\odot}$
- Temperature $> 25000\text{K}$
(Sun $\sim 5700\text{K}$)
- Powerful stellar winds
Mass loss $\sim 10^{-5} M_{\odot}/\text{year}$
Winds can reach 3000 km/s
- Short lifetime $< 10^7 \text{ yrs}$ while Sun $\sim 10^{10} \text{ yrs}$ **Rare**
- O \rightarrow LBV/RSG \rightarrow WR \rightarrow BH/SNe




Massive stars :

- Luminosity $> 10^{4-5} L_{\odot}$
- Temperature $> 25000\text{K}$
(Sun $\sim 5700\text{K}$)
- Powerful stellar winds
Mass loss $\sim 10^{-5} M_{\odot}/\text{year}$
Winds can reach 3000 km/s
- Short lifetime $< 10^7 \text{ yrs}$ while Sun $\sim 10^{10} \text{ yrs}$ **Rare**
- O \rightarrow LBV/RSG \rightarrow WR \rightarrow BH/SNe





They play a crucial role into the ecology (evolution and dynamic) of the galaxies

= COSMIC ENGINES





Open questions: Binary interactions

- > 70% are binary or multiple systems



Open questions: Nature of the stellar winds



- Stellar winds are not homogeneous but composed of clumps.
- The mass-loss rates are thus impacted by the clumping which affects the evolutionary theory of massive stars




Sundqvist et al. (2009)

Other questions: Formation - Evolution - Death

- History of the star-forming region
- Use nebulae around evolved massive stars as tracers for their evolution
- Many more !!!





- One way to provide answers is to use : **ATMOSPHERE CODES** (Radiative transfer)

Allow to derive the internal structure of the stars
Allow to determine the stellar and wind parameters
as well as their surface abundances

- When we want to include the whole physics, very time consuming !
- For one simple star:
Temperature, gravity, luminosity,
mass-loss, terminal velocity of the wind, clumping, wind velocity law,
Helium, Carbon, Nitrogen, Oxygen surface abundances
➡ 11 parameters, min. 22 models
➡ Each model needs 1 day **1 Month of calculations**

- One way to provide answers is to use : **ATMOSPHERIC CODES** (Radiative transfer)

Allow to derive the internal structure of stars
Allow to determine the stellar and wind parameters
as well as their surface abundances

- When we want to include the hydrodynamics physics, very time consuming !

- For one simple case:
Temperature, luminosity,
mass-loss rate, velocity of the wind, clumping, wind velocity law,
Helium, Nitrogen, Oxygen surface abundances

➔ parameters, min. 22 models

➔ Each model needs 1 day

1 Month of calculations

Super calculators are required



Thank you