

SMBMIP

over the Greenland ice sheet

**Intercomparison of 11 models
over 1980-2012 forced by ERA-Interim**

1. List of models




Model Name	Model type				Reso. (km)	Modis Albedo	Institute (Country)		Contacts
	EBM	GCM	PDD	RCM					
BESSI	X				10		Bergen Univ.	NO	Zolles T.
BOX13 (MM5)				X	5		GEUS	DN	Box J.
CESM		X			~1 (100?)		Utrecht Univ.	NL	Kamphenhout L., Lenaerts J., ...
dEBM	X				1		AWI	DE	Krebs-Kanzow U.
HIRHAM				X	5	X	DMI	DN	Mottram R., Langen P., ...
MAR (ISMIP6)				X	15		Liège Univ.	BE	Fettweis X.
MPI-ESM		X			~1 (150)		Max Planck Inst.	DE	Kapsch M.
PDD1km			X		1		Sheffield Univ.	UK	Wilton D., Hanna E.
PDD5km			X		5		Lincoln Univ.	UK	Hanna E., Huybrechts P.
RACMO				X	~1 (5)	X	Utrecht. Univ	NL	Noel B., van den Broeke M., van de Berg W.J., ...
SNOWMODEL	X				5		Colorado Univ.	US	Liston G., Mernild S.

- All the models have been interpolated on the same [ISMIP6 based 1km grid](#) by using the downscaling technique of Franco et al. (2012), taking into account the sub-grid 1km topo.
- The comparison is done on the ["common" ice sheet mask](#) and common period ([1980-2012](#)).
- Except both GCMs, all the models use [ERA-Interim as forcing](#).

2. Mean 2000-2012 SMB


Mean snowline over 2000-2012 based on MODIS
(source: jonathan_ryan@brown.edu)

Units: m WE/yr




2. Mean 2000-2012 SMB


A priori, the mean snowline should be in the ablation zone !!




BOX13_box




BESS1_zolles




CESM_kamphenhout




dEBM_krebs




HIRHAM_mottram




MAR_fettweis




MPI-ESM_kapsch




PDD1km_wilton



PDD5km_hanna




RACMO_noel



SNOWMODEL_liston


3. Ensemble mean




3. Ensemble mean

Models vs ensemble mean

Units: m WE/yr



4. Evaluation



PROMICE SMB data base
(Machguth et al., 2016)

	Model	Bias (mWE)	RMSE	Correl.
Ice cores (#260)	BESSI_zolles	-0.07	0.11	0.87
	BOX13_box	0.02	0.13	0.83
	CESM_kamphenhout	0.05	0.14	0.80
	dEBM_krebs	-0.01	0.08	0.89
	HIRHAM_mottram	0.02	0.13	0.82
	MAR_fettweis	0.01	0.08	0.93
	MPI-ESM_kapsch	0.01	0.12	0.76
	PDD1km_wilton	-0.01	0.04	0.97
	PDD5km_hanna	-0.01	0.04	0.96
	RACMO_noel	-0.02	0.08	0.88
	SNOWMODEL_liston	-0.05	0.12	0.87
Full PROMICE SMB data Base (#1694)	BESSI_zolles	0.37	0.87	0.79
	BOX13_box	-0.01	0.61	0.88
	CESM_kamphenhout	0.11	0.66	0.86
	dEBM_krebs	0.03	0.70	0.84
	HIRHAM_mottram	0.13	0.58	0.90
	MAR_fettweis	0.10	0.49	0.93
	MPI-ESM_kapsch	-0.06	0.75	0.81
	RACMO_noel	-0.07	0.61	0.89
SNOWMODEL_liston	-0.30	0.67	0.89	
Only main ice sheet SMB data base (#1439)	BESSI_zolles	0.45	0.89	0.81
	BOX13_box	-0.01	0.58	0.90
	CESM_kamphenhout	0.11	0.61	0.89
	dEBM_krebs	0.05	0.76	0.82
	HIRHAM_mottram	0.09	0.57	0.91
	MAR_fettweis	0.10	0.48	0.93
	MPI-ESM_kapsch	-0.05	0.69	0.85
	PDD1km_wilton	-0.18	0.69	0.89
	PDD5km_hanna	-0.17	0.72	0.86
	RACMO_noel	-0.05	0.62	0.90
	SNOWMODEL_liston	-0.31	0.61	0.92

Observations are not representative of the whole ice sheet !!

4. Evaluation

Mean SMB vs Bare ice area

Comparison of the mean 2000-2012 bare ice area from MODIS with respect to the mean 2000-2012 SMB from models.











	Percentage of the main ice sheet		
	SMB > 200mm/yr and bare ice in MODIS	SMB < -200mm/yr and no bare ice in MODIS	Total Error
BOX13_box	1.0	2.3	3.3
dEBM_krebs	1.6	2.8	4.4
MAR_fettweis	0.5	2.5	2.9
CESM_kampenhout	1.5	0.9	2.4
BESSI_zolles	3.3	0.8	4.1
PDD1km_wilton	1.6	1.7	3.3
PDD5km_hanna	1.2	4.2	5.4
RACMO_noel	0.9	2.1	3.0
SNOWMODEL_liston	0.4	9.7	10.1
HIRHAM_mottram	0.9	1.6	2.5
MPI-ESM_kapsch	0.6	1.5	2.1

In the bare ice area, SMB should be negative!

SMB can be negative even if bare ice is not detected by MODIS

5. SMB vs GRACE

Source GRACE:
andreas.groh@tu-dresden.de




on the common
ice sheet mask

2003 - 2012

5. SMB vs GRACE

Source GRACE:
andreas.groh@tu-dresden.de



GrIS	Trend.	R ²	RMSE
BESSI	-143.8	0.98	86.9
BOX13	-83.6	0.98	79.7
CESM	-188.1	0.98	94.9
dEBM	-41.2	0.98	79.1
HIRHAM	-81.1	0.99	51.9
MAR	-73.6	0.99	48.4
MPI-ESM	-177.2	0.93	186.5
PDD1km	-91.4	0.98	82.6
PDD5km	-80.5	0.98	74.6
RACMO	-85.8	0.99	51.8
SNOWMODEL	44.2	0.98	90

Units: GT/yr


Basin 2	Trend.	R ²	RMSE
BESSI	1.3	0.72	8.4
BOX13	2.8	0.77	7.4
CESM	3.5	0.6	10.2
dEBM	13.3	0.63	12.3
HIRHAM	17.4	0.63	9.9
MAR	5.4	0.77	7.4
MPI-ESM	-0.5	0.29	13.3
PDD1km	5.2	0.71	8.8
PDD5km	8.1	0.69	10.8
RACMO	7.7	0.74	7.8
SNOWMODEL	25.5	0.5	21.9

Basin 5	Trend.	R ²	RMSE
BESSI	-11.1	0.94	16.2
BOX13	-9.9	0.97	12.2
CESM	-17	0.96	14.2
dEBM	-2.6	0.95	14.3
HIRHAM	-2.9	0.93	17.4
MAR	-8	0.98	9.9
MPI-ESM	-10.8	0.92	19.6
PDD1km	-6.7	0.95	14.7
PDD5km	-8.8	0.95	14.4
RACMO	-13.1	0.97	11.2
SNOWMODEL	-0.1	0.97	11.5

Basin 6	Trend.	R ²	RMSE
BESSI	-10.5	0.89	36
BOX13	13.5	0.95	24.8
CESM	-11.8	0.87	39.6
dEBM	17.8	0.93	27.9
HIRHAM	-11.8	0.79	49.6
MAR	9.8	0.97	18.9
MPI-ESM	-10.9	0.65	66.3
PDD1km	1.5	0.92	31.9
PDD5km	4.5	0.92	30.5
RACMO	2.9	0.97	18.3
SNOWMODEL	27.6	0.92	29.7




Trend of the GRACE - SMB
i.e. "trend" of the ice dynamics



RMSE if we add this trend to SMB

Next steps:


- Comparison with ~ 200 snow pits from Jason Box.
- Comparison with ICEsat ?
- Comparison with other SMB products you have ?
- Addition of the ensemble mean in the comparison.



Next SMB measurements would be done in the "red" areas ;-)

Thanks for your attention!

x. Mean SMB



Mean SMB over 1980-2012 (GT/yr)

	GrIS	B 1	B 2	B 3	B 4	B 5	B 6	B 7	B 8
BESSI	386	14	18	74	109	42	45	53	32
BOX13	425	12	15	84	151	32	6	63	61
CESM	419	19	43	82	108	34	36	56	40
dEBM	387	10	14	81	115	37	28	58	44
HIRHAM	391	23	25	67	114	26	4	65	67
MAR	369	8	21	89	119	20	5	56	51
MPI-ESM	283	24	27	63	82	9	-25	46	57
PDD1km	329	26	26	62	65	28	13	65	43
PDD5km	282	12	15	63	66	28	4	62	30
RACMO	352	9	21	67	104	24	5	65	56
SNOWMODEL	90	-45	-28	51	90	36	-26	24	-11
<i>Mean</i>	337.5	10	18	71	102	29	9	56	43