



ISSN: 0960-2011 (Print) 1464-0694 (Online) Journal homepage: https://www.tandfonline.com/loi/pnrh20

## Corrigendum

To cite this article: (2009) Corrigendum, , 19:5, 783-783, DOI: <u>10.1080/09602010903206177</u>

To link to this article: <a href="https://doi.org/10.1080/09602010903206177">https://doi.org/10.1080/09602010903206177</a>

	Published online: 02 Sep 2009.
	Submit your article to this journal 🗗
ılıl	Article views: 102
Q <sup>L</sup>	View related articles 🗹



## Corrigendum

Volume 18, Issue 3 (June 2008), pp. 343–371, Face—name association learning in early Alzheimer's disease: A comparison of learning methods and their underlying mechanisms by Nathalie Bier, Martial Van der Linden, Lise Gagnon, Johanne Desrosiers, Stéphane Adam, Stéphanie Louveaux, and Julie Saint-Mleux.

Please note that in Table 4 on page 359, the wrong data was submitted in the 'Delayed-Total score combining free and cued recall (/30): AD and Controls' rows. The statistical analyses are not affected by this error. The correct version of the table is reprinted below.

TABLE 4
Test performances of the AD and Control groups on immediate and delayed recall

	Learning methods				
Performance measures	Spaced retrieval	Vanishing cues	Errorless	Explicit trial-and- error	Implicit trial-and- error
Immediate					
Free recall (/5)					
AD: Mean $\pm SD$	$1.5 \pm 1.5$	$1.1 \pm 1.4$	$.80 \pm .86$	$1.3 \pm 1.2$	$.60 \pm .74$
Max	4	5	3	3	2
Controls: Mean $\pm SD$	$4.5 \pm 1.1$	$4.5 \pm 1.1$	$3.9 \pm 1.1$	$4.5 \pm .83$	$4.3 \pm 1.0$
Total score combining free and c	ued recall (/30):				
AD: Mean $\pm SD$	$22 \pm 6.5$	$22.7 \pm 4.9$	$18.9 \pm 5.9$	$21.9 \pm 5.4$	$21.5 \pm 3.2$
Controls: Mean $\pm SD$	$29.4 \pm 1.5$	$29.5 \pm 1.1$	$28.7 \pm 1.3$	$29.5 \pm .83$	$29.3 \pm 1.0$
Recognition (/5)					
AD: Mean $\pm SD$	$1.8 \pm 1.7$	$1.3 \pm 1.5$	$1.6 \pm 1.0$	$1.6 \pm 1.6$	$1.0 \pm 1.7$
Max	5	4	3	5	5
Controls: Mean $\pm SD$	$4.7 \pm 1.0$	$4.3 \pm 1.4$	$4.9 \pm .51$	$4.6 \pm 1.1$	$4.6 \pm .91$
Delayed					
Free recall (/5)					
AD: Mean $\pm SD$	$.10 \pm .29$	$.30 \pm .45$	$0 \pm 0$	$0\pm0$	$0\pm0$
Proportion of associations retained	7%	27%	0%	0%	0%
Controls: Mean $\pm SD$	$.40 \pm .92$	$.50 \pm .69$	$.40 \pm 1.2$	$.50 \pm .67$	$.40 \pm .82$
Total score combining free and c	ued recall (/30):				
AD: Mean $\pm SD$	$14.3 \pm 4.5$	$14.7 \pm 4$	$12.6 \pm 3.6$	$13.6 \pm 3.6$	$14.4 \pm 4$
Controls: Mean $\pm SD$	$17.5 \pm 4.6$	$17.3 \pm 2.9$	$16.2 \pm 5.5$	$15.8 \pm 5.5$	$17.5 \pm 4.8$
Recognition (/5)					
AD: Mean $\pm SD$	$.80 \pm 1.1$	$1.4 \pm 1.3$	$.70 \pm .65$	$1.7 \pm 1.6$	$.80 \pm 1.1$
Controls: $Mean \pm SD$	$2.7 \pm 2.0$	$3.5 \pm 1.8$	$2.1 \pm 1.8$	$2.7 \pm 2.0$	$2.7 \pm 1.8$

SD = standard deviation; Max = Maximum score obtained by the best participant.

<sup>© 2009</sup> Psychology Press, an imprint of the Taylor & Francis Group, an Informa business http://www.psypress.com/neurorehab DOI:10.1080/09602010903206177