

GDNF/GFR α 1 Complex Abrogates Self-Renewing Activity of Cortical Neural Precursors Inducing Their Differentiation

Antonela Bonafina,¹ Paula Aldana Fontanet,¹ Gustavo Paratcha,¹ and Fernanda Ledda^{1,*}

¹Division of Molecular and Cellular Neuroscience, Institute of Cell Biology and Neuroscience (IBCN)-CONICET-UBA, School of Medicine, University of Buenos Aires (UBA), Buenos Aires CP 1121, Argentina

*Correspondence: mledda@fmed.uba.ar

<https://doi.org/10.1016/j.stemcr.2018.01.019>

SUMMARY

The balance between factors leading to proliferation and differentiation of cortical neural precursors (CNP) determines the correct cortical development. In this work, we show that GDNF and its receptor GFR α 1 are expressed in the neocortex during the period of cortical neurogenesis. We show that the GDNF/GFR α 1 complex inhibits the self-renewal capacity of mouse CNP cells induced by fibroblast growth factor 2 (FGF2), promoting neuronal differentiation. While GDNF leads to decreased proliferation of cultured cortical precursor cells, ablation of GFR α 1 in glutamatergic cortical precursors enhances its proliferation. We show that GDNF treatment of CNPs promoted morphological differentiation even in the presence of the self-renewal-promoting factor, FGF2. Analysis of GFR α 1-deficient mice shows an increase in the number of cycling cells during cortical development and a reduction in dendrite development of cortical GFR α 1-expressing neurons. Together, these results indicate that GDNF/GFR α 1 signaling plays an essential role in regulating the proliferative condition and the differentiation of cortical progenitors.

INTRODUCTION

The mammalian cerebral cortex consists of diverse areas with specific functions and its cellular organization must be strictly regulated to ensure proper functionality (Bystron et al., 2008; Kowalczyk et al., 2009; Rakic, 2006). The development of this structure follows an organized generation of neurons and glial cells from local neural stem cells (NSCs). These cells are critical for the generation of the two main classes of neurons that populate the cerebral cortex, the excitatory projection neurons, which use glutamate as neurotransmitter, and the inhibitory interneurons, which use γ -amino butyric acid (GABA) (Anderson et al., 1997, 2002). While excitatory cortical neurons are generated from NSCs from the dorsal telencephalon, inhibitory cortical interneurons are mainly generated from the germinal zones of the ventral telencephalon. Different subtypes of excitatory cortical projection neurons are born in overlapping temporal waves from local NSCs to generate defined cortical layers establishing the six-layered structure of the mature neocortex with layers VI and V generated first, followed by layers IV, III, and II (Molyneux et al., 2007). On the other hand, interneurons generated from NSCs of the ventral telencephalon, migrate tangentially a long distance to populate the dorsal cortical structure. A correct balance between excitatory glutamatergic neurons and inhibitory (GABAergic) interneurons is necessary for the correct functioning of the cerebral cortex. Thus, a correct regulation of this process is necessary, since any inconvenience may cause possible nervous system disorders, such as cerebral malformations or psychiatric diseases. Emerging evidence indicates that intrinsic and extrinsic

cues are involved in cortical precursor development. Particularly, fibroblast growth factors (FGFs) are secreted signaling molecules that play cue roles in the cerebral cortex formation (Vaccarino et al., 1999a) and patterning (Fukuchi-Shimogori and Grove, 2001; Garel et al., 2003; Hebert and Fishell, 2008). Several FGF ligands, including FGF2, 3, 7, 8, 10, 15, 17, and 18 are expressed in the rostral telencephalic midline and early cortical primordium (Cholfin and Rubenstein, 2007; Garel et al., 2003; Vaccarino et al., 1999b), and have been involved in neurogenesis (Borello et al., 2008; Raballo et al., 2000). Among the four FGF receptors (FGFRs), FGFR 1–3 are expressed in the developing CNS (Ford-Perriss et al., 2001). Despite FGFRs having been involved in the control of CNS growth including the hippocampus (Ohkubo et al., 2004) and the cerebral cortex (Kang et al., 2009; Stevens et al., 2010), the cellular and molecular mechanisms of FGFR function in the process of neurogenesis and cortical surface area expansion is still unclear (Rash et al., 2011). Most FGF ligand studies focused on the FGF2 or basic FGF ligand, which binds with high affinity to FGFR1 and certain FGFR2, 3, and 4 isoforms. FGF2 is highly expressed in the developing nervous system, especially in the ventricular zone of the cerebral cortex (Weise et al., 1993), and also functions as a mitogen for neural progenitors in culture (Qian et al., 1997). Other extrinsic cues such as BMPs, Wnts, as well as neurotrophins NT3 and BDNF, have been involved in the control of cortical progenitor development (Barnabe-Heider and Miller, 2003; Bartkowska et al., 2007). Recent evidence suggests that members of the glial cell line-derived neurotrophic factor (GDNF) family could have an important role in cortical development (Yuzwa et al., 2016).



GDNF was the first identified member of the family of neurotrophic factors, consisting of GDNF, neurturin (NTRN), persephin (PSPN), and artemin, and together make up the GDNF family of ligands (GFLs) (Saarma, 2000). These factors are best known for regulating the biology of peripheral and CNS neurons. Signaling by GFLs is mediated by alternative multicomponent receptor complex containing a glycosylphosphatidylinositol-anchored subunit named GDNF family receptor alpha (GFR α) together with either RET receptor tyrosine kinase or the neural cell adhesion molecule (NCAM). Four GPI-GFR α (GFR α 1-4) have been described which provide ligand-specific binding activity for each GFLs. Thus, GDNF binds preferentially to GFR α 1 receptor, while NRTN signals through GFR α 2, ART through GFR α 3, and PSPN through GFR α 4. Throughout the nervous system and, particularly in the forebrain, GFR α receptors are more widely expressed than RET, suggesting that GFR α s may signal independently of RET, presumably in collaboration with NCAM. Physiological functions of GDNF signaling in the absence of RET have been reported in different systems including GABAergic neuron development and commissural neurons of the spinal cord (Canty et al., 2009; Charoy et al., 2012; Paratcha et al., 2003, 2006; Pozas and Ibanez, 2005). Interestingly, syndecan-3 has been described as an alternative signaling partner of GDNF and NTRN in the brain without the involvement of their conventional receptors. GDNF signaling through syndecan-3 has been involved in migration of GABAergic neurons during cortical development (Bespalov et al., 2011). During the last years, we have described that GDNF through GFR α 1 and NCAM receptors play a crucial role in glutamatergic hippocampal dendrite development, and that this complex is required for proper hippocampal connectivity (Irala et al., 2016).

Based on the expression pattern of GDNF and its receptor, GFR α 1, at early stages in cortical development, we investigated the function of this molecular system in cortical glutamatergic precursor development. In the present work, we provide evidence indicating that GDNF and its GPI-anchored receptor, GFR α 1, expressed in the forebrain during the period of cortical neurogenesis, play an important role during embryonic cortical precursor development.

RESULTS

GFR α 1 Localization during Early Cortical Development

To determine whether GFR α 1 is expressed in neural precursors of the embryonic cortex, we analyzed expression of *Gfra1* mRNA. RT-PCR of RNA isolated from rat neocortex

at embryonic day 13.5 (E13.5) showed that *Gfra1* is expressed at low levels at this time point, when the cortex is predominantly comprised of neural precursors, but its expression increased throughout embryogenesis and decreased in the cortex at postnatal day 0 (P0, Figure 1A). Expression of *Gdnf* and *Ncam* mRNA remained similar during the analyzed period, while the expression of other member of the GFR α receptor family, *Gfra2* mRNA, was very low during the early cortical development increasing at birth (P0) (Figure 1A) and during postnatal cortical development as described previously (Burazin and Gundlach, 1999). In agreement with other works, we did not detect expression of the GDNF canonical receptor *Ret* mRNA during cortical development (Glazner et al., 1998; Golden et al., 1999; Lenhard and Suter-Crazzolara, 1998; Trupp et al., 1999). The expression of *Fgfr1* mRNA, the most expressed FGFR in the CNS (Ford-Perriss et al., 2001), was clearly detected at early cortical developmental stages, and its expression was markedly decreased at later embryonic stages dropping at P0 (Figure 1A). The expression pattern of *Fgfr1* mRNA is in agreement with its role as a mediator of FGF2 in early precursor proliferation (Maric et al., 2007). The realization that *Gfra1* mRNA was expressed exclusively during early cortical stages prompted us to investigate a possible role of this complex in cortical development.

To define GFR α 1 localization, we performed immunofluorescence analysis of E13.5 mouse cortical sections and compared the distribution of GFR α 1 with SOX2, which is expressed in the proliferative ventricular zone and subventricular zone (VZ and SVZ, respectively) by NSCs (Gotz et al., 1998) and β III-tubulin, which is expressed in the cortical plate by postmitotic neurons. We observed that GFR α 1 expression was very low in cortical precursors from VZ/SVZ and high in postmitotic cells. A similar distribution was observed for NCAM (Figure 1B). GFR α 1 was clearly expressed in cells positive for the postmitotic nuclear marker, Tbr1 (Figure 1C). RET immunoreactivity could not be detected in embryonic cortical sections, but neurons from cervical dorsal root ganglia, used as positive expression control, were clearly labeled (See Figure S1A). Antibody specificity was confirmed in heterologous cells transfected with *Ret* cDNA by immunofluorescence (Figure S1B). Moreover, we could not detect either *Ret* mRNA expression, or RET activity in cultured cortical precursors stimulated with GDNF and analyzed by immunoblot (Figure S1C).

Expression of GFR α 1 was analyzed in dissociated cultures of rat E14.5 cortical precursors. After 5 days *in vitro* (DIV), most cortical precursors differentiated into postmitotic neurons in the absence of ligand, whereas cultures treated with saturating concentrations of the mitotic ligand FGF2, were composed predominantly by proliferating precursors,




Figure 1. Developmental Expression and Localization of GFR α 1 during Early Cortical Development

(A) Analysis of developmental expression of *Gdnf*, *Gfra1*, *Gfra2*, *Ncam*, *Fgfr1*, and *Ret* mRNA by semiquantitative RT-PCR (27 cycles) in rat cortex at embryonic day (E) 13.5, E15.5, E17.5, and newborn (P0). Expression of the housekeeping gene TATA binding protein (*Tbp*) was analyzed as control. *Ret* mRNA expression was analyzed in rat E18.5 dorsal root ganglia (DRG) and in PC12 cells stimulated with nerve growth factor (NGF), a treatment that induces *Ret* expression.

(B) Localization by immunofluorescence of GFR α 1 in coronal sections of mouse E14.5 cortex. Expression of GFR α 1 (red) and Sox2 (green) are shown in the upper panel, GFR α 1 (red) and β III-tubulin (green) in the middle panel, GFR α 1 (red), and NCAM (green) in the lower panel. VZ, ventricular zone; SVZ, subventricular zone; CP, cortical plate. Scale bar, 100 μ m.

(C) Localization by immunofluorescence of GFR α 1 (green) and the neuronal nuclear marker Tbr1 (red) in CP. Scale bar, 50 μ m. High-magnification image of boxed area is shown. The arrows indicate cells expressing GFR α 1 with nuclei positive for Tbr1. Scale bar, 20 μ m.

(D) Analysis of cortical progenitors cultured in the absence or presence of FGF2 (40 ng/mL) for 5 days *in vitro* (DIV) and stained for nestin (red) or GFR α 1 (green). Scale bar, 50 μ m.

(E) Upper bar graph, quantification of the percentage of nestin⁺ cells in the presence and absence of FGF2 (40 ng/mL). Lower bar graph, quantification of the percentage of GFR α 1⁺ cells in the presence and absence of FGF2 (40 ng/mL). Data are mean \pm SEM of n = 3 independent experiments. *p < 0.05, ***p < 0.001 (Student's t test).

(F) Co-localization of GFR α 1 (green) and the neuronal marker β III-tubulin (red) in dissociated cortical precursors maintained in the absence of FGF2. Scale bar, 50 μ m.

(G) Analysis by semiquantitative RT-PCR of *Gfra1*, *Ncam* and *Ret* in proliferating (+FGF2, 40 ng/mL) and non-proliferating (–FGF2) cortical precursors. Analysis of *Tbp* mRNA was used as control. *Ret* mRNA expression analyzed in E18.5 rat DRG and in PC12 cells treated with NGF, which induced *Ret* expression, is shown as positive control. See also Figure S1.

positive for nestin, a specific intermediate filament of neural precursors. To determine which cells were expressing GFR α 1 receptor, we performed double-label immunofluorescence analysis for GFR α 1 and nestin, or β III-tubulin. In the cultures maintained in the presence of FGF2 the major-

ity of the cells were nestin⁺ (\approx 70%), while cultures maintained in the absence of FGF2 resulted in an enhanced neuronal differentiation and less nestin⁺ cells (\approx 20%). Expression of GFR α 1 was observed in \approx 40% of total cells grown in the absence of FGF2 and in \approx 8% of the cells



maintained in the presence of FGF2 (Figures 1D and 1E). All of the β III-tubulin cells of these cultures were highly positive for GFR α 1 (Figure 1F). Although the majority of the nestin⁺ cells were negative for GFR α 1, many of them show low levels of GFR α 1 expression. Analysis by RT-PCR revealed that *Gfra1* and *Ncam* mRNA expression was higher in cultures maintained in the absence of FGF2 and lower in proliferating cultures, maintained in the presence of FGF2, while *Ret* mRNA could not be detected in these cells independently of the presence of FGF2 (Figure 1G).

GDNF and FGF Regulate Opposite Processes in Cortical Precursor Cells

It is known that FGF2 is widely distributed throughout the dorsal telencephalon. High expression of FGF2 in the VZ correlates with the initial expansion of NSCs, whereas its low expression near the pial surface is associated with neuronal differentiation. Thus, the distribution of FGF2 *in vivo* is consistent with the dose-dependent effect of FGF2 on self-renewing of NSCs *in vitro*. Similar to FGF2, FGFRs are widely distributed throughout the dorsal telencephalon and exhibit progressively reduced levels of expression along the ventriculopial gradient, being lower in the pial surface (Vaccarino et al., 1999a; Weise et al., 1993). As the expression of GFR α 1 was higher in differentiated cells and lower in cells positive for SOX2 (Figure 1B), we decided to study whether FGF2 could regulate the expression levels of GFR α 1. To this end, cortical precursor cells isolated from E14.5 rat cortex were cultured in the presence of different concentrations of FGF2, mimicking what happens during cortical neurogenesis. After 5 DIV, the levels of *Gfra1* mRNA were analyzed by real-time PCR. We observed that cultures maintained in the presence of different concentrations of FGF2, which are enough to maintain cortical precursors in proliferation (Qian et al., 1997), expressed very low levels of *Gfra1* and high levels of *Fgfr1* mRNA, while cultures maintained in the absence of FGF2 resulted in high levels of *Gfra1* and low levels of *Fgfr1* mRNA (Figure 2A). This result indicates that *Gfra1* and *Fgfr1* mRNA expression levels are modulated by FGF2 in an opposite manner (Figure 2A).

To analyze the effect of GDNF on the expression of *Gfra1* and *Fgfr1* mRNA, cortical cultures were grown in the presence of different concentrations of FGF2 plus GDNF. We observed that, in the presence of low concentrations of FGF2 and GDNF, *Gfra1* mRNA expression does not decrease as much as when the cultures were grown in the presence of FGF2 alone, indicating that GDNF could modulate the expression of its own receptor, making cortical precursor cells more responsive to the neurotrophic factor (Figure 2B). However, the presence of a high concentration of FGF2 (25 ng/mL) was enough to decrease substantially the levels of *Gfra1* mRNA, even in the presence of GDNF. Similar re-

sults were observed for *Gfra2* mRNA. As it has been described that GDNF induces RET expression in diverse systems (Arvidsson et al., 2001), we analyzed the levels of *Ret* mRNA expression in cortical precursors maintained with different concentrations of FGF2 and GDNF, but we were not able to detect expression of *Ret* mRNA in these conditions (Figure S1D).

We next examined the effects of GDNF on precursor cells maintained under self-renewing conditions in the presence of FGF2 for proliferation and neural differentiation. Immunostaining with antibodies against Ki67, a protein that is present during all active phases of cell cycle, but is absent from resting cells (G₀) (Burdon et al., 2002), indicated that, under low concentrations of FGF2 (5 ng/mL), the GDNF treatment decreases the proportion of proliferating cells compared with controls maintained in the presence of FGF2 alone (Figure 2D). The decreased proliferation observed after GDNF exposure correlated with an increase in neuronal differentiation of precursors to postmitotic neurons positive for β III-tubulin (Figure 2D). In the presence of higher concentrations of FGF2, the effect of GDNF on cell proliferation and neuronal differentiation was abrogated (Figure 2E). This result is in agreement with the low levels of *Gfra1* mRNA observed when the precursors were exposed to high concentrations of FGF2 (Figures 1G, 2A, and 2B).

As it is known that Akt is involved in neuronal precursor proliferation, and it is activated by FGF2 (Burdon et al., 2002; Jin et al., 2005), we analyzed the effect of GDNF on FGF2-induced Akt phosphorylation. As shown in Figure 2F, stimulation of cortical precursors with 5 ng/mL of FGF2 induced activation of Akt, but when the cells were treated with FGF2 in the presence of GDNF, the phosphorylation levels of Akt were reduced. The activation of MAPK, which is also known to be induced by FGF2 was not affected by GDNF, indicating that GDNF specifically inhibits FGF2-induced Akt pathway in cortical precursors (Figure 2F).

Altogether these results indicate that GDNF counteracts the mitogenic stimulation of low concentrations of FGF2 on cortical progenitors by modulating Akt activity and promoting neuronal differentiation.

GDNF Decreases Proliferation of Cortical Progenitors

To test whether culture conditions, such as adhesion can affect GDNF function, we assayed the effects of GDNF on cortical precursor grown maintained as neurospheres in the presence of FGF2 (25 ng/mL). The neurosphere assay is a useful tool for determining the proliferative capacity of a cell population because only progenitor cells with high proliferative capacity have the ability to form neurospheres in presence of FGF2. Our data indicate a significant decrease in the number and diameter of spheres formed in the medium containing FGF2 and GDNF compared with




Figure 2. GDNF Regulates Cortical Precursor Response to FGF2

(A and B) Analysis of *Gfra1* and *Fgfr1* mRNA expression by semiquantitative RT-PCR (27 cycles) in cortical precursor cultures maintained in the absence (A) or presence of GDNF (B), plus different concentrations of FGF2. *Tbp* mRNA expression was analyzed as control. The bar graphs show the levels of *Gfra1* mRNA measured by real-time PCR using the expression of the housekeeping gene *Tbp* for normalization. Shown are mean \pm SEM of triplicate determinations. *** $p < 0.001$, * $p < 0.05$ (Student's *t* test).

(C) Representative images of proliferating cortical progenitors grown for 5 DIV in the absence or presence of GDNF (100 ng/mL) plus different concentrations of FGF2 stained with antibodies against β III-tubulin (green) and Ki67 (red). Scale bar, 20 μ m.

(D and E) Bar graphs showing the percent of Ki67⁺ (left) and β III-tubulin⁺ (right) cells in cultures maintained in the presence of 5 ng/mL (D) or 25 ng/mL (E) of FGF2 with or without GDNF (100 ng/mL). Data are mean \pm SEM of *n* = 3 independent experiments * $p < 0.05$, ** $p < 0.005$ (Student's *t* test).

(F) Analysis of Akt and MAPK activation induced by FGF2 and GDNF on cortical progenitors. After starving overnight, cultures were stimulated for 15 min with 5 ng/mL of FGF2 or FGF2 (FGF 5 ng/mL plus GDNF 100 ng/mL). The lysates were analyzed by pAkt and pMAPK immunoblotting. The same membrane was reprobbed with anti-actin antibodies. Fold of change in pAkt and pMAPK relative to total actin is indicated.

the medium containing only FGF2 (Figure 3A). Interestingly, in this culture condition, GDNF was able to inhibit the proliferative effects induced by high concentrations of FGF2. We evaluated the proportion of cells per sphere that were positive for Ki67⁺ and nestin⁺ by immunofluorescence. We observed that the addition of GDNF to these cultures decreased the proportion of Ki67-expressing cells, and also decreased the percentage of nestin⁺ progenitors expressing Ki67 (proliferating precursors) (Figure 3B). Interestingly, a robust increase in the number of differentiated β III-tubulin⁺ cells was observed in neurospheres grown in

the presence of GDNF (see Figure S3). Cell death was not affected as similar results were obtained in the presence of the apoptosis inhibitor, Z-VAD-FMK (see Figures S2A–S2C).

As the GDNF transmembrane signaling receptor NCAM, was detected in cortical precursors (Figure 1B), we analyzed whether NCAM could be involved in the effects of GDNF on proliferation. To investigate this, neurospheres were grown in the presence of NCAM^{ECD} function-blocking antibody. Interestingly, the effects of GDNF on proliferation could be reverted in the presence of the anti-NCAM^{ECD}




Figure 3. GDNF Treatment Decreases Proliferation of Cortical Precursor Cells

(A) Neurospheres from cortical precursor cells grown for 5 DIV in the presence or absence of GDNF (100 ng/mL) plus FGF2 (25 ng/mL) stained with DAPI. Scale bar, 200 µm. The bar graphs show the quantification of the number and diameter of neurospheres that were generated. The results are expressed as mean ± SEM of three independent experiments. * $p < 0.05$ (Student's t test, see also Figure S2). (B) Images of neurospheres derived from cortical precursor cells grown as described in (A) stained with nestin (red), Ki67 (green), and DAPI. Scale bar, 50 µm. The bar graphs show the quantification of the percentage of Ki67⁺ (Ki67⁺/DAPI) cells as well as proliferating precursor (Ki67⁺ nestin⁺/nestin⁺) cells per sphere grown in the presence of FGF2 (25 ng/mL) or FGF2 (25 ng/mL) plus GDNF (100 ng/mL). The results are expressed as the mean ± SEM of three independent experiments. * $p < 0.05$ (Student's t test). (C) Bar graphs showing the quantification of the percent and diameter of neurospheres that were generated from cortical precursor cells grown for 5 DIV in the presence of the indicated trophic factors and blocking anti-NCAM^{ECD} antibodies. The results are expressed as the average percentage of reduction in the number of neurospheres grown in the presence of GDNF relative to the neurospheres grown in the presence of FGF in the absence of GDNF (indicated as dotted line). The data are expressed as the mean ± SEM three independent experiments. * $p < 0.05$ (Student's t test). (D) Analysis of *CyclinD1* and *Cyclin E*, *p21* and *p27*, and *GFRα1* mRNA from neurospheres cultured in the presence of FGF2 or FGF2 plus GDNF by semiquantitative PCR. (E) Bar graph showing the levels of *CyclinD1* and *GFRα1* mRNA analyzed by real-time PCR. The expression of the specific mRNA was normalized to that of the housekeeping gene *Tbp*. Shown are averages ± SEM of triplicate determinations, ** $p < 0.001$ (Student's t test).

antibodies, indicating the requirement of NCAM in this process (Figure 3C).

To analyze the role of GDNF on mitogenic cortical cultures, we investigated the influence of GDNF on molecules known to regulate the cell cycle. To this end, neurospheres were grown in the presence of FGF2, or FGF2 plus GDNF, and the levels of different cyclins and negative regulators of the cell cycle were analyzed by real-time PCR. Neurospheres grown in the presence of FGF2 are characterized by an upregulation of the levels of cyclins as well as by a strong decrease of the cell-cycle inhibitors (Andreu et al., 2015; Mira et al., 2010). We observed that the decrease pro-

liferation after GDNF exposure correlated with the downregulation of *Cyclin D1* and *Cyclin E* mRNA and the upregulation of the cyclin-dependent kinase inhibitors *p21* and *p27* mRNAs (Figures 3D and 3E). This result indicates that GDNF has opposite effects with respect to FGF2 on the induction of the levels of positive and negative regulators of the cell cycle. Interestingly, we also observed an upregulation of *GFRα1* mRNA in cells cultured in the presence of GDNF compared with neurospheres maintained only in the presence of FGF2 (Figures 3D and 3E).

Neurospheres contain cells at different stages of differentiation, with only a fraction of cells that maintains the



initial proliferative characteristics. This small fraction of cells is the one that maintains the capacity to generate secondary neurospheres. When primary neurospheres, which were grown in the presence or absence of GDNF, were dissociated and cultured only in the presence of FGF2, we did not find differences in the number or in the diameter of secondary neurospheres. These results indicated that GDNF treatment inhibits the proliferation of neural precursors inducing its differentiation and maintaining NSCs in a resting state (see [Figures S3C and S3D](#)).

GDNF/GFR α 1 Complex Is Required to Control Cortical Precursor Proliferation in Response to FGF

To analyze the contribution of GDNF through its GPI receptor GFR α 1 to the inhibition of cortical precursor proliferation, the neurosphere assay was performed using cortical progenitors derived from control or GFR α 1 mutant mice. To this end, we used mice carrying a cassette composed of floxed *Gfra1* cDNA followed by *GFP* cDNA. In these mice, Cre-mediated excision of floxed *Gfra1* cDNA converts the floxed *Gfra1* allele into a *GFP* reporter allele ([Uesaka et al., 2007](#)). We generated conditional *Gfra1*-mutant mice by breeding *Emx1-Cre* ones ([Iwasato et al., 2000](#); [Weisstaub et al., 2006](#)) with mice carrying floxed *Gfra1* alleles. Because *Emx1* is expressed in glutamatergic neuronal progenitors, *Emx1-Cre:Gfra1^{fllox/fllox}* mice (conditional *Gfra1* mutants) represent a model in which *Gfra1* is deleted from glutamatergic precursors ([Gorski et al., 2002](#); [Irala et al., 2016](#)). To analyze the contribution of GDNF/GFR α 1 to cortical precursor development, we used *Gfra1*-mutant mice homozygous for *Gfra1* deletion, *Emx1-Cre:GFR α ^{fllox/fllox}*, and heterozygous *Emx1-Cre:GFR^{fllox/+}* as control. There was a significant decrease in the number of neurospheres formed from E13.5 cortical precursors derived from control *Emx1-Cre:GFR α 1^{fllox/+}* when they were grown in the presence of FGF2 plus GDNF compared with the neurospheres grown in the presence of FGF2 alone. However, cortical precursor cells derived from animals deficient in GFR α 1 gave rise to an increased number of spheres compared with the ones derived from control animals, independently of the presence of GDNF ([Figures 4A and 4B](#)). In addition, the average diameter of the spheres derived from GFR α 1 mutant mice was bigger than the diameter of the spheres derived from control animals. The average diameter of the spheres derived from GFR α 1-deficient mice cortical progenitors grown in the presence of FGF2 was similar to the spheres grown in the presence of FGF2 plus GDNF ([Figures 4A and 4B](#)). The absence of GFR α 1 expression in the neocortex of *Emx1-Cre:GFR α ^{fllox/fllox}* mice was controlled by immunofluorescence (see [Figure S4](#)). Although GFR α 1 is the main binding receptor for GDNF, this neurotrophic factor can also bind with lower affinity to GFR α 2. However, the low levels of

Gfra2 mRNA expression during the early cortical development ([Figure 1A](#)), together with the complete absence of effect of GDNF in the neurosphere assay from cells coming from mice lacking *Gfra1*, suggested that GDNF specifically inhibits FGF2-induced cortical precursor proliferation acting through GFR α 1.

We also analyzed, by qPCR, the expression levels of cell-cycle regulatory genes in neurospheres derived from GFR α 1 mutant mice compared with control mice. In neurospheres derived from GFR α 1-deficient mice, the mRNA expression level of the cell-cycle promoter *CyclinD1* increased significantly compared with the levels detected in neurospheres derived from control animals in the presence of FGF2. In contrast, the mRNA expression level of cell-cycle inhibitor *p27* was decreased ([Figure 4C](#)). These results indicate that GDNF, through its receptor GFR α 1, negatively controls the proliferation of neural precursor cortical progenitors.

To analyze the requirement of the GDNF/GFR α 1 system in the proliferation of neuronal precursors *in vivo*, we performed immunofluorescence against Ki67 in cortical sections of GFR α 1-deficient or control mice at E13.5. The expression of this proliferation marker was analyzed in the different layers of the cortex, the high proliferative layer, VZ/SVZ (where the apical progenitors resides); the intermediate layer, IZ ([intermediate zone] populated mostly by basal progenitors); and the cortical plate where the neurons generated in this period are located ([Molyneaux et al., 2007](#)). In agreement with the *in vitro* results, a significant increase in Ki67-expressing cells was observed in GFR α 1-deficient animal sections, both in the VZ/SVZ and IZ regions, suggesting that GFR α 1 is required *in vivo* for a correct regulation of the proliferation of apical and basal neuronal progenitors ([Figures 4D and 4E](#)).

GDNF and Its Receptor GFR α 1 Are Involved in Neuronal Differentiation of Cortical Precursors

As we mentioned earlier, GDNF counteracts the mitogenic stimulation of FGF2 on cortical progenitors promoting neuronal differentiation *in vitro*. To characterize the contribution of GDNF on neuronal differentiation, cortical progenitors were cultured in the presence of a low concentration of FGF2 (5 ng/mL), with or without GDNF, followed by staining with the neuronal marker β III-tubulin and glutamatergic (VGlut, glutamate vesicular transporter) or GABAergic (VGAT, GABA vesicular transporter) markers. We observed that, under low concentrations of FGF2, GDNF induced an increase in the VGlut⁺ neuronal population. Few VGAT⁺ cells were detected in the cultures, and no difference was detected in their number in the presence or absence of GDNF ([Figures 5A and 5B](#)). Then, we analyzed whether GDNF treatment had any effect on neurite development of postmitotic neurons. The complexity of the β III-tubulin cells was




Figure 4. The Complex GDNF/GFR α 1 Is Required for Proper Cortical Progenitor Proliferation

(A) Neurospheres from cortical precursor cells derived from GFR α 1-deficient (*Emx1-Cre:GFR α 1^{fllox/fllox}*) or control mice (*Emx1-Cre:GFR α 1^{fllox/+}*) were grown for 5 DIV in the presence or absence of GDNF plus FGF2. Scale bar, 200 μ m.

(B) The bar graphs show the quantification of the number and diameter of neurospheres that were generated as described in (A). The results are expressed as the average of three independent experiments \pm SEM. * $p < 0.05$, *** $p < 0.001$ (ANOVA, followed by a Newman-Keuls test).

(C) Analysis of *Cyclin D1* and *p21* mRNA from neurospheres derived from GFR α 1-deficient (*Emx1-Cre:GFR α 1^{fllox/fllox}*) or control mice (*Emx1-Cre:GFR α 1^{fllox/+}*) cultured in the presence of FGF2, by real-time PCR. The expression of the specific mRNA was normalized to *Tbp*. Shown are mean \pm SEM of triplicate determinations, *** $p < 0.001$ (Student's *t* test).

(D) Ki67 immunostaining in coronal sections of E13.5 mice cortex from GFR α 1-deficient (*Emx1-Cre:GFR α 1^{fllox/fllox}*) or control (*Emx1-Cre:GFR α 1^{fllox/+}*) mice. VZ, ventricular zone; SVZ, subventricular zone, CP, cortical plate. Scale bar, 100 μ m.

(E) Quantitative analysis showing the percentage of Ki67⁺ cells in GFR α 1-deficient (*Emx1-Cre:GFR α 1^{fllox/fllox}*) respect to control (*Emx1-Cre:GFR α 1^{fllox/+}*) mice in the VZ/SVZ/IZ. The results are shown as mean \pm SEM of independent determinations in three separate mice of each genotype. * $p < 0.05$, ** $p < 0.005$ (Student's *t* test).

evaluated by Sholl analysis, which quantifies the number of neuritic branches intersecting concentric circles of increasing radius centering the reference point on the cell body (Sholl, 1953). Morphological parameters, such as total neurite length and branching points, were evaluated (Figures 5C–5E). We found that postmitotic neurons treated with GDNF showed a significant increase in neuritic arbor development compared with neurons non-treated with exogenous GDNF (Figures 5C–5E and S2D–S2F). The morphological effect on postmitotic neurons was also demonstrated by staining the cells with the dendrite marker MAP2 (see Figure S5). The effect of GDNF on these cells was observed in the presence of low concentrations of FGF2, but was abrogated under high concentrations of this growth factor (data not

shown), indicating that GDNF promotes morphological neuronal differentiation of cortical precursors depending on the levels of GFR α 1 expression modulated by FGF2.

Differentiation during brain development involves not only the growth of neurites but also the correct establishment of synaptic connections. Because of this, we decided to evaluate whether GDNF could affect the ability of postmitotic neurons to form dendritic spines, where the majority of excitatory synapses are established. A significant increase in spine density was observed in cells maintained in the presence of GDNF and low concentration of FGF2 relative to cultures maintained only in the presence of FGF2 (Figure 5F). These results collectively are in agreement with a role of GDNF as an inhibitor of cortical precursor proliferation during early development and as a




Figure 5. GDNF Promotes Neuronal Differentiation in Cortical Precursors

(A) Representative pictures of dissociated rat cortical progenitors cultured in the absence or presence of GDNF (100 ng/mL) plus FGF2 (5 ng/mL). After 5 DIV, the cells were fixed and stained with antibodies against β III-tubulin (green) and VGlut (blue) or VGAT (red). Arrows indicate individual cells showing VGAT/ β III-tubulin or VGlut/ β III-tubulin co-expression. Scale bar, 100 μ m.

(B) Quantitative analysis of the percent of VGlut⁺ or VGAT⁺ postmitotic neurons from the experiment described in (A). Shown are averages \pm SEM of three experiments, * p < 0.05 (Student's t test).

(C) Representative inverted images of rat cortical precursors maintained for 5 DIV in the absence or presence of GDNF (100 ng/mL) plus FGF2 (5 ng/mL) stained with anti- β III-tubulin. Scale bar, 30 μ m. See also Figure S5.

(D) Quantification of neurite complexity by Sholl analysis of cortical precursors grown as described in (C). Graph bar represents the cumulative neurite crossings obtained by Sholl analysis. * p < 0.05 (Student's t test).

(E) Bar graphs show the quantification of total neuritic length and branching from cortical precursors maintained as described in (C) (see also Figure S2E). Data of (D) and (E) represent mean \pm SEM of n = 3 independent experiments. *** p < 0.001 (ANOVA, followed by a Newman-Keuls test).

(F) Representative confocal images showing dendritic spines from postmitotic neurons obtained from cortical progenitors treated as indicated in (C). Scale bar, 5 μ m. The graph bar shows the quantification of the number of total dendritic spines along 100 μ m of dendritic length of postmitotic neurons. Data represent mean \pm SEM of n = 3 independent experiments. The arrowheads indicate dendritic spines. *** p < 0.001 (Student's t test).

promoter of neuronal specification and morphological differentiation.

GFR α 1 Is Required for Proper Glutamatergic Neuronal Differentiation

To analyze the contribution of GDNF/GFR α 1 to neuronal differentiation we cultured cortical progenitors derived

from GFR α 1 mutant mice *Emx1-Cre:GFR α 1^{fllox/fllox}*, in which Cre-mediated excision of floxed *GFR α 1* cDNA converts the floxed GFR α 1 allele into a GFP reporter (*Emx1-Cre:GFR α 1^{GFP/GFP}*). We performed cultures from GFR α 1-deficient mice homozygous for GFP, *Emx1-Cre:GFR α 1^{fllox/fllox}* (*Emx1-Cre:GFR α 1^{GFP/GFP}*) and heterozygous, *Emx1-Cre:GFR α 1^{fllox/+}* (*Emx1-Cre:GFR α 1^{+GFP}*) as control, in the




Figure 6. The Complex GDNF/GFRα1 Is Required for Neuronal Differentiation from Cortical Progenitors

(A) Representative pictures of dissociated rat cortical progenitors obtained from GFRα1-deficient (*Emx1-Cre:GFRα1^{flox/flox}*, *Emx1-Cre:GFRα1^{GFP/GFP}*) and control (*Emx1-Cre:GFRα1^{flox/+}*, *Emx1-Cre:GFRα1^{GFP/+}*) mice cultured in the presence of GDNF (100 ng/mL) plus FGF2 (5 ng/mL). After 5 DIV the cells were stained with antibodies against βIII-tubulin (red) and GFP. Scale bar, 30 μm.

(B) Quantification of dendrite complexity by Sholl analysis of postmitotic GFP⁺ neurons described in (A) maintained in the indicated conditions. Graph bar represents the cumulative neurite crossings obtained by Sholl analysis.

(C) Bar graphs show the quantification of total neuritic length and branching from cortical precursors obtained from GFRα1-deficient (*Emx1-Cre:GFRα1^{flox/flox}*, *Emx1-Cre:GFRα1^{GFP/GFP}*) and control (*Emx1-Cre:GFRα1^{flox/+}*, *Emx1-Cre:GFRα1^{GFP/+}*) mice maintained in the indicated conditions. Data of (B) and (C) represent mean ± SEM of n = 3 independent experiments. **p < 0.001, ***p < 0.0001; ns, non-significant (ANOVA, followed by a Newman-Keuls test).

(D) GFRα1 deletion results in a reduction of cortical dendrite development *in vivo*. Representative images of cortical neurons from P0 GFRα1-deficient and control littermate mice. Scale bars, 500 μm (left panel) and 100 μm (right panel). Bar graph shows the quantification of total dendritic length of cortical neurons from GFRα1-deficient (*Emx1-Cre:GFRα1^{flox/flox}*, *Emx1-Cre:GFRα1^{GFP/GFP}*) versus control (*Emx1-Cre:GFRα1^{flox/+}*, *Emx1-Cre:GFRα1^{GFP/+}*) mice. The results shown are mean ± SEM, *p < 0.05 (Student's t test). Quantifications were performed in neurons from n = 3 GFRα1-deficient (*Emx1-Cre:GFRα1^{flox/flox}*, *Emx1-Cre:GFRα1^{GFP/GFP}*) and control (*Emx1-Cre:GFRα1^{flox/+}*, *Emx1-Cre:GFRα1^{GFP/+}*) littermate mice.

presence of GDNF and low concentrations of FGF2 (Figure 6A). No difference in the number of βIII-tubulin⁺ cells was observed in cultures derived from GFRα1-deficient mice compared with the control mice. Morphological parameters such as total neuritic length and branching were also evaluated. We found a significant reduction in total neurite complexity as demonstrated by Sholl analysis,

and length and branchpoint number, in neurons derived from GFRα1 mutant mice compared with cells obtained from control mice when they were maintained in the presence of GDNF, indicating that GDNF/GFRα1 is required for proper neuronal development (Figures 6A–6C).

Finally, we analyzed the morphological features of GFRα1-deficient neurons in the postnatal cortex of GFRα1



mutant mice. Although GFR α 1 is expressed at very low levels in the majority of postnatal cortical neurons, a subpopulation of neurons from the cingulate cortex showed a clear GFP expression in transgenic *Emx1:Cre-GFR α 1^{fllox/fllox}* (*Emx1:Cre-GFR α 1^{GFP/GFP}*) or *Emx1:Cre:GFR α 1^{+fllox}* (*Emx1:Cre-GFR α 1^{GFP/+}*), indicating that GFR α 1 is expressed in these neurons. The complexity of the dendritic arbors was examined using an anti-GFP antibody, which allows visualization of individual cells and follows dendritic arbors on P0 cortical sections. We analyzed whether loss of GFR α 1 expression affects dendritic morphological features in GFP-expressing neurons. In control mice, a typical pyramidal morphology of these neurons was observed with a single apical dendrite extending toward the pia and secondary branches. In GFR α 1 mutant mice *Emx1:Cre-GFR α 1^{fllox/fllox}*, the dendrite morphology was abnormal, a significant reduction in the total length of apical dendritic arbors was observed, suggesting that GDNF/GFR α 1 signaling may be required *in vivo* for normal dendritic elaboration (Figure 6D). Taken together, these results indicate that the GDNF/GFR α 1 system is involved in the neuronal differentiation of glutamatergic neurons from cortical precursors.

DISCUSSION

During embryogenesis, neural precursor cells (NPCs) in the developing cortex are exposed to different intrinsic and extrinsic cues, the expression of which is under strict spatial and temporal control. External cues include growth factors such as FGF2, epidermal growth factor, Wnts, and neurotrophins, neurotransmitters such as GABA, glutamate, acetylcholine, and PACAP (Antonopoulos et al., 1997; Bartkowska et al., 2007; Haydar et al., 2000; Li et al., 2001; LoTurco et al., 1995; Ma et al., 2000), and contact-mediated signals, such as Notch (Chambers et al., 2001; Shen et al., 2002).

In the present work, we described the expression of *Gdnf* mRNA during early cortical developmental stages and provided evidence indicating that GDNF counteracts FGF2 self-renewal activity on NSCs, promoting the neural differentiation. Our data show that the GDNF receptor, *Gfra1*, mRNA is expressed during cortical development, increasing between E12 and E18, while the levels of the main FGF2 receptor in the CNS, FGFR1, decrease during this period, indicating a negative correlation between the expression of the two receptors. While *Fgfr1* is highly expressed in proliferating cultures of NPCs maintained in the presence of FGF2, *Gfra1* is mainly expressed in non-proliferating cultures maintained in the absence of FGF2. Using both adherent and neurosphere cell cultures, we found that GDNF acts by restricting NPC division in the presence of a mitogenic stimulus. Our work provides

evidence showing that GDNF reduces Akt activation and inhibits the expression of *Cyclin D1* and *E*, which are induced by FGF2, and promotes the expression levels of cell-cycle inhibitors, such as *p27* and *p21*. Selective ablation of GFR α 1 in glutamatergic cortical precursors results in an increase in the number of cortical precursors positive for the proliferation marker Ki67 *in vivo*. In agreement with this, we obtained more and bigger neurospheres from GFR α 1-deficient mice than the ones obtained from control animals. The decrease in proliferation induced by GDNF/GFR α 1 is due in part to an increase in neuronal differentiation of the NPCs and to maintaining stem cells in a resting state. Addition of GDNF enhances the number of postmitotic neurons on cultures of cortical precursor cells. However, although our experiments indicate that GDNF can influence neuronal differentiation, it is likely that other factors can contribute to the process as well, since animals with target disruptions of GFR α 1 in glutamatergic precursors do not have a clear deficit in neuronal number in the postnatal cortex or hippocampus (Irala et al., 2016). Although it is possible that other factors can compensate for the loss of GFR α 1, it could also be possible that the deficit in GFR α 1 could delay neuronal differentiation. In addition to promoting the withdrawal of precursor cells from cell cycle and inducing an increase in postmitotic neurons, GDNF also regulates dendrite development of these neurons. In agreement with a previous report (Irala et al., 2016) in which we described that GDNF through GFR α 1 promotes dendrite arborization and neurite complexity in hippocampal pyramidal cells, in the present work we show that the addition of GDNF to cultures of NPCs induced an increase in dendrite length and complexity, as well in dendritic spine-like protrusion, even in the presence of FGF2. These results revealed an important role of GDNF and GFR α 1 in hippocampal and cortical connectivity. Recently it has been described that glutamatergic cortical neurons from patients with Alzheimer's disease (AD) lacked a response to GDNF. In that work, the authors showed evidence indicating that cultured cortical neurons and postmortem brain tissue from patients with AD have lower levels of GFR α 1 receptors compared with neurons derived from normal brains (Konishi et al., 2014). Based on all these data it will be important to continue investigating the contribution of GDNF/GFR α 1 complex in neurodevelopmental disease characterized by cognitive impairments.

GDNF in Cortical Development

Previous reports have described a role for GDNF in GABAergic neuron migration and differentiation during early cortical development. These reports indicate that GDNF promotes differentiation and tangential migration of cortical GABAergic neuroblast from the ganglionic



eminence toward the cortex (Pozas and Ibanez, 2005; Bepalov et al., 2011). In the present work, we described a role of GDNF as a differentiation factor of glutamatergic cortical precursors acting through its receptor GFR α 1. In agreement with our results, a recent work using a transcriptoma profiling comparing ligands produced by NPCs and neurons, identified GDNF and another member of the GDNF family, NRTN, as proneurogenic factors. The authors claim that these effects are mediated by the tyrosine kinase receptor, RET; however, we could not detect the expression of this receptor by RT-PCR (Figure 1), immunostaining or immunoprecipitation (Figure S1). To define the mechanism involved in GDNF effect on cortical development, we consider that it will be necessary to perform the *in vivo* experiments in RET-deficient mice.

In the present work, we described that GDNF, through GFR α 1, promotes dendritic growth and a complexity of glutamatergic neurons derived from cortical precursors. Although GFR α 1 is highly expressed in hippocampal neurons, the expression of GFR α 1 in the postnatal cortex is restricted to a small population of cortical neurons. Analysis of newborn conditional GFR α 1-deficient mice shows a reduction in dendritic length in a subpopulation of cingulate cortical neurons. This result is in agreement with our previous results indicating that the GDNF/GFR α 1 complex plays a crucial role in the development of dendritic arbors and establishment of excitatory synaptic contacts. In hippocampal neurons these effects are mediated by the neural cell adhesion molecules NCAM-180 (Irala et al., 2016).

GDNF as a Proliferative or Differentiation Factor

Interestingly, GDNF has been described as a promotor of proliferation in different systems such as enteric progenitors and spermatogonial stem cells (SSCs) (Meng et al., 2000; Sasselli et al., 2012; Yan et al., 2004). In these systems, the GDNF proliferative effects depend on tyrosine kinase receptor, RET, signaling. Addition of GDNF to enteric neuron precursors, which express RET, increases the number of neurons and enteric glia. Furthermore, it has also been described that GDNF signaling is indispensable for SSC self-renewal by binding to the GFR α 1/RET receptor. In this system, GDNF and FGF2 induce SSC self-renewal by different molecular mechanisms (Takashima et al., 2015).

Our present results indicate that GDNF inhibits the proliferation of cortical precursors. The mechanism by which GDNF suppress NPCs proliferation it is not clear yet. Our findings indicate that GDNF antagonizes the proliferative effect of FGF by inhibiting intracellular signaling, such as the Akt pathway, which leads to increased levels of the cell-cycle inhibitor p27 and reduction in the levels of the cyclin D1 expression, as well as downregulation of FGFR1

expression. We provide evidence indicating that this effect could be mediated by GDNF signaling acting through GFR α 1 and NCAM receptors. Recently, NCAM and NCAM-derived peptides have been reported to interact with FGFRs (Christensen et al., 2006; Francavilla et al., 2009; Kiselyov et al., 2003), modulating FGFR-mediated cellular functions (Hansen et al., 2008). Interestingly, it has been demonstrated that NCAM expression reduces FGF-stimulated cell proliferation in fibroblasts (Francavilla et al., 2007). Thus, one possibility is that GDNF acting through its receptors GFR α 1 and NCAM could interfere FGFR signaling in NPCs.

The proliferation of cortical precursor cells appears to be regulated by FGF during early cortical development in the ventricular zone, when FGF2 prevents NPC differentiation. Subsequently, GDNF acts on precursor cells inducing cell-cycle withdrawal and neuronal differentiation. Later in the development GDNF acts on postmitotic neurons promoting dendrite complexity. The dual effect of GDNF depends on the levels of GFR α 1 expression. Our findings show that the expression of GFR α 1 is regulated by FGF2 treatment in cortical precursors, while the expression of FGFR1 is modulated by GDNF. Thus, during cortical development, the levels of FGF2 decrease downregulating the expression of its receptor FGFR1 and upregulating the levels of GFR α 1 in postmitotic neurons. The absence of RET in cortical postmitotic neurons suggests that the effect on neurite complexity could be mediated by NCAM signaling, as we have described previously in CA1 and CA3 pyramidal neurons (Irala et al., 2016). The evidence obtained during the last years about the role of GDNF on hippocampal and cortical neurons gives new opportunities to study the role of GDNF in neurodevelopmental diseases characterized by cognitive deficits.

EXPERIMENTAL PROCEDURES

Transgenic Mice

Gfr α 1^{flx/flx} mice were generously provided by Dr J. Milbrandt (Washington University School of Medicine, St Louis, MO, USA) (Uesaka et al., 2007). *Gfr α 1^{flx/flx}* mice were mated with *Emx1-Cre* mice generously provided by Dr N. Weisstaub (School of Medicine, University of Buenos Aires, Argentina) (Iwasato et al., 2000; Weisstaub et al., 2006). All transgenic strains were genotyped by PCR. PCR primer sequences are available upon request. The use of animals was approved by the Animal Care and Use Committee of the School of Medicine, University of Buenos Aires, ethical permit number 67341/2013.

PCR and Real-Time PCR

PCR and real-time PCR were performed using standard methodologies. See [Supplemental Experimental Procedures](#) for details.



DNA Constructs, Cell Transfection, and Recombinant Proteins

Details of DNA constructs, cell transfection, and recombinant proteins used in the work can be found in [Supplemental Experimental Procedures](#).

Primary Cortical Progenitor Cultures and Neurosphere Assay

Rat and mouse cortical progenitor cells were isolated from E14.5 Wistar rats and E13.5 C57/BL6 mice, respectively (School of Medicine, University of Buenos Aires, Argentina) to perform primary cultures. Details about the cultures can be found in [Supplemental Experimental Procedures](#).

Immunostaining and Confocal Microscopy

For immunofluorescence, cells were washed, fixed with 4% paraformaldehyde (PFA) in PBS, permeabilized with 0.3% Triton X-100, blocked with 10% donkey normal serum (Jackson ImmunoResearch) in PBS, and incubated overnight at 4°C with the indicated antibodies (see [Supplemental Experimental Procedures](#)).

For tissue immunofluorescence assays, rat and mouse brains were isolated from animals fixed with 4% PFA for 4 hr, maintained in sucrose 30% in PBS overnight, and then embedded in O.C.T. (Tissue-Tek) and sectioned at 25 μm . Cryostat sections were permeabilized with 0.1% Triton X-100, blocked with 10% donkey normal serum, and incubated overnight at 4°C with primary antibodies. The primary antibodies used for these stainings are described in [Supplemental Experimental Procedures](#). The nuclear marker DAPI (1/10000, Sigma) was used to stain cells in culture and tissue sections. The secondary antibodies were from Jackson ImmunoResearch.

After immunostaining, microscopy was performed using an Olympus IX-81 inverted microscope or an Olympus FV-1000 confocal microscope, using identical settings between control and experimental images.

Total Cell Lysates and Western Blot

Cell cultures were lysed at 4°C in TNE buffer (25 mM Tris-ClH [pH 7.4]; 1 mM EDTA; 137 mM NaCl) containing 0.5% Triton X-100, plus protease and phosphatase inhibitors. Proteins lysates were clarified by centrifugation and analyzed by western blotting as described previously ([Paratcha et al., 2003](#)). The antibodies used for western blotting are described in [Supplemental Experimental Procedures](#). The blots were scanned in a Storm 845 PhosphorImager (GE Healthcare Life Sciences), and quantifications were done with ImageQuant software (GE Healthcare Life Sciences). Numbers below the lanes indicate fold of induction relative to control normalized to total levels of target protein.

Image Analysis and Quantification

Neuritic growth assays for neurons derived from cortical precursors were performed in primary E14.5 rat or E13.5 mouse dissociated cultures. Images of dissociated neurons to perform the morphological analyzes were obtained using an Olympus IX-81 microscope, in a 20 \times objective. Neuritic complexity measurements were per-

formed, as described in [Supplemental Experimental Procedures](#), on neuronal cells that showed a pyramidal type morphology, containing a principal dendrite and several fine dendrites.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism. Details are provided in each figure legend. For quantification of *in vitro* or *in vivo* assays, 10–30 neurons were chosen randomly per condition in each experiment from at least three independent experiments. The results are shown as mean \pm SEM as indicated in the figure legends, and significance was accepted at $p < 0.05$. Statistical significance was calculated using two-tailed Student's *t* test or ANOVA followed by *post hoc* multiple comparison test as indicated.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures and five figures and can be found with this article online at <https://doi.org/10.1016/j.stemcr.2018.01.019>.

AUTHOR CONTRIBUTIONS

A.B., G.P., and F.L. designed the experiments. A.B. and P.A.F. performed the experiments. A.B., P.A.F., G.P., and F.L. analyzed and discussed the experiment results. F.L. wrote the manuscript.

ACKNOWLEDGMENTS

We thank Dr. N. Weisstaub and Dr. J. Milbrand for sharing mice, A. Pecile and M. Ponce for animal care, D. Macaria and F.C. Alsina for technical help, Roux-Ocefa for reagent supply, and Innova-t and UBATEC for research grant administration. Financial support was obtained from the Argentine Agency for Promotion of Science and Technology (ANPCyT, PICT-PROBITEC-2011-2731, PICT-2014-2155, PICTV-2015-3814) and from the University of Buenos Aires (UBACyT-2014-2017, 20020130100779BA). G.P. and F.L. were supported by an Independent Research Career Position from the Argentine Medical Research Council (CONICET). A.B. was supported by a fellowship from ANPCyT and CONICET and P.A.F. by a fellowship from CONICET.

Received: June 17, 2017

Revised: January 17, 2018

Accepted: January 18, 2018

Published: February 22, 2018

REFERENCES

- Anderson, S.A., Eisenstat, D.D., Shi, L., and Rubenstein, J.L. (1997). Interneuron migration from basal forebrain to neocortex: dependence on *Dlx* genes. *Science* 278, 474–476.
- Anderson, S.A., Kaznowski, C.E., Horn, C., Rubenstein, J.L., and McConnell, S.K. (2002). Distinct origins of neocortical projection neurons and interneurons *in vivo*. *Cereb. Cortex* 12, 702–709.
- Andreu, Z., Khan, M.A., Gonzalez-Gomez, P., Negueruela, S., Hortiguera, R., San Emeterio, J., Ferron, S.R., Martinez, G., Vidal, A., Farinas, I., et al. (2015). The cyclin-dependent kinase inhibitor p27 *kip1* regulates radial stem cell quiescence and neurogenesis in the adult hippocampus. *Stem Cells* 33, 219–229.



- Antonopoulos, J., Pappas, I.S., and Parnavelas, J.G. (1997). Activation of the GABAA receptor inhibits the proliferative effects of bFGF in cortical progenitor cells. *Eur. J. Neurosci.* *9*, 291–298.
- Arvidsson, A., Kokaia, Z., Airaksinen, M.S., Saarma, M., and Lindvall, O. (2001). Stroke induces widespread changes of gene expression for glial cell line-derived neurotrophic factor family receptors in the adult rat brain. *Neuroscience* *106*, 27–41.
- Barnabe-Heider, F., and Miller, F.D. (2003). Endogenously produced neurotrophins regulate survival and differentiation of cortical progenitors via distinct signaling pathways. *J. Neurosci.* *23*, 5149–5160.
- Bartkowska, K., Paquin, A., Gauthier, A.S., Kaplan, D.R., and Miller, F.D. (2007). Trk signaling regulates neural precursor cell proliferation and differentiation during cortical development. *Development* *134*, 4369–4380.
- Bespalov, M.M., Sidorova, Y.A., Tumova, S., Ahonen-Bishopp, A., Magalhaes, A.C., Kuleskiy, E., Paveliev, M., Rivera, C., Rauvala, H., and Saarma, M. (2011). Heparan sulfate proteoglycan syndecan-3 is a novel receptor for GDNF, neurturin, and artemin. *J. Cell Biol.* *192*, 153–169.
- Borello, U., Cobos, I., Long, J.E., McWhirter, J.R., Murre, C., and Rubenstein, J.L. (2008). FGF15 promotes neurogenesis and opposes FGF8 function during neocortical development. *Neural Dev.* *3*, 17.
- Burazin, T.C., and Gundlach, A.L. (1999). Localization of GDNF/neurturin receptor (c-ret, GFRalpha-1 and alpha-2) mRNAs in postnatal rat brain: differential regional and temporal expression in hippocampus, cortex and cerebellum. *Brain Res. Mol. Brain Res.* *73*, 151–171.
- Burdon, T., Smith, A., and Savatier, P. (2002). Signalling, cell cycle and pluripotency in embryonic stem cells. *Trends Cell Biol.* *12*, 432–438.
- Bystron, I., Blakemore, C., and Rakic, P. (2008). Development of the human cerebral cortex: Boulder Committee revisited. *Nat. Rev. Neurosci.* *9*, 110–122.
- Canty, A.J., Dietze, J., Harvey, M., Enomoto, H., Milbrandt, J., and Ibanez, C.F. (2009). Regionalized loss of parvalbumin interneurons in the cerebral cortex of mice with deficits in GFRalpha1 signaling. *J. Neurosci.* *29*, 10695–10705.
- Chambers, C.B., Peng, Y., Nguyen, H., Gaiano, N., Fishell, G., and Nye, J.S. (2001). Spatiotemporal selectivity of response to Notch1 signals in mammalian forebrain precursors. *Development* *128*, 689–702.
- Charoy, C., Nawabi, H., Reynaud, F., Derrington, E., Bozon, M., Wright, K., Falk, J., Helmbacher, F., Kindbeiter, K., and Castellani, V. (2012). Gdnf activates midline repulsion by Semaphorin3B via NCAM during commissural axon guidance. *Neuron* *75*, 1051–1066.
- Cholfin, J.A., and Rubenstein, J.L. (2007). Patterning of frontal cortex subdivisions by Fgf17. *Proc. Natl. Acad. Sci. USA* *104*, 7652–7657.
- Christensen, C., Lauridsen, J.B., Berezin, V., Bock, E., and Kiselyov, V.V. (2006). The neural cell adhesion molecule binds to fibroblast growth factor receptor 2. *FEBS Lett.* *580*, 3386–3390.
- Ford-Perriss, M., Abud, H., and Murphy, M. (2001). Fibroblast growth factors in the developing central nervous system. *Clin. Exp. Pharmacol. Physiol.* *28*, 493–503.
- Francavilla, C., Cattaneo, P., Berezin, V., Bock, E., Ami, D., de Marco, A., Christofori, G., and Cavallaro, U. (2009). The binding of NCAM to FGFR1 induces a specific cellular response mediated by receptor trafficking. *J. Cell Biol.* *187*, 1101–1116.
- Francavilla, C., Loeffler, S., Piccini, D., Kren, A., Christofori, G., and Cavallaro, U. (2007). Neural cell adhesion molecule regulates the cellular response to fibroblast growth factor. *J. Cell Sci.* *120*, 4388–4394.
- Fukuchi-Shimogori, T., and Grove, E.A. (2001). Neocortex patterning by the secreted signaling molecule FGF8. *Science* *294*, 1071–1074.
- Garel, S., Huffman, K.J., and Rubenstein, J.L. (2003). Molecular regionalization of the neocortex is disrupted in Fgf8 hypomorphic mutants. *Development* *130*, 1903–1914.
- Glazner, G.W., Mu, X., and Springer, J.E. (1998). Localization of glial cell line-derived neurotrophic factor receptor alpha and c-ret mRNA in rat central nervous system. *J. Comp. Neurol.* *391*, 42–49.
- Golden, J.P., DeMaro, J.A., Osborne, P.A., Milbrandt, J., and Johnson, E.M., Jr. (1999). Expression of neurturin, GDNF, and GDNF family-receptor mRNA in the developing and mature mouse. *Exp. Neurol.* *158*, 504–528.
- Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L., and Jones, K.R. (2002). Cortical excitatory neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage. *J. Neurosci.* *22*, 6309–6314.
- Gotz, M., Stoykova, A., and Gruss, P. (1998). Pax6 controls radial glia differentiation in the cerebral cortex. *Neuron* *21*, 1031–1044.
- Hansen, S.M., Kohler, L.B., Li, S., Kiselyov, V., Christensen, C., Owczarek, S., Bock, E., and Berezin, V. (2008). NCAM-derived peptides function as agonists for the fibroblast growth factor receptor. *J. Neurochem.* *106*, 2030–2041.
- Haydar, T.F., Wang, F., Schwartz, M.L., and Rakic, P. (2000). Differential modulation of proliferation in the neocortical ventricular and subventricular zones. *J. Neurosci.* *20*, 5764–5774.
- Hebert, J.M., and Fishell, G. (2008). The genetics of early telencephalon patterning: some assembly required. *Nat. Rev. Neurosci.* *9*, 678–685.
- Irala, D., Bonafina, A., Fontanet, P.A., Alsina, F.C., Paratcha, G., and Ledda, F. (2016). The GDNF-GFRalpha1 complex promotes the development of hippocampal dendritic arbors and spines via NCAM. *Development* *143*, 4224–4235.
- Iwasato, T., Datwani, A., Wolf, A.M., Nishiyama, H., Taguchi, Y., Tonogawa, S., Knopfel, T., Erzurumlu, R.S., and Itohara, S. (2000). Cortex-restricted disruption of NMDAR1 impairs neuronal patterns in the barrel cortex. *Nature* *406*, 726–731.
- Jin, L., Hu, X., and Feng, L. (2005). NT3 inhibits FGF2-induced neural progenitor cell proliferation via the PI3K/GSK3 pathway. *J. Neurochem.* *93*, 1251–1261.
- Kang, W., Wong, L.C., Shi, S.H., and Hebert, J.M. (2009). The transition from radial glial to intermediate progenitor cell is inhibited by FGF signaling during corticogenesis. *J. Neurosci.* *29*, 14571–14580.



- Kiselyov, V.V., Skladchikova, G., Hinsby, A.M., Jensen, P.H., Kula-hin, N., Soroka, V., Pedersen, N., Tsetlin, V., Poulsen, F.M., Berezin, V., et al. (2003). Structural basis for a direct interaction between FGFR1 and NCAM and evidence for a regulatory role of ATP. *Structure* *11*, 691–701.
- Konishi, Y., Yang, L.B., He, P., Lindhol, K., Lu, B., Li, R., and Shen, Y. (2014). Deficiency of GDNF receptor GFR α 1 in Alzheimer's neurons results in neuronal death. *J. Neurosci.* *34*, 13127–13138.
- Kowalczyk, T., Pontious, A., Englund, C., Daza, R.A., Bedogni, F., Hodge, R., Attardo, A., Bell, C., Huttner, W.B., and Hevner, R.F. (2009). Intermediate neuronal progenitors (basal progenitors) produce pyramidal-projection neurons for all layers of cerebral cortex. *Cereb. Cortex* *19*, 2439–2450.
- Lenhard, T., and Suter-Crazzolara, C. (1998). Developmental expression of GDNF, neurturin and their receptors in rat hippocampus. *Neuroreport* *9*, 2927–2931.
- Li, M., Price, D., and Smith, A. (2001). Lineage selection and isolation of neural precursors from embryonic stem cells. *Symp. Soc. Exp. Biol.*, 29–42.
- LoTurco, J.J., Owens, D.F., Heath, M.J., Davis, M.B., and Kriegstein, A.R. (1995). GABA and glutamate depolarize cortical progenitor cells and inhibit DNA synthesis. *Neuron* *15*, 1287–1298.
- Ma, W., Maric, D., Li, B.S., Hu, Q., Andreadis, J.D., Grant, G.M., Liu, Q.Y., Shaffer, K.M., Chang, Y.H., Zhang, L., et al. (2000). Acetylcholine stimulates cortical precursor cell proliferation in vitro via muscarinic receptor activation and MAP kinase phosphorylation. *Eur. J. Neurosci.* *12*, 1227–1240.
- Maric, D., Fiorio Pla, A., Chang, Y.H., and Barker, J.L. (2007). Self-renewing and differentiating properties of cortical neural stem cells are selectively regulated by basic fibroblast growth factor (FGF) signaling via specific FGF receptors. *J. Neurosci.* *27*, 1836–1852.
- Meng, X., Lindahl, M., Hyvonen, M.E., Parvinen, M., de Rooij, D.G., Hess, M.W., Raatikainen-Ahokas, A., Sainio, K., Rauvala, H., Lakso, M., et al. (2000). Regulation of cell fate decision of undifferentiated spermatogonia by GDNF. *Science* *287*, 1489–1493.
- Mira, H., Andreu, Z., Suh, H., Lie, D.C., Jessberger, S., Consiglio, A., San Emeterio, J., Hortiguera, R., Marques-Torres, M.A., Nakashima, K., et al. (2010). Signaling through BMPRII regulates quiescence and long-term activity of neural stem cells in the adult hippocampus. *Cell Stem Cell* *7*, 78–89.
- Molyneux, B.J., Arlotta, P., and Macklis, J.D. (2007). Molecular development of corticospinal motor neuron circuitry. *Novartis Found. Symp.* *288*, 3–15, discussion 15–20, 96–18.
- Ohkubo, Y., Uchida, A.O., Shin, D., Partanen, J., and Vaccarino, F.M. (2004). Fibroblast growth factor receptor 1 is required for the proliferation of hippocampal progenitor cells and for hippocampal growth in mouse. *J. Neurosci.* *24*, 6057–6069.
- Paratcha, G., Ibanez, C.F., and Ledda, F. (2006). GDNF is a chemoattractant factor for neuronal precursor cells in the rostral migratory stream. *Mol. Cell Neurosci.* *31*, 505–514.
- Paratcha, G., Ledda, F., and Ibanez, C.F. (2003). The neural cell adhesion molecule NCAM is an alternative signaling receptor for GDNF family ligands. *Cell* *113*, 867–879.
- Pozas, E., and Ibanez, C.F. (2005). GDNF and GFR α 1 promote differentiation and tangential migration of cortical GABAergic neurons. *Neuron* *45*, 701–713.
- Qian, X., Davis, A.A., Goderie, S.K., and Temple, S. (1997). FGF2 concentration regulates the generation of neurons and glia from multipotent cortical stem cells. *Neuron* *18*, 81–93.
- Raballo, R., Rhee, J., Lyn-Cook, R., Leckman, J.F., Schwartz, M.L., and Vaccarino, F.M. (2000). Basic fibroblast growth factor (Fgf2) is necessary for cell proliferation and neurogenesis in the developing cerebral cortex. *J. Neurosci.* *20*, 5012–5023.
- Rakic, P. (2006). A century of progress in corticogenesis: from silver impregnation to genetic engineering. *Cereb. Cortex* *16* (Suppl 1), i3–17.
- Rash, B.G., Lim, H.D., Breunig, J.J., and Vaccarino, F.M. (2011). FGF signaling expands embryonic cortical surface area by regulating Notch-dependent neurogenesis. *J. Neurosci.* *31*, 15604–15617.
- Saarma, M. (2000). GDNF - a stranger in the TGF-beta superfamily? *Eur. J. Biochem.* *267*, 6968–6971.
- Sasselli, V., Pachnis, V., and Burns, A.J. (2012). The enteric nervous system. *Dev. Biol.* *366*, 64–73.
- Shen, Q., Zhong, W., Jan, Y.N., and Temple, S. (2002). Asymmetric Numb distribution is critical for asymmetric cell division of mouse cerebral cortical stem cells and neuroblasts. *Development* *129*, 4843–4853.
- Sholl, D.A. (1953). Dendritic organization in the neurons of the visual and motor cortices of the cat. *J. Anat.* *87*, 387–406.
- Stevens, H.E., Smith, K.M., Maragnoli, M.E., Fagel, D., Borok, E., Shanabrough, M., Horvath, T.L., and Vaccarino, F.M. (2010). Fgfr2 is required for the development of the medial prefrontal cortex and its connections with limbic circuits. *J. Neurosci.* *30*, 5590–5602.
- Takashima, S., Kanatsu-Shinohara, M., Tanaka, T., Morimoto, H., Inoue, K., Ogonuki, N., Jijiwa, M., Takahashi, M., Ogura, A., and Shinohara, T. (2015). Functional differences between GDNF-dependent and FGF2-dependent mouse spermatogonial stem cell self-renewal. *Stem Cell Reports* *4*, 489–502.
- Trupp, M., Scott, R., Whittmore, S.R., and Ibanez, C.F. (1999). Ret-dependent and -independent mechanisms of glial cell line-derived neurotrophic factor signaling in neuronal cells. *J. Biol. Chem.* *274*, 20885–20894.
- Uesaka, T., Jain, S., Yonemura, S., Uchiyama, Y., Milbrandt, J., and Enomoto, H. (2007). Conditional ablation of GFR α 1 in post-migratory enteric neurons triggers unconventional neuronal death in the colon and causes a Hirschsprung's disease phenotype. *Development* *134*, 2171–2181.
- Vaccarino, F.M., Schwartz, M.L., Raballo, R., Nilsen, J., Rhee, J., Zhou, M., Doetschman, T., Coffin, J.D., Wyland, J.J., and Hung, Y.T. (1999a). Changes in cerebral cortex size are governed by fibroblast growth factor during embryogenesis. *Nat. Neurosci.* *2*, 848.
- Vaccarino, F.M., Schwartz, M.L., Raballo, R., Rhee, J., and Lyn-Cook, R. (1999b). Fibroblast growth factor signaling regulates growth and morphogenesis at multiple steps during brain development. *Curr. Top. Dev. Biol.* *46*, 179–200.



Weise, B., Janet, T., and Grothe, C. (1993). Localization of bFGF and FGF-receptor in the developing nervous system of the embryonic and newborn rat. *J. Neurosci. Res.* *34*, 442–453.

Weisstaub, N.V., Zhou, M., Lira, A., Lambe, E., Gonzalez-Maeso, J., Hornung, J.P., Sibille, E., Underwood, M., Itohara, S., Dauer, W.T., et al. (2006). Cortical 5-HT_{2A} receptor signaling modulates anxiety-like behaviors in mice. *Science* *313*, 536–540.

Yan, H., Bergner, A.J., Enomoto, H., Milbrandt, J., Newgreen, D.F., and Young, H.M. (2004). Neural cells in the esophagus respond to glial cell line-derived neurotrophic factor and neurturin, and are RET-dependent. *Dev. Biol.* *272*, 118–133.

Yuzwa, S.A., Yang, G., Borrett, M.J., Clarke, G., Cancino, G.I., Zahr, S.K., Zandstra, P.W., Kaplan, D.R., and Miller, F.D. (2016). Pro-neurogenic ligands defined by modeling developing cortex growth factor communication networks. *Neuron* *91*, 988–1004.