





ORIGINAL ARTICLE

Plant, Cell & Environment

WILEY

The two copies of the zinc and cadmium ZIP6 transporter of *Arabidopsis halleri* have distinct effects on cadmium tolerance

Julien Spielmann¹  | Hassan Ahmadi² | Maxime Scheepers¹ | Michael Weber²  | Sarah Nitsche² | Monique Carnol³ | Bernard Bosman³ | Juergen Kroymann⁴ | Patrick Motte¹ | Stephan Clemens²  | Marc Hanikenne¹ 

¹InBioS-PhytoSystems, Functional Genomics and Plant Molecular Imaging, University of Liège, Liège, Belgium

²Department of Plant Physiology, University of Bayreuth, Bayreuth, Germany

³InBioS-PhytoSystems, Laboratory of Plant and Microbial Ecology, University of Liège, Liège, Belgium

⁴CNRS, AgroParisTech, Ecologie Systématique et Evolution, Université Paris-Saclay, Orsay, France

Correspondence

Marc Hanikenne, InBioS-PhytoSystems, University of Liège, Quartier de la Vallée, 1, Chemin de la Vallée, 4-Bât B22, B4000 Liège, Belgium.

Email: marc.hanikenne@uliege.be

Stephan Clemens, Department of Plant Physiology, and Bayreuth Centre of Ecology and Environmental Research, University of Bayreuth, Universitätsstrasse 30, 95447 Bayreuth, Germany
Email: stephan.clemens@uni-bayreuth.de

Funding information

Belgian Federal Science Policy Office, Grant/Award Number: IAP no. P7/44; Deutsche Forschungsgemeinschaft, Grant/Award Numbers: SPP1529 ADAPTOMICS, Cl152/9-1, Cl152/9-2; Fonds De La Recherche Scientifique - FNRS, Grant/Award Numbers: PDR-T.0206.13, MIS-F.4511.16, CDR J.0009.17, PDR T0120.18; Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture; University of Liège, Grant/Award Number: SFRD-12/03

Abstract

Plants have the ability to colonize highly diverse environments. The zinc and cadmium hyperaccumulator *Arabidopsis halleri* has adapted to establish populations on soils covering an extreme range of metal availabilities. The *A. halleri* ZIP6 gene presents several hallmarks of hyperaccumulation candidate genes: it is constitutively highly expressed in roots and shoots and is associated with a zinc accumulation quantitative trait locus. Here, we show that *AhZIP6* is duplicated in the *A. halleri* genome. The two copies are expressed mainly in the vasculature in both *A. halleri* and *Arabidopsis thaliana*, indicative of conserved *cis* regulation, and acquired partial organ specialization. Yeast complementation assays determined that *AhZIP6* is a zinc and cadmium transporter. *AhZIP6* silencing in *A. halleri* or expression in *A. thaliana* alters cadmium tolerance, but has no impact on zinc and cadmium accumulation. *AhZIP6*-silenced plants display reduced cadmium uptake upon short-term exposure, adding *AhZIP6* to the limited number of Cd transporters supported by *in planta* evidence. Altogether, our data suggest that *AhZIP6* is key to fine-tune metal homeostasis in specific cell types. This study additionally highlights the distinct fates of duplicated genes in *A. halleri*.

KEYWORDS

Arabidopsis halleri, cadmium tolerance, gene duplication, hyperaccumulation, male sterility, sub-functionalization, zinc

1 | INTRODUCTION

All organisms have to maintain adequate metal ion homeostasis to ensure growth and development in varying environments. For sessile

organisms such as plants, this represents an even more complicated challenge. Plants have developed a sophisticated and tightly regulated metal homeostasis network of transporters and chelating molecules to ensure sufficient metal supply while avoiding their toxicity (Palmer & Guerinot, 2009; Ricachenevsky, Menguer, Sperotto, & Fett, 2015; Sinclair & Krämer, 2012). This allows plants to develop in

Julien Spielmann and Hassan Ahmadi contributed equally to this study.

environments that substantially differ in metal availability. Hence, plants employ two main strategies to colonize natural or anthropogenic metal-polluted soils (Baker, 1981). The majority of plant species are excluders, limiting metal absorption, exporting metal ions in excess out of roots and limiting root-to-shoot translocation to protect the aerial photosynthetically active tissues. In contrast, about 720 species have developed a so-called hyperaccumulation strategy, where large amounts of metals are stored in shoot tissues without toxicity symptoms (Hanikenne & Nouet, 2011; Krämer, 2010; Merlot, de la Torre, & Hanikenne, 2018; Reeves et al., 2018; Verbruggen, Hermans, & Schat, 2009). Among those, only a few are zinc and cadmium hyperaccumulators, with commonly accepted hyperaccumulation thresholds of $3,000 \mu\text{g g}^{-1}$ of zinc and $100 \mu\text{g g}^{-1}$ of cadmium in leaf dry weight (Krämer, 2010; van der Ent, Baker, Reeves, Pollard, & Schat, 2013). Hyperaccumulator plants are regarded as models to unveil adaptation to extreme environments, but also to reveal homeostatic mechanisms controlling metal distribution in plants (Hanikenne & Nouet, 2011; Krämer, 2010).

With *Noccaea caerulea* and two *Sedum* species, *Arabidopsis halleri* is a metal hyperaccumulator model (Merlot et al., 2018). It is a highly zinc and cadmium tolerant species and displays species-wide zinc hyperaccumulation, both in metallicolous (living on polluted soils) and in the more common non-metallicolous populations (Bert, Bonnin, Saumitou-Laprade, De Laguérie, & Petit, 2002; Stein et al., 2017). Cadmium hyperaccumulation is highly variable and restricted to a more limited number of *A. halleri* populations (Meyer et al., 2015; Stein et al., 2017). This suggests that zinc and cadmium hyperaccumulation rely on at least partially distinct pathways.

Physiologically, the hyperaccumulation process may be summarized to four main steps: (a) an active and efficient metal uptake from the soil, (b) an efficient radial transfer of metals to vascular tissues combined with low vacuolar storage in roots, (c) an enhanced xylem loading and root-to-shoot translocation and finally (d) an efficient detoxification process enabled by efficient shoot metal distribution and leaf vacuolar storage (Clemens, 2016; Hanikenne & Nouet, 2011; Merlot et al., 2018).

A. halleri genes with putative functions in these processes have been identified using quantitative trait locus (QTL) and/or cross-species transcriptomic analyses. Interspecific crosses between *A. halleri* and *Arabidopsis lyrata* ssp. *petraea*, a non-hyperaccumulator relative, were used to map QTLs for zinc tolerance (Willems et al., 2007) and accumulation (Frérot et al., 2010), as well as cadmium tolerance (Baliardini, Meyer, Salis, Saumitou-Laprade, & Verbruggen, 2015; Courbot et al., 2007) and accumulation (Willems et al., 2010), in back-cross 1 or F2 populations. Transcriptomics comparing *A. halleri* with *Arabidopsis thaliana* allowed the identification of about 50 metal homeostasis genes, encoding transporters or metal chelator synthesis enzymes, which are more highly expressed in *A. halleri*. High expression levels of a number of these genes, but not all (see Charlier et al., 2015) were linked to gene copy number expansion and/or cis-regulatory changes (Dräger et al., 2004; Fasani et al., 2017; Hanikenne et al., 2008; Shahzad et al., 2010; Suryawanshi et al., 2016). More recently, these approaches were also employed to

examine the molecular basis of intraspecific variation in the zinc and cadmium hyperaccumulation and hypertolerance traits in *A. halleri* (Corso et al., 2018; Karam et al., 2019; Schwartzman et al., 2018).

A number of candidate genes were commonly identified by the QTL and transcriptomics approaches (Hanikenne & Nouet, 2011; Krämer, 2010) but so far a direct contribution to hypertolerance and/or hyperaccumulation has been experimentally confirmed for only three of these genes by gene silencing in *A. halleri*. *HMA4* (HEAVY METAL ATPase 4) has a major contribution to both zinc and cadmium hypertolerance and hyperaccumulation (Hanikenne et al., 2008), whereas *NAS2* (Nicotianamine Synthase 2) contributes to zinc root-to-shoot translocation (Cornu et al., 2014; Deinlein et al., 2012; Uraguchi, Weber, & Clemens, 2019). Finally, *CAX1* (CATION/HYDROGEN EXCHANGER 1) is required to minimize cadmium-triggered oxidative damages at low calcium supply (Ahmadi, Corso, Weber, Verbruggen, & Clemens, 2018; Baliardini et al., 2015).

Among uncharacterized candidate genes is *ZIP6* (ZRT-IRT-like PROTEIN 6). It is 9- and 24-fold more highly expressed in roots and shoots of *A. halleri* compared with *A. thaliana*, respectively (Becher, Talke, Krall, & Krämer, 2004; Talke, Hanikenne, & Krämer, 2006) and is associated with a QTL for zinc accumulation in the presence of cadmium (Willems et al., 2010). It may be present in multiple copies in the *A. halleri* genome (Suryawanshi et al., 2016; Talke et al., 2006). Moreover, it was found as highly expressed in roots and shoots of zinc hyperaccumulating F3 progenies of a *A. halleri* and *A. lyrata* ssp. *petraea* cross (Filatov et al., 2006). *ZIP6* is also highly expressed in shoots of *N. caerulea*, another zinc and cadmium hyperaccumulator (Hammond et al., 2006; Krämer, Talke, & Hanikenne, 2007) and, interestingly, it was part of a list of 12 genes whose expression correlated with cadmium accumulation among *N. caerulea* populations (Halimaa et al., 2014; Wu et al., 2009).

ZIP6 belongs to the ZIP family of divalent metal cation transporters (Guerinot, 2000). ZIPs are described as cytoplasmic influx antiporters, which use the energy of the proton gradient to transport a wide variety of cations (zinc, iron, manganese, copper and cadmium) (Guerinot, 2000). The knowledge on metal specificity, subcellular localization and overall function of individual ZIP transporters in plant metal homeostasis remains very fragmentary (Ricachenevsky et al., 2015).

In contrast to most of its family members, *ZIP6* expression is not induced upon zinc deficiency in *A. thaliana* (Talke et al., 2006) and *zip6* mutant plants do not display obvious phenotypes (Wu et al., 2009). Overexpression of the *N. caerulea* *ZIP6* (*ZNT6*) in *A. thaliana* resulted in a weak increase of root cadmium sensitivity in one of three independent transgenic lines (Wu et al., 2009).

Here, we examined the contribution of the *A. halleri* *ZIP6* (*AhZIP6*) gene to zinc and cadmium tolerance and hyperaccumulation. We showed a tandem duplication of *ZIP6* in the genome of different *A. halleri* ecotypes and a partial tissue specialization of the two copies. We determined that *AhZIP6* is a zinc and cadmium transporter, mainly expressed in vascular tissues. Using *ZIP6* RNAi *A. halleri* lines and *A. thaliana* transgenic plants overexpressing *AhZIP6*, we demonstrated that *AhZIP6* alters cadmium tolerance.

2 | RESULTS

2.1 | ZIP6 is duplicated in tandem in the *A. halleri* genome

Several highly expressed metal homeostasis genes are present in multiple copies in the *A. halleri* genome (e.g. *MTP1* or *HMA4*) (Dräger et al., 2004; Hanikenne et al., 2008; Shahzad et al., 2010; Suryawanshi et al., 2016). To determine *AhZIP6* gene copy number at a species-wide level, six ecotypes were selected from the two genetic units and the hybrid zone identified within the geographic distribution of

A. halleri in Europe to include both metallicolous (I16, Lan3.1, Lan5, PI22) and non-metallicolous (I30, Sk13) populations (Pauwels et al., 2012). As expected, a single *ZIP6* copy was detected in the *A. thaliana* and *A. lyrata* genomes (Figure 1a), whereas *ZIP6* was duplicated in all *A. halleri* ecotypes, except in I16 where it was triplicated (Figure 1a).

To further examine the genomic architecture of the *ZIP6* loci in the *A. halleri* genome, bacterial artificial chromosome (BAC) clones (Hanikenne et al., 2013) harbouring *AhZIP6* were isolated and analysed by quantitative PCR, which confirmed the gene duplication and indicated that both *AhZIP6* copies were on the same genome




FIGURE 1 Analysis of the *ZIP6* locus in different *Arabidopsis* species. (a) Determination of *ZIP6* gene copy numbers in *A. thaliana* (Col-0), *A. lyrata* (Lyr) and different populations of *A. halleri* (I16, I30, Lan3.1, Lan5, PI22 and Sk13) originating from the South-East (SE), North-West (NW) and Hybrid Zone (HZ) genetic units defined by Pauwels et al. (2012). Quantitative PCR was performed in triplicate on genomic DNA of pools of 10 individuals per population using two sets of primers (I and II). *A. thaliana* (Col-0) and two single copy genes (*FRD3* and *S13*) were used to normalise gene copy numbers (means \pm SEM; from one experiment representative of two independent experiments). (b) Schematic representation of the *ZIP6* promoters. Based on multiple alignments, coloured boxes represent conserved DNA sequences with more than 75% of identity and a minimal length of 30 bp. Red, orange and yellow indicate the presence of conserved boxes in four, three and two promoters, respectively. Letters and numbers show boxes correspondence and position. Arrowheads represent the orientation of the boxes. All *ZIP6* coding sequences are represented by black boxes and are not at the same scale. *ZIP6* promoter sequences were collected from Phytozome (<https://phytozome.jgi.doe.gov/>) for *A. thaliana* and *A. lyrata* and were cloned from the Lan3.1 population for *A. halleri* in this study [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]

segment (Figure S1). PCR analyses determined a tandem and forward organization of the two copies separated by ~5 kb in the Lan3.1 genome (Figure 1b).

The promoter of the *AhZIP6* copy 1 (pAhZIP6-1, 3,038 bases upstream the ATG codon) of Lan3.1 was cloned by PCR thanks to sequence conservation with the *A. thaliana* (pAtZIP6) and *A. lyrata* (pAlZIP6) genomic sequences (Figure 1b). No sequence information was available for the *AhZIP6* copy 2 promoter. BAC clone Illumina sequencing and draft assemblies identified putative sequences that were then cloned by PCR from Lan3.1 genomic DNA (pAhZIP6-2, 3,116 bases upstream the ATG codon). Sequence analysis revealed that pAhZIP6-1 was closely

related to pAlZIP6 (seven blocks covering >2 kb and sharing an average of 89.3% of identity, excluding gaps) (Figure 1b and Figure S2, blocks a–g) and to a lesser extent to pAtZIP6 (five blocks covering ~1.1 kb with 80.4% average identity) (Figure 1b and Figure S2, blocks a, b, d, g and j). In contrast, the sequence of pAhZIP6-2 was highly divergent compared to the three others, with only two small conserved segments of 67 bp located in the 5'UTR and 34 bp including a putative TATA box, respectively (Figure 1b and Figure S2, red blocks). Using differences in promoter sequences and transcriptome data (from the PI22 ecotype) (Schvartzman et al., 2018), the coding sequences (CDS) of the 2 *AhZIP6* copies were cloned from Lan3.1 genomic DNA and were highly similar with 99.4% of




FIGURE 2 Localization of promoter activity and transcript level of *AtZIP6* or two *AhZIP6* copies in *A. thaliana* and *A. halleri*. (a) Histochemical detection of GUS activity (blue) directed by the pAtZIP6, pAhZIP6-1 or pAhZIP6-2 promoters in roots and leaves in *A. thaliana* (Top) and *A. halleri* (Bottom). Pictures are representative of six independent lines per construct. (b) Relative transcript levels (RTL) of *AhZIP6-1*, *AhZIP6-2* and both copies (*AhZIP6-Tot*) in wild-type (Lan3.1) *A. halleri* plants grown 7 weeks (long days) in Hoagland hydroponic medium in control condition (5 μ M Zn). Values (mean \pm SEM; from three biological replicates of two plants each) are relative to *EF1 α* and *UBQ13*. (c) ZIP6 RTL in transgenic homozygous *A. thaliana* plants expressing pAhZIP6-1::AhZIP6 (*AhZIP6-1*), pAhZIP6-2::AhZIP6 (*AhZIP6-2*) or in Col-0 (*AtZIP6*) grown 6 weeks (short days) in Hoagland hydroponic medium control condition (1 μ M Zn). Values (means \pm SEM; from two biological replicates of two plants each for three independent lines per construct) are relative to *EF1 α* and *At1g58050*. Data were analysed by two-way ANOVA followed by Bonferroni multiple comparison post-test. Statistically significant differences between means are indicated by asterisks (** p < .01, *** p < .001) or different letters (p < .05) [Colour figure can be viewed at wileyonlinelibrary.com]

amino acid sequence identity and closely related to *A. thaliana* (~97%) and *A. lyrata* (~97%) ZIP6 sequences (Figure S3). The two sequences only diverged in a poly-glycine stretch (Figure S3C), outside conserved motifs found in transmembrane segments IV and V in ZIP proteins (Eng, Gueriot, Eide, & Saier, 1998; Gueriot, 2000), suggesting that polymorphisms among the two AhZIP6 proteins sequences are unlikely to result in functional differences. The version with the shortest Gly-stretch (8 Gly) was used in all downstream functional analysis.

2.2 | The AhZIP6 copies display partial tissue specialization

Next, the expression profile and expression level of the AhZIP6 copies were determined. Firstly, the pAhZIP6-1 and pAhZIP6-2 promoters, as well as pAtZIP6 for comparison, were fused to the *GUS* reporter gene for stable transformation in *A. thaliana* and *A. halleri*. A highly similar expression profile was observed for all three promoters in vegetative tissues of both species: ZIP6 was mainly expressed in the root and leaf vascular tissues, but was also expressed in the root cap, in leaf mesophyll cells and in hydathodes (Figure 2a and Figure S4). This profile was consistent with the AtZIP6 expression data available at Genevestigator (<https://genevestigator.com/gv/>).

Secondly, copy-specific primers were designed and the expression level of each AhZIP6 copy was determined in wild-type *A. halleri* plants (ecotype Lan3.1). In roots, AhZIP6-2 was threefold more expressed than AhZIP6-1. In contrast, AhZIP6-1 was 1.7-fold more expressed than AhZIP6-2 in cauline leaves and both copies showed similar expression level in shoots (Figure 2b). As control, the AhZIP6 expression level detected with primers targeting simultaneously both copies was roughly similar to the addition of the AhZIP6-1 and AhZIP6-2 specific expression.

Finally, *A. thaliana* T3 homozygous transgenic lines expressing AhZIP6 under the control of either pAhZIP6-1 (AhZIP6-1 lines) or pAhZIP6-2 (AhZIP6-2 lines) were obtained. Both AhZIP6 copies were more highly expressed than AtZIP6 in roots and shoots of *A. thaliana* (Figure 2c). In roots, AhZIP6-2 was again predominant and was, respectively, 2- and 10-fold more expressed than AhZIP6-1 and AtZIP6. In contrast, in shoots and cauline leaves, AhZIP6-1 was dominant (1.5-fold higher than AhZIP6-2 and 20-33-fold higher than AtZIP6, respectively) (Figure 2c).

Altogether, these analyses indicated a higher expression of both AhZIP6 copies compared with AtZIP6 and suggested overall conserved *cis* regulatory mechanisms between *A. halleri* and *A. thaliana*, with a partial organ specialization of each copy: AhZIP6-1 was predominant in aerial parts and AhZIP6-2 in roots in both species.

2.3 | The localization of the AhZIP6 protein remains elusive

GUS staining and qRT-PCR provide relevant information on expression profile and expression level. However, subcellular localization is

important to elucidate the protein function. Green fluorescent protein (GFP) was thus fused to the ZIP6 genomic DNA sequence (AhZIP6_g). Genomic DNA was used instead of cDNA to resolve a problem of apparent bacterial toxicity during cloning. *A. thaliana* plants were stably transformed with two constructs (pAhZIP6-1::AhZIP6_g::GFP and pAhZIP6-2::AhZIP6_g::GFP) and five independent T2 lines per construct were observed by confocal microscopy. Intense fluorescence was only distinguished for pAhZIP6-1::AhZIP6_g::GFP lines. In the epidermis of mature roots, fluorescent spots were observed on the cell edges and in some points co-localized with propidium iodide, a fluorescent dye staining the cell wall (Figure 3). However, observation in developing roots (root tip and elongation zone) where vacuoles are smaller and the cytoplasm wider, detected a dotted cytoplasmic fluorescence that did not co-localize with plasma membrane (Figure 3). AhZIP6_g::GFP was probably expressed in small cytoplasmic vesicles, which were squeezed against the plasma membrane when vacuoles were fully developed. Additional efforts using *A. thaliana* protoplasts transformed with various tagged versions of AhZIP6-1 under control of the 35S promoter yielded equally ambiguous results. Protoplasts showed both cytosolic signals and in some cases, a staining pattern that could be interpreted as plasma membrane (Figure S5A–C). Note, however, that the functionality of the GFP fusion has to be questioned based on the expression in yeast (see next section).

2.4 | AhZIP6 is a zinc and cadmium transporter in yeast

As *A. halleri* is a zinc and cadmium hyperaccumulator plant, we next tested the transport ability of the AhZIP6 protein upon constitutive expression in zinc- (*zrc1cot1*) and cadmium- (*ycf1*) sensitive yeast mutants, defective for vacuolar metal transport, and the corresponding wild-type strains. Metal tolerance of the transformed yeast cells was assessed upon growth both on agar plates and in liquid medium contaminated with a range of zinc or cadmium concentrations, respectively (Figure 4). The expression of AhZIP6 had no effect on growth of either strain in control conditions (Figure 4). In contrast, AhZIP6 increased the zinc sensitivity of the *zrc1cot1* strain, both in drop tests and in liquid cultures, with a complete inhibition of growth at 200 µM zinc in liquid culture for instance (Figure 4a,b). However, zinc concentration did not differ between AhZIP6-expressing and empty vector-transformed yeast cells, respectively (Figure S6A). Similarly, AhZIP6 expression increased cadmium toxicity in the *ycf1* mutant (Figure 4c,d). Additionally, the growth of the wild-type strain expressing AhZIP6 was also affected when exposed to the highest cadmium concentration (25 µM) in liquid cultures (Figure 4d). Again, no major difference in cadmium concentration in cells was found between the different yeast genotypes (Figure S6). As AhZIP6 seemed to localize to endomembranes in yeast cells when fused to GFP (Figure S5D), these data suggested that AhZIP6 was able to transport zinc and cadmium and was possibly involved in compartmentalization rather than cellular zinc and cadmium uptake. Note, however, that expression of the AhZIP6::GFP fusion in yeast did not cause the zinc




FIGURE 3 AhZIP6 localization in *A. thaliana* roots. Plants expressing pAhZIP6-1::AhZIP6::GFP were grown vertically in Petri plates (control condition 1 μ M Zn) for 6 days. Green fluorescent protein (GFP) was detected as a dotted cytoplasmic pattern (green). Propidium iodide (PI) was used to stain the cell wall (red). Pictures were taken with a Leica SP2 confocal microscope. GFP and PI were excited at 488 nm and detected between 500 and 550 or 585 and 650 nm, respectively. Scale bars: 20 μ m [Colour figure can be viewed at wileyonlinelibrary.com]

and cadmium sensitivity phenotype observed with the native protein, questioning its functionality.

2.5 | Silencing ZIP6 expression in *A. halleri* does not affect zinc and cadmium accumulation

As expected from previous observations (Talke et al., 2006), AhZIP6 expression levels were only marginally regulated in *A. halleri* upon zinc deficiency (0 μ M Zn), high zinc (300 μ M) or cadmium (5 μ M) exposure (Figure 5). Noticeably, the two AhZIP6 copies displayed a similar behaviour, with the exception of a repression of AhZIP6-2 at high zinc, amounting to a global down-regulation of AhZIP6 in shoots (Figure 5b).

To more directly address the physiological role of AhZIP6 in metal hyperaccumulation and hypertolerance, a vegetatively propagated

individual from the Langelsheim population was transformed with an RNAi construct targeting the two AhZIP6 copies. Thirty-one independent transgenic lines were obtained and screened for AhZIP6 total transcript abundance in shoots using primers matching both AhZIP6 copies. Two lines with nearly 90% suppression of AhZIP6 transcript (lines 7 and 11) were selected for further analysis (Figure 6a). Analysis of the RNAi effect on the two AhZIP6 copies separately using specific primers confirmed a suppression of about 90% for both copies in the selected transgenic lines compared with the wild type (Figure S7). The RNAi effect was also confirmed in comparison to a transgenic line expressing GFP under control of the 35S promoter as an additional control besides wild-type plants (Figure 6 and Figure S7) since it had undergone the transformation procedure including prolonged phases of tissue culture.

Wild-type and transgenic lines were grown in hydroponic culture under control conditions (10 μ M Zn). After 3 weeks, micronutrient




FIGURE 4 Functional analysis of the AhZIP6 cDNA in zinc- and cadmium-sensitive yeast strains. Expression of AhZIP6 or the empty vector (ev, pFL38) in the zinc-sensitive *zrc1cot1* double mutant (a, b) or in the cadmium-sensitive *ycf1* mutant and their respective parental strains (BY4741 and BY4742) (c, d). (a, c) Drop tests. Transformants were serially diluted at OD₆₀₀ of 0.2, 0.02, 0.002 (left to right) and spotted on LSP-Leu medium with different zinc or cadmium concentrations. Plates were incubated at room temperature (~20°C) for 4 days. The pictures are representative of two independent experiments, each including four independent transformants. (b, d) Growth curves. Transformants were inoculated at an initial OD₆₀₀ of 0.003 in liquid LSP-Leu medium with different zinc or cadmium concentrations, then grown for 4 days (250 rpm) at room temperature (~20°C) before OD₆₀₀ was measured. Values (means ± SEM; from four independent cultures) were normalized to the parental wild-type strain expressing the empty vector in control condition (2 μM Zn or no added cadmium). Data were analysed by one-way ANOVA followed by Tukey multiple comparison post-test. Statistically significant differences between means are indicated by letters ($p < .05$)

contents were determined for roots and leaves. Typical strong accumulation of zinc in leaves was apparent but no differences between lines were observed for zinc, iron and manganese (Figure S8). Similarly, no genotype-dependent differences were found when plants were cultivated in excess zinc (300 μM Zn) (Figure S8).

Next, the two AhZIP6-suppressed lines and the two controls were grown in native *A. halleri* soil from the metalliferous site in Bestwig (for GPS and soil data, see Höreth et al., 2020) and in the phenotyping soil used for the characterization of field-collected individuals of European *A. halleri* populations (Stein et al., 2017). Growth as determined by shoot biomass was comparable between the metalliferous site soil and the phenotyping soil, indicating that the metal-contaminated soil did not exert toxic effects on the plants. No significant differences between genotypes were apparent (Figure S9). After cultivation in Bestwig soil, zinc accumulation in all lines exceeded by far the zinc hyperaccumulation threshold of 3,000 μg g⁻¹ of dry weight, whereas in phenotyping soil with very limited zinc phytoavailability, accumulation stayed below this threshold (Figure S10). In contrast, cadmium accumulation was higher in phenotyping soil exceeding the hyperaccumulation threshold of 100 μg g⁻¹ of dry weight, but not in Bestwig soil (Figure S10). Again,

however, no significant differences were observed between the lines, arguing against a major direct impact of AhZIP6 on zinc or cadmium hyperaccumulation.

2.6 | Silencing ZIP6 expression in *A. halleri* specifically increases cadmium tolerance

Shoot growth in Bestwig soil had not been inhibited by high metal content. To assess possible metal tolerance effects of AhZIP6 silencing directly on roots and in a more targeted, element-specific manner, plants were exposed to growth-inhibiting concentrations of different metals in hydroponic culture. Conditions tested were 600 μM zinc, 5 and 10 μM nickel and 10 and 20 μM cadmium. These concentrations were chosen in order to cause a comparable range of root growth inhibition effects on wild-type plants. Zinc concentrations could not be increased any further without precipitation of zinc salts. Importantly, AhZIP6-suppressed plants grew significantly better than control plants specifically in the presence of cadmium (Figure 6b–d). The gain in tolerance was restricted to root length and root fresh weight, and seemed specific to cadmium as all genotypes




FIGURE 5 Relative transcript levels of the two *AhZIP6* copies in *A. halleri* upon different metal exposures. Relative transcript levels (RTL) of *AhZIP6-1*, *AhZIP6-2* and both copies (*AhZIP6-Tot*) in roots (a) and shoots (b) of in wild-type (Lan3.1) *A. halleri* plants grown (in long days) 4 weeks in Hoagland hydroponic medium (5 μM Zn) then exposed for 3 weeks to the following conditions: zinc deficiency (0 μM Zn), control condition (5 μM Zn), high zinc (300 μM Zn) and exposure to cadmium (5 μM Cd). Values (means ± SEM; from three biological replicates of two plants each) are relative to *EF1α* and *UBQ13*. Data were analysed by two-way ANOVA followed by Bonferroni multiple comparison post-test. Statistically significant differences between means are indicated by asterisks (* $p < .05$, ** $p < .01$, *** $p < .001$) or different letters ($p < .05$)

were equally sensitive to zinc and nickel (Figure 6b–d, Figures S11 and S12).

The higher cadmium tolerance of *AhZIP6*-suppressed lines could be due to a reduction in root cadmium uptake activity. Thus, we measured short-term cadmium uptake by roots of hydroponically grown plants. Experiments performed with non-toxic concentrations of 2 and 5 μM cadmium showed indeed that uptake by roots of *AhZIP6*-suppressed plants was strongly reduced compared with control plants (Figure 7). Thus, *AhZIP6* appears to represent a cadmium uptake pathway in *A. halleri*.

When cadmium accumulation after long-term exposure in hydroponic culture was assayed, however, no differences between control and *AhZIP6*-RNAi plants were observed (Figure S13). This confirmed the findings of the soil experiments.

2.7 | In *A. thaliana*, *AhZIP6* impacts the growth under cadmium exposure

To define the function of each *AhZIP6* copy, wild-type (Col-0) and T3 homozygous *A. thaliana* plants expressing *AhZIP6* under the control of either p*AhZIP6-1* (*AhZIP6-1* lines) or p*AhZIP6-2* (*AhZIP6-2* lines) were grown for 3 weeks in control hydroponic medium (1 μM Zn), then exposed for 3 weeks to the following conditions: control (1 μM Zn), zinc deficiency (0 μM Zn), zinc excess (20 μM) or presence of cadmium (0.05 μM). Although the expression of both *AhZIP6* copies was not altered by the zinc status or by cadmium concentration in the medium (Figure 8a,b), the fresh shoot biomass of all genotypes was impacted by the treatments (Figure 8c). Zinc deficiency and zinc excess decreased the shoot biomass with, for instance, 22 and 35% biomass reduction for Col-0, respectively, but no differences were observed between genotypes (Figure 8c). *AhZIP6* expression in *A. thaliana* had overall a very limited impact on the plant ionome, across all culture conditions (Figure S14). Indeed, only a small but significant reduction (8%) of zinc concentration was found in shoots of *AhZIP6* lines upon zinc excess (Figure S14A).

In contrast to zinc, cadmium exposure differentially impacted shoot biomass in the different genotypes: shoots of *AhZIP6-2* plants had indeed 23% higher biomass than Col-0, which in turn had 30% higher biomass than *AhZIP6-1* shoots (Figure 8c). Thus, high expression of *AhZIP6* in roots triggered by the *AhZIP6-2* construct contributed to increased shoot cadmium tolerance, whereas high expression of *AhZIP6* in shoots triggered by the *AhZIP6-1* construct increased shoot cadmium sensitivity. This occurred despite similar cadmium accumulation in roots and shoots among genotypes (Figure S14C). As for *A. halleri* RNAi lines, short-term cadmium uptake by roots of hydroponically grown plants was measured in Col-0 and in *AhZIP6-2* lines. *AhZIP6-2* lines were used as this copy is the most highly expressed in roots (Figure 2b,c). Cadmium uptake by roots was found to be similar between the two genotypes (Figure S15).

3 | DISCUSSION

In spite of their long suspected importance for zinc acquisition and other aspects of zinc homeostasis, very little is known about the physiological roles of ZIP transporters (Ricachenevsky et al., 2015). Plant ZIP proteins were detected in different cellular membranes and various, sometimes contradictory, phenotypes were reported for yeast cells expressing ZIPs. With the exception of the IRT proteins, hardly any phenotype could to date be associated with a loss or gain of ZIP function in plants.

We analysed *AhZIP6* because ZIP6 genes from *A. halleri* and *N. caerulea* show all the known hallmarks of a metal hyperaccumulation candidate gene. As many other metal homeostasis genes, they are highly expressed in *A. halleri* and *N. caerulea* compared with non-accumulator relatives and, more importantly, *AhZIP6* was found to be associated with a QTL for zinc accumulation in *A. halleri* (Becher et al., 2004; Filatov et al., 2006; Hammond




FIGURE 6 Identification of *AhZIP6*-RNAi *A. halleri* lines and cadmium tolerance test. (a) *AhZIP6* transcript levels were analysed in shoots of wild-type, a control GFP-transformed line and two *AhZIP6*-suppressed (7-2 and 11-1) *A. halleri* lines grown on normal soil. Tissues were harvested after 6 weeks of cultivation and analysed by quantitative RT-PCR. Transcript abundance was expressed relative to *EF1α*. Values are means \pm SD from two independent experiments including a pool of six replicate clones per genotype for each data point. Cadmium tolerance was determined by measuring root length (b), and root weight (c) relative to control conditions of plants grown hydroponically for 3 weeks in 1/10 Hoagland medium supplemented with 0 (control), 10 or 20 μ M cadmium. Pictures of the representative plants at the end of the experiment are shown (d). Values (means \pm SD, $N = 12$ replicates from three independent experiments) are relative to the control condition and were analysed by ANOVA followed by Bonferroni multiple comparison post-test. Statistically significant differences between means are indicated by asterisks (* $p < .05$, ** $p < .01$, *** $p < .001$) or different letters ($p < .05$) [Colour figure can be viewed at wileyonlinelibrary.com]

et al., 2006; Talke et al., 2006; Weber, Harada, Vess, Roepenack-Lahaye, & Clemens, 2004; Willems et al., 2010).

High expression of these metal homeostasis genes was often found to be associated with gene copy number amplification (Dräger et al., 2004; Fasani et al., 2017; Hanikenne et al., 2008; Shahzad et al., 2010; Suryawanshi et al., 2016). For instance, both *HMA4*, critical for zinc root-to-shoot translocation (Courbot et al., 2007; Hanikenne et al., 2008; Hanikenne et al., 2013; Nouet et al., 2015), and *MTP1*, involved in zinc vacuolar storage in shoots (Dräger et al., 2004; Fasani et al., 2017; Krämer, 2005; Shahzad et al., 2010; Willems et al., 2007), are present in multiple copies in the *A. halleri* genome.

Here, we showed that *ZIP6* is a multicopy gene (Figure 1a) in both metallicolous and non-metallicolous *A. halleri* populations representing the geographic distribution of the species in Europe (Pauwels et al., 2012). For most genotypes, our analysis revealed that *ZIP6* is duplicated (Figure 1a). This suggests that the duplication occurred before the last glacial age (around 100 000–15 000 years ago), which isolated the *A. halleri* genetic units (Pauwels et al., 2012). However, the number of *ZIP6* copies in the *A. halleri* genome is not fixed. Three *ZIP6* copies were indeed detected in the I16 Italian metallicolous population (Figure 1a) that colonizes a highly zinc-, cadmium- and lead-contaminated environment and displays a cadmium exclusion strategy




FIGURE 7 Short-term cadmium uptake in wild-type and *AhZIP6*-RNAi *A. halleri* lines. Cadmium concentration in roots of wild-type, control GFP-transformed line and two independent *AhZIP6*-suppressed lines (7-2 and 11-1) grown hydroponically for 2 weeks in 1/10 Hoagland medium and then exposed for a short term (15 or 30 min) to media containing 2 µM (a) and 5 µM (b) cadmium at 4°C or 25°C. Values (means ± SD) are from three independent experiments, each including 4–5 replicate clones per genotype. Data were analysed by two-way ANOVA followed by Bonferroni multiple comparison post-test. Statistically significant differences between means are indicated by different letters ($p < .05$)

(Corso et al., 2018; Fasani et al., 2017; Meyer et al., 2015; Schwartzman et al., 2018). Unfixed gene copy number among populations was also found for *MTP1* (Fasani et al., 2017; Meyer et al., 2016; Shahzad et al., 2010). Gene copy number variation among and within populations may reflect ongoing local adaptation processes. For instance, the third copy might differentially contribute to cadmium tolerance or accumulation in the I16 population (Corso et al., 2018).

As for *HMA4* (Hanikenne et al., 2008), the two *ZIP6* copies are found in tandem in the *A. halleri* genome (Figure 1b), at an even closer distance, and remain highly conserved (Figure S3). This suggests, as for *HMA4* (Hanikenne et al., 2013), that the *ZIP6* copies may undergo ectopic gene conversion enabling concerted evolution of the paralogous *ZIP6* coding sequences. In contrast, the promoter sequences of the *ZIP6* copies have considerably diverged, again reminiscent of *HMA4* and *MTP1* (Fasani et al., 2017; Hanikenne et al., 2008; Shahzad et al., 2010), with p*AhZIP6*-1 retaining the ancestral sequence most similar to the *A. lyrata* and *A. thaliana* *ZIP6* promoters (Figure 1b). However, unlike *MTP1* (Shahzad et al., 2010), both *ZIP6* copies are highly expressed in *A. halleri*, compared with *A. thaliana* (Figure 2). In contrast to *HMA4* (Hanikenne et al., 2008), a (partial) sub-functionalization of the two copies occurred after the duplication. While the two copies retained a highly similar tissue-specificity of expression, at least in vegetative tissues (Figure 2a), *AhZIP6*-1 and *AhZIP6*-2 are differentially expressed in root and shoot tissues (Figure 2b,c).

Taken together, the genomic situation and expression analyses further substantiated *AhZIP6* as an important candidate gene possibly contributing to zinc and/or cadmium hyperaccumulation and hyper-tolerance in *A. halleri*. We therefore transformed an *A. halleri* individual from the metalcolous Langelshiem population with an RNAi construct and obtained lines with strong suppression of both *AhZIP6* transcripts (Figure 6a, Figure S7). However, no significant changes in

root or leaf ionomes were detected in these lines after hydroponic or soil cultivation in a wide range of conditions (Figures S7–S10), arguing against a direct role of *AhZIP6* in metal hyperaccumulation and against *AhZIP6* being the causal gene detected in hyperaccumulation QTLs (Filatov et al., 2006; Willems et al., 2010).

Surprisingly, tolerance assays with *AhZIP6*-suppressed plants and respective controls revealed a gain in cadmium tolerance when the two *AhZIP6* copies are knocked down (Figure 6). This effect appeared to be specific for cadmium, over zinc or nickel (Figure S12). A gain in cadmium tolerance upon the suppression of a gene could be indicative of cadmium uptake activity. ZIP transporter-mediated cadmium transport has been described for ZIP proteins from other kingdoms before. For example, loss-of-function mutations in mouse ZIP8 (SLC39A8) explain the resistance of particular inbred strains to cadmium-induced testicular necrosis (Dalton et al., 2005) because ZIP8 represents the main cadmium uptake pathway. Mouse ZIP14 was shown to transport cadmium into mouse kidney cells (Fujishiro, Yano, Takada, Tanihara, & Himeno, 2012).

Short-term cadmium uptake assays indeed demonstrated a significant reduction in roots of *AhZIP6*-suppressed plants (Figure 6). Thus, *AhZIP6* can be added to the small list of plant transporters for which cadmium transport activity has been directly shown. These include IRT1 in *A. thaliana* (Vert et al., 2002) and OsNramp5 in rice (Ishimaru et al., 2012; Sasaki, Yamaji, Yokosho, & Ma, 2012). Indirect support for a cadmium, and zinc, transport activity of *AhZIP6* was also derived from yeast expression (Figure 4). Furthermore, *AhZIP6*-dependent cadmium uptake is consistent with the reported higher cadmium sensitivity of *A. thaliana* plants overexpressing the ZIP6 ortholog from *N. caerulescens* (Wu et al., 2009).

Unlike OsNramp5 loss-of-function mutants, however, which accumulate much less cadmium in above-ground tissues than respective wild-type plants (Ishimaru et al., 2012; Sasaki et al., 2012), *AhZIP6*-knock-down plants did not show any alterations in cadmium




FIGURE 8 Analysis of AhZIP6 expression in *A. thaliana* upon different metal exposure. Transgenic homozygous *A. thaliana* plants expressing pAhZIP6-1::AhZIP6 (AhZIP6-1), pAhZIP6-2::AhZIP6 (AhZIP6-2) or in Col-0 plants (AtZIP6) were grown (in short days) for 3 weeks in Hoagland hydroponic medium (1 μ M Zn) then exposed for 3 weeks to the following conditions: zinc deficiency (0 μ M Zn), control condition (1 μ M Zn), zinc excess (20 μ M Zn) and exposure to cadmium (0.05 μ M Cd). ZIP6 relative transcript levels (RTL) were quantified in roots (a) or shoots (b). Values (means \pm SEM; from two biological replicates of two plants each for three independent lines per construct) are relative to *EF1 α* and *At1g58050*. (c) Measurement of shoot fresh weight. Values (means \pm SEM; from one experiment representative of two independent experiments, with 14–24 plants for each three independent line per construct). Data were analysed by two-way ANOVA followed by Bonferroni multiple comparison post-test. Statistically significant differences between means are indicated by asterisks (* p < .05, *** p < .001) or different letters (p < .05)

accumulation behaviour (Figures S10 and S13). We therefore hypothesize that (a) several pathways for cadmium uptake into *A. halleri* roots exist that mask the effects of AhZIP6 knock-down when integrating uptake over a long period of time in accumulation experiments, and that (b) AhZIP6 may mediate transport of cadmium into particularly sensitive root cells, thus explaining the tolerance phenotype.

Cadmium hypertolerance is a species-wide trait in *A. halleri* (Meyer et al., 2015; Stein et al., 2017). Thus, it is difficult to explain why AhZIP6 should be more highly expressed in *A. halleri* and possibly still subject to local adaptation processes (see above) when the activity of AhZIP6 renders plants more cadmium sensitive. Arguably, the simplest explanation would be that AhZIP6 is involved in other processes important for metal hyperaccumulation, for example, by fine-tuning metal distribution within tissues. Cadmium transport by AhZIP6 would in such a scenario be a tolerable side effect, consistent with the lack of any biomass reduction of AhZIP6 RNAi plants after cultivation in native metalliferous *A. halleri* soil (Figure S9).

Most plant ZIP transporters have so far been indirectly implicated in zinc transport, making zinc the most likely candidate metal to guide the search for other suspected ZIP6 functions. As was the case for practically all studies on plant ZIP transporters to date, however, our *A. halleri* experiments did not uncover zinc ionome phenotypes upon suppression of ZIP gene function. Most available evidence for zinc transport by plant ZIPs has been obtained in overexpression studies, for example, OsZIP7 (Ricachenevsky et al., 2018) and HvZIP7 (Tiong et al., 2014). We therefore generated *A. thaliana* lines expressing the AhZIP6 gene copies under their native promoters (Figure 8), allowing copy-specific analyses not possible in *A. halleri*.

Overall, subtle metal-related phenotypes were observed for the transgenic lines. AhZIP6 expression affected cadmium tolerance, albeit in opposite directions for the two copies. AhZIP6-2 is responsible for higher ZIP6 expression in roots and conferred increased cadmium tolerance to the entire plant without modifying the ionomics profile in tissues (Figures 2b and 8, Figure S14) or contributing measurably to root cadmium uptake (Figure S15). AhZIP6-1 expression, in contrast, rendered plants slightly more cadmium sensitive (Figure 8). Taken together, these observations provided further indirect support for a role of AhZIP6 in cadmium transport. It thus seems that high expression of AhZIP6 in an organ conferred cadmium sensitivity locally (e.g. in roots of *A. halleri*, Figure 6, or in shoots for AhZIP6-1 in *A. thaliana*, Figure 8c), but could also act on cadmium tolerance of another organ (i.e. AhZIP6-2 in *A. thaliana*, Figure 8c). This distinction could only be revealed by analysing the two copies separately in *A. thaliana*, whereas the phenotypic output in *A. halleri* resulted from the combined actions of the two copies.

At the end of this study, the cellular localization of AhZIP6 remains unclear. A protein mediating short-term cadmium uptake would be expected to reside in the plasma membrane (Figure 7). On the contrary, the absence of ionome phenotype in yeast (Figure S6), *A. thaliana* (Figures S14 and 15) or *A. halleri* (Figures S10 and S13) over or underexpressing AhZIP6 may equally rule out plasma membrane localization and rather support a role for AhZIP6 in intracellular cadmium distribution. In this case, altered cadmium distribution in

cells may indirectly trigger the increased cadmium uptake observed at short term in *A. halleri*. Neither of these hypotheses could be confirmed or falsified by our localization experiments. AhZIP6 fused to a fluorescent reporter protein in C-terminal seems to localize in vesicular endomembranes in yeast cells (Figure S5D) and in Arabidopsis plants or protoplasts (Figure 3, Figure S5A–C). The localization of AhZIP6 is somewhat reminiscent of those of IRT1 (Dubeaux, Neveu, Zelazny, & Vert, 2018) and NRAMP1 (Agorio et al., 2017), which were described to undergo intracellular trafficking and whose localization on endomembranes or the plasma membrane is regulated by iron and other metal availability (Agorio et al., 2017; Dubeaux et al., 2018). Whether the localization of AhZIP6 is submitted to such dynamic processes depending on cadmium and/or zinc availability will need to be examined in the future. The observations reported here have, however, to be interpreted with caution as the AhZIP6 fusion protein appeared no longer functional in yeast. These ambiguous observations may result from a possible toxicity of the protein—molecular cloning of tagged or untagged versions of AhZIP6 has been strongly hampered by toxicity in *E. coli* in the course of this study.

A working hypothesis is that AhZIP6 is key to fine-tune metal distribution at local scale, controlling metal homeostasis in specific cell types. Considering two scenarios for the protein localization, high expression of AhZIP6 would act on metal distribution between tissues (plasma-membrane localization) or between compartments within cells (endomembrane localization), producing the tolerance phenotype reported in this study. Moreover, AhZIP6 contributes to root cadmium uptake, although it is mostly expressed in vascular tissues (Figure 2). It has been shown for other transporters that expression in the vascular system can contribute to metal uptake. For instance, high expression of HMA4 in *A. halleri* root vascular tissues activates zinc uptake by ZIPs (Hanikenne et al., 2008). Another example is NRAMP1, which although mostly expressed in the endodermis, contributes together with IRT1 to iron uptake in *A. thaliana* roots (Cailliatte, Schikora, Briat, Mari, & Curie, 2010; Castaings, Caquot, Loubet, & Curie, 2016).

In conclusion, ZIP6 is a duplicated gene in *A. halleri*, expressed mainly in the vasculature and encoding a zinc and cadmium transporter. Each copy acquired some extent of organ specialization, in aerial parts (AhZIP6-1) and in roots (AhZIP6-2). The expression pattern of the AhZIP6 copies, and the activity of their promoters, was overall conserved in both *A. halleri* and *A. thaliana*, suggesting a conservation of *cis* regulatory mechanisms, as for HMA4 (Hanikenne et al., 2008). Overexpression of the AhZIP6 copies in *A. thaliana* or their extinction in *A. halleri* altered cadmium tolerance but had no impact on zinc tolerance or the ionome of the plants, suggesting a function of AhZIP6 in controlling tissular or cellular metal distribution. Altogether, this study highlights the diversity and complexity of mechanisms underlying the evolution of hyperaccumulation and the distinct fates of duplicated genes in *A. halleri*. Obvious hyperaccumulation candidate genes do not necessarily show a directly measurable loss-of-function impact on metal concentrations when the still tedious generation of respective knock-down lines is achieved. Our study also confirms that functional characterization of ZIPs in plants remains a challenge.

4 | METHODS

4.1 | Plant material, growth conditions and transformation

A. thaliana (accession Columbia-0, Col-0) and *A. halleri* ssp. *halleri* (accession Langelsheim) were used for all experiments. Growth conditions in hydroponic Hoagland medium and on soil are detailed in Text S1. *A. thaliana* transformation was performed by floral dip (Clough & Bent, 1998) and *Agrobacterium tumefaciens* (GV3101, pMP90) mediated stable transformation of *A. halleri* was performed using a tissue-culture based procedure (Ahmadi et al., 2018; Hanikenne et al., 2008).

4.2 | Determination of gene copy number and gene expression analysis

Gene copy numbers were determined by quantitative PCR as described (Hanikenne et al., 2013) using primers listed in Table S1. Gene expression analyses were performed by quantitative RT-PCR (Table S1). Data analysis and normalization were conducted as described (Talke et al., 2006; Nouet et al., 2015; Lekeux et al., 2018). Procedures are detailed in Text S1.

4.3 | Cloning, DNA manipulation and sequence analysis

The identification and characterization of the two AhZIP6 copies from *A. halleri* BACs (Hanikenne et al., 2008) using quantitative PCR, BAC sequencing and phylogenetic analysis (Kumar, Stecher, Li, Knyaz, & Tamura, 2018; Nei & Kumar, 2000) are detailed in Text S1. Vector constructs for expression of (un)tagged versions of AhZIP6 in yeast or plants and reporter constructs obtained using standard molecular cloning procedures and primers listed in Table S2 are described in Text S1.

4.4 | Yeast tolerance and complementation assays

Zinc and cadmium yeast assays were conducted using the *zrc1cot1* and *ycf1* mutants and their respective parental strains BY4741 and BY4742 (Euroscarf), the pFL38 vectors (Text S1) and LSP-Leu medium added with a range of zinc or cadmium concentrations (Talke et al., 2006).

4.5 | Protoplast isolation and transformation

Protoplasts were isolated from *A. thaliana* wild-type (Col-0) leaves and were transformed using a PEG solution as described in Text S1.

4.6 | Short-term uptake experiment

To examine the effect of *AhZIP6* suppression or over-expression on transport activity for cadmium in *A. halleri* or *A. thaliana*, respectively, a short-term (15 and 30 min) cadmium uptake experiment, as described in (Sasaki et al., 2012), was performed (Text S1).

4.7 | Analysis of metal contents

After desorption, plant and yeast samples were digested with HNO_3 (Sigma-Aldrich) as described (Scheepers et al., 2020) and analysed by inductively coupled plasma atomic emission spectroscopy (ICP-AES) with a Vista-AX instrument (Varian, Melbourne, Australia) or a iCAP 6,500 Series instrument (Thermo-Fisher, Waltham, MA) (yeast, roots and shoots) (Text S1).

4.8 | Confocal microscopy analysis

Confocal images were recorded using a Leica SP2 confocal microscope (Leica Microsystems, Wetzlar, Germany) or Leica TCS SP5 (Leica Microsystems) (Text S1).

4.9 | Histochemical staining

GUS staining was performed as described (Jefferson, Kavanagh, & Bevan, 1987) and tissues were then fixed and discoloured (Hanikenne et al. (2008).

4.10 | Accession numbers

The genomic sequences of *AhZIP6-1* and *pAhZIP6-2*, including the promoter sequences, were deposited in the NCBI Genbank database with the accession numbers MT155929 and MT155930, respectively.

ACKNOWLEDGMENTS

We thank Prof. N. Verbruggen, Prof. U. Krämer and S. Fanara for helpful discussions, as well as V. Deblander, V. Crutzen and A. Jadoul for technical assistance. *A. thaliana* HMA4-expressing lines were provided by Dr. C. Nouet. Funding was provided by the "Fonds de la Recherche Scientifique-FNRS" (PDR-T.0206.13, MIS-F.4511.16, CDR J.0009.17, PDR T0120.18 (M.H.), the University of Liège (SFRD-12/03) (M.H.) and the Belgian Program on Interuniversity Attraction Poles (IAP no. P7/44) (M.H.). S.C. gratefully acknowledges support from the Deutsche Forschungsgemeinschaft (SPP1529 ADAPTOMICS, CI152/9-1, -2). M.H. is Senior Research Associate of the F.R.S.-FNRS. J.S. and M.S. were doctoral fellows of the FNRS and the F.R.I.A., respectively. No conflict of interest declared.

AUTHOR CONTRIBUTIONS

Marc Hanikenne, Julien Spielmann and Stephan Clemens designed the research. Julien Spielmann, Hassan Ahmadi, Maxime Scheepers, Michael Weber, Sarah Nitsche, Monique Carnol, Bernard Bosman, Juergen Kroymann performed the experiments. Julien Spielmann, Hassan Ahmadi, Marc Hanikenne, Michael Weber, Stephan Clemens, Patrick Motte analysed the data. Julien Spielmann and Hassan Ahmadi made the Figures. Julien Spielmann, Marc Hanikenne, Stephan Clemens wrote the manuscript. All authors read and approved the manuscript.

ORCID

Julien Spielmann  <https://orcid.org/0000-0003-1327-3975>

Michael Weber  <https://orcid.org/0000-0001-5647-6217>

Stephan Clemens  <https://orcid.org/0000-0003-0570-1060>

Marc Hanikenne  <https://orcid.org/0000-0002-8964-9601>

REFERENCES

- Agorio, A., Giraudat, J., Bianchi, M. W., Marion, J., Espagne, C., Castaings, L., ... Merlot, S. (2017). Phosphatidylinositol 3-phosphate-binding protein AtPH1 controls the localization of the metal transporter NRAMP1 in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, 114, E3354–E3363.
- Ahmadi, H., Corso, M., Weber, M., Verbruggen, N., & Clemens, S. (2018). CAX1 suppresses Cd-induced generation of reactive oxygen species in *Arabidopsis halleri*. *Plant, Cell and Environment*, 41, 2435–2448.
- Baker, A. J. M. (1981). Accumulators and excluders-strategies in the response of plants to heavy metals. *Journal of Plant Nutrition*, 3, 643–654.
- Baliardini, C., Meyer, C. L., Salis, P., Saumitou-Laprade, P., & Verbruggen, N. (2015). CATION EXCHANGER1 cosegregates with cadmium tolerance in the metal hyperaccumulator *Arabidopsis halleri* and plays a role in limiting oxidative stress in *Arabidopsis* spp. *Plant Physiology*, 169, 549–559.
- Becher, M., Talke, I. N., Krall, L., & Krämer, U. (2004). Cross-species microarray transcript profiling reveals high constitutive expression of metal homeostasis genes in shoots of the zinc hyperaccumulator *Arabidopsis halleri*. *Plant Journal*, 37, 251–268.
- Bert, V., Bonnin, I., Saumitou-Laprade, P., De Laguérie, P., & Petit, D. (2002). Do *Arabidopsis halleri* from nonmetalliferous populations accumulate zinc and cadmium more effectively than those from metalliferous populations? *New Phytologist*, 155, 47–57.
- Cailliatte, R., Schikora, A., Briat, J. F., Mari, S., & Curie, C. (2010). High-affinity manganese uptake by the metal transporter NRAMP1 is essential for *Arabidopsis* growth in low manganese conditions. *The Plant Cell*, 22, 904–917.
- Castaings, L., Caquot, A., Loubet, S., & Curie, C. (2016). The high-affinity metal transporters NRAMP1 and IRT1 team up to take up iron under sufficient metal provision. *Scientific Reports*, 6, 37222.
- Charlier, J. B., Polese, C., Nouet, C., Carnol, M., Bosman, B., Krämer, U., ... Hanikenne, M. (2015). Zinc triggers a complex transcriptional and post-transcriptional regulation of the metal homeostasis gene *FRD3* in *Arabidopsis* relatives. *Journal of Experimental Botany*, 66, 3865–3878.
- Clemens, S. (2016). How metal hyperaccumulating plants can advance Zn biofortification. *Plant and Soil*, 411, 111–120.
- Clough, S. J., & Bent, A. F. (1998). Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant Journal*, 16, 735–743.
- Cornu, J. Y., Deinlein, U., Horeth, S., Braun, M., Schmidt, H., Weber, M., ... Clemens, S. (2014). Contrasting effects of nicotianamine synthase

- knockdown on zinc and nickel tolerance and accumulation in the zinc/cadmium hyperaccumulator *Arabidopsis halleri*. *New Phytologist*, 206, 738–750.
- Corso, M., Schwartzman, M. S., Guzzo, F., Souard, F., Malkowski, E., Hanikenne, M., & Verbruggen, N. (2018). Contrasting cadmium resistance strategies in two metalcolous populations of *Arabidopsis halleri*. *New Phytologist*, 218, 283–297.
- Courbot, M., Willems, G., Motte, P., Arvidsson, S., Roosens, N., Saumitou-Laprade, P., & Verbruggen, N. (2007). A major quantitative trait locus for cadmium tolerance in *Arabidopsis halleri* colocalizes with *HMA4*, a gene encoding a heavy metal ATPase. *Plant Physiology*, 144, 1052–1065.
- Dalton, T. P., He, L., Wang, B., Miller, M. L., Jin, L., Stringer, K. F., ... Nebert, D. W. (2005). Identification of mouse SLC39A8 as the transporter responsible for cadmium-induced toxicity in the testis. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 3401–3406.
- Deinlein, U., Weber, M., Schmidt, H., Rensch, S., Trampczynska, A., Hansen, T. H., ... Clemens, S. (2012). Elevated nicotianamine levels in *Arabidopsis halleri* roots play a key role in zinc hyperaccumulation. *The Plant Cell*, 24, 708–723.
- Dräger, D. B., Desbrosses-Fonrouge, A. G., Krach, C., Chardonens, A. N., Meyer, R. C., Saumitou-Laprade, P., & Krämer, U. (2004). Two genes encoding *Arabidopsis halleri* MTP1 metal transport proteins cosegregate with zinc tolerance and account for high MTP1 transcript levels. *Plant Journal*, 39, 425–439.
- Dubeaux, G., Neveu, J., Zelazny, E., & Vert, G. (2018). Metal sensing by the IRT1 transporter-receptor orchestrates its own degradation and plant metal nutrition. *Molecular Cell*, 69(953–964), e955.
- Eng, B. H., Gueriot, M. L., Eide, D., & Saier, M. H. (1998). Sequence analyses and phylogenetic characterization of the ZIP family of metal ion transport proteins. *Journal of Membrane Biology*, 166, 1–7.
- Fasani, E., DalCorso, G., Varotto, C., Li, M., Visioli, G., Mattarozzi, M., & Furini, A. (2017). The MTP1 promoters from *Arabidopsis halleri* reveal cis-regulating elements for the evolution of metal tolerance. *New Phytologist*, 214, 1614–1630.
- Filatov, V., Dowdle, J., Smirnov, N., Ford-Lloyd, B., Newbury, H. J., & Macnair, M. R. (2006). Comparison of gene expression in segregating families identifies genes and genomic regions involved in a novel adaptation, zinc hyperaccumulation. *Molecular Ecology*, 15, 3045–3059.
- Frérot, H., Faucon, M. P., Willems, G., Gode, C., Courseaux, A., Darracq, A., ... Saumitou-Laprade, P. (2010). Genetic architecture of zinc hyperaccumulation in *Arabidopsis halleri*: The essential role of QTL x environment interactions. *New Phytologist*, 187, 355–367.
- Fujishiro, H., Yano, Y., Takada, Y., Tanihara, M., & Himeno, S. (2012). Roles of ZIP8, ZIP14, and DMT1 in transport of cadmium and manganese in mouse kidney proximal tubule cells. *Metallomics*, 4, 700–708.
- Gueriot, M. L. (2000). The ZIP family of metal transporters. *Biochimica et Biophysica Acta*, 1465, 190–198.
- Halima, P., Lin, Y. F., Ahonen, V. H., Blande, D., Clemens, S., Gyenesi, A., ... Tervahauta, A. I. (2014). Gene expression differences between *Noccaea caerulea* ecotypes help to identify candidate genes for metal phytoremediation. *Environmental Science & Technology*, 48, 3344–3353.
- Hammond, J. P., Bowen, H. C., White, P. J., Mills, V., Pyke, K. A., Baker, A. J., ... Broadley, M. R. (2006). A comparison of the *Thlaspi caerulescens* and *Thlaspi arvense* shoot transcriptomes. *New Phytologist*, 170, 239–260.
- Hanikenne, M., Kroymann, J., Trampczynska, A., Bernal, M., Motte, P., Clemens, S., & Krämer, U. (2013). Hard selective sweep and ectopic gene conversion in a gene cluster affording environmental adaptation. *PLoS Genetics*, 9, e1003707.
- Hanikenne, M., & Nouet, C. (2011). Metal hyperaccumulation and hyper-tolerance: A model for plant evolutionary genomics. *Current Opinion in Plant Biology*, 14, 252–259.
- Hanikenne, M., Talke, I. N., Haydon, M. J., Lanz, C., Nolte, A., Motte, P., ... Krämer, U. (2008). Evolution of metal hyperaccumulation required cis-regulatory changes and triplication of *HMA4*. *Nature*, 453, 391–395.
- Höreth, S., Pongrac, P., van Elteren, J. T., Debeljak, M., Vogel-Mikuš, K., Weber, M., ... Clemens, S. (2020). *Arabidopsis halleri* shows hyperbioindicator behaviour for Pb and leaf Pb accumulation spatially separated from Zn. *New Phytologist*, 226, 492–506. <https://doi.org/10.1111/nph.16373>
- Ishimaru, Y., Takahashi, R., Bashir, K., Shimo, H., Senoura, T., Sugimoto, K., ... Nishizawa, N. K. (2012). Characterizing the role of rice NRAMP5 in manganese, iron and cadmium transport. *Scientific Reports*, 2, 286.
- Jefferson, R. A. K., Kavanagh, T. A., & Bevan, M. W. (1987). GUS fusions: β -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO Journal*, 6, 3901–3907.
- Karam, M.-J., Souleman, D., Schwartzman, M.-S., Gallina, S., Spielmann, J., Poncet, C., ... Frérot, H. (2019). Genetic architecture of a plant adaptive trait: QTL mapping of intraspecific variation for tolerance to metal pollution in *Arabidopsis halleri*. *Heredity*, 122, 877–892.
- Krämer, U. (2005). MTP1 mops up excess zinc in *Arabidopsis* cells. *Trends in Plant Science*, 10, 313–315.
- Krämer, U. (2010). Metal hyperaccumulation in plants. *Annual Review of Plant Biology*, 61, 517–534.
- Krämer, U., Talke, I. N., & Hanikenne, M. (2007). Transition metal transport. *FEBS Letters*, 581, 2263–2272.
- Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution*, 35, 1547–1549.
- Lekeux, G., Laurent, C., Joris, M., Jadoul, A., Jiang, D., Bosman, B., ... & Hanikenne, M. (2018). di-Cysteine motifs in the C-terminus of plant HMA4 proteins confer nanomolar affinity for zinc and are essential for HMA4 function in vivo. *Journal of Experimental Botany*, 69, 5547–5560.
- Merlot, S., de la Garcia de la Torre, V. S. G., & Hanikenne, M. (2018). Physiology and molecular biology of trace element hyperaccumulation. G. Echevarria A. van der Ent J. L. Morel & A. Baker (Eds.), *Agromining: Farming for Metals. Extracting unconventional resources using plants*, Mineral Resource Reviews (pp. 93–116). Switzerland: Springer.
- Meyer, C. L., Juraniec, M., Hugué, S., Chaves-Rodriguez, E., Salis, P., Isaure, M. P., ... Verbruggen, N. (2015). Intraspecific variability of cadmium tolerance and accumulation, and cadmium-induced cell wall modifications in the metal hyperaccumulator *Arabidopsis halleri*. *Journal of Experimental Botany*, 66, 3215–3227.
- Meyer, C. L., Pauwels, M., Brisset, L., Gode, C., Salis, P., Bourceaux, A., ... Verbruggen, N. (2016). Potential preadaptation to anthropogenic pollution: Evidence from a common quantitative trait locus for zinc and cadmium tolerance in metalcolous and nonmetalcolous accessions of *Arabidopsis halleri*. *New Phytologist*, 212, 934–943.
- Nei, M., & Kumar, S. (2000). *Molecular evolution and Phylogenetics*. New York, NY: Oxford University Press.
- Nouet, C., Charlier, J.-B., Carnol, M., Bosman, B., Farnir, F., Motte, P., & Hanikenne, M. (2015). Functional analysis of the three *HMA4* copies of the metal hyperaccumulator *Arabidopsis halleri*. *Journal of Experimental Botany*, 66, 5783–5795.
- Palmer, C. M., & Gueriot, M. L. (2009). Facing the challenges of Cu, Fe and Zn homeostasis in plants. *Nature Chemical Biology*, 5, 333–340.
- Pauwels, M., Vekemans, X., Gode, C., Frérot, H., Castric, V., & Saumitou-Laprade, P. (2012). Nuclear and chloroplast DNA phylogeography reveals vicariance among European populations of the model species for the study of metal tolerance, *Arabidopsis halleri* (Brassicaceae). *New Phytologist*, 193, 916–928.
- Reeves, R. D., Baker, A. J., Jaffré, T., Erskine, P. D., Echevarria, G., & van der Ent, A. (2018). A global database for plants that hyperaccumulate metal and metalloid trace elements. *New Phytologist*, 218, 407–411.
- Ricachenevsky, F. K., Menguer, P. K., Sperotto, R. A., & Fett, J. P. (2015). Got to hide your Zn away: Molecular control of Zn accumulation and biotechnological applications. *Plant Science*, 236, 1–17.

- Ricachenevsky, F. K., Punshon, T., Lee, S., Oliveira, B. H. N., Trenz, T. S., Maraschin, F. D. S., ... Guerinot, M. L. (2018). Elemental profiling of rice FOX lines leads to characterization of a new Zn plasma membrane transporter, OsZIP7. *Frontiers in Plant Science*, 9, 865.
- Sasaki, A., Yamaji, N., Yokosho, K., & Ma, J. F. (2012). Nramp5 is a major transporter responsible for manganese and cadmium uptake in rice. *The Plant Cell*, 24, 2155–2167.
- Scheepers, M., Spielmann, J., Boulanger, M., Carnol, M., Bosman, B., De Pauw, E., ... Hanikenne, M. (2020). Intertwined metal homeostasis, oxidative and biotic stress responses in the *Arabidopsis frd3* mutant. *Plant Journal*, 102, 34–52.
- Schvartzman, M. S., Corso, M., Fataftah, N., Scheepers, M., Nouet, C., Bosman, B., ... Hanikenne, M. (2018). Adaptation to high zinc depends on distinct mechanisms in metallicolous populations of *Arabidopsis halleri*. *New Phytologist*, 218, 269–282.
- Shahzad, Z., Gosti, F., Frérot, H., Lacombe, E., Roosens, N., Saumitou-Laprade, P., & Berthomieu, P. (2010). The five *AhMTP1* zinc transporters undergo different evolutionary fates towards adaptive evolution to zinc tolerance in *Arabidopsis halleri*. *PLoS Genetics*, 6, e1000911.
- Sinclair, S. A., & Krämer, U. (2012). The zinc homeostasis network of land plants. *Biochimica et Biophysica Acta*, 1823, 1553–1567.
- Stein, R. J., Horeth, S., de Melo, J. R., Syllwasschy, L., Lee, G., Garbin, M. L., ... Krämer, U. (2017). Relationships between soil and leaf mineral composition are element-specific, environment-dependent and geographically structured in the emerging model *Arabidopsis halleri*. *New Phytologist*, 213, 1274–1286.
- Suryawanshi, V., Talke, I. N., Weber, M., Eils, R., Brors, B., Clemens, S., & Krämer, U. (2016). Between-species differences in gene copy number are enriched among functions critical for adaptive evolution in *Arabidopsis halleri*. *BMC Genomics*, 17, 1034.
- Talke, I. N., Hanikenne, M., & Krämer, U. (2006). Zinc-dependent global transcriptional control, transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of the hyperaccumulator *Arabidopsis halleri*. *Plant Physiology*, 142, 148–167.
- Tiong, J., McDonald, G. K., Genc, Y., Pedas, P., Hayes, J. E., Toubia, J., ... Huang, C. Y. (2014). HvZIP7 mediates zinc accumulation in barley (*Hordeum vulgare*) at moderately high zinc supply. *New Phytologist*, 201, 131–143.
- Uraguchi, S., Weber, M., & Clemens, S. (2019). Elevated root nicotianamine concentrations are critical for Zn hyperaccumulation across diverse edaphic environments. *Plant, Cell & Environment*, 42, 2003–2014.
- van der Ent, A., Baker, A. J., Reeves, R. D., Pollard, A. J., & Schat, H. (2013). Hyperaccumulators of metal and metalloid trace elements: Facts and fiction. *Plant and Soil*, 362, 319–334.
- Verbruggen, N., Hermans, C., & Schat, H. (2009). Molecular mechanisms of metal hyperaccumulation in plants. *New Phytologist*, 181, 759–776.
- Vert, G., Grotz, N., Dédaldéchamp, F., Gaymard, F., Guerinot, M. L., Briat, J.-F., & Curie, C. (2002). IRT1, an *Arabidopsis* transporter essential for iron uptake from the soil and for plant growth. *The Plant Cell*, 14, 1223–1233.
- Weber, M., Harada, E., Vess, C., Roepenack-Lahaye, E. V., & Clemens, S. (2004). Comparative microarray analysis of *Arabidopsis thaliana* and *Arabidopsis halleri* roots identifies nicotianamine synthase, a ZIP transporter and other genes as potential metal hyperaccumulation factors. *Plant Journal*, 37, 269–281.
- Willems, G., Dräger, D. B., Courbot, M., Gode, C., Verbruggen, N., & Saumitou-Laprade, P. (2007). The genetic basis of zinc tolerance in the metallophyte *Arabidopsis halleri* ssp. *halleri* (Brassicaceae): An analysis of quantitative trait loci. *Genetics*, 176, 659–674.
- Willems, G., Frérot, H., Gennen, J., Salis, P., Saumitou-Laprade, P., & Verbruggen, N. (2010). Quantitative trait loci analysis of mineral element concentrations in an *Arabidopsis halleri* x *Arabidopsis lyrata* *petraea* F₂ progeny grown on cadmium-contaminated soil. *New Phytologist*, 187, 368–379.
- Wu, J., Zhao, F.-J., Ghandilyan, A., Logoteta, B., Guzman, M. O., Schat, H., ... Aarts, M. G. (2009). Identification and functional analysis of two ZIP metal transporters of the hyperaccumulator *Thlaspi caerulescens*. *Plant and Soil*, 325, 79–95.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Spielmann J, Ahmadi H, Scheepers M, et al. The two copies of the zinc and cadmium ZIP6 transporter of *Arabidopsis halleri* have distinct effects on cadmium tolerance. *Plant Cell Environ*. 2020;1–15. <https://doi.org/10.1111/pce.13806>