

Supporting evidence for PCB pollution threatening global killer whale population



Joseph G. Schnitzler^{a,*}, Anja Reckendorf^a, Marianna Pinzone^b, Marijke Autenrieth^c,
Ralph Tiedemann^c, Adrian Covaci^d, Govindan Malarvannan^d, Andreas Ruser^a, Krishna Das^b,
Ursula Siebert^a

^a Institute for Terrestrial and Aquatic Wildlife Research, University of Veterinary Medicine Hannover, Foundation, 25761, Büsum, Schleswig-Holstein, Germany

^b Freshwater and Oceanic sciences Unit of reSearch (FOCUS)-Oceanology, University of Liège, Allée du 6 Août, B6C, 4000, Liège, Belgium

^c Unit of Evolutionary Biology/Systematic Zoology, Institute for Biochemistry and Biology, University of Potsdam, Karl-Liebknecht-Str. 24-25, 14476, Potsdam, Germany

^d Toxicological Centre, University of Antwerp, 2610, Wilrijk, Belgium

ARTICLE INFO

Keywords:

Killer whale
PCB
DDT
PBDE
Mercury;
North Sea

ABSTRACT

A recent Science report predicted the global killer whale population to collapse due to PCB pollution. Here we present empirical evidence, which supports and extends the reports' statement. In 2016, a neonate male killer whale stranded on the German island of Sylt. Neonatal attributes indicated an age of at least 3 days. The stomach contained ~20 mL milk residue and no pathologies explaining the cause of death could be detected. Blubber samples presenting low lipid concentrations were analysed for persistent organic pollutants. Skin samples were collected for genotyping of the mitochondrial control region. The blubber PCB concentrations were very high [SPCBs, 225 mg/kg lipid weight (lw)], largely exceeding the PCB toxicity thresholds reported for the onset of immunosuppression [9 mg/kg lw ΣPCB] and for severe reproductive impairment [41 mg/kg lw ΣPCB] reported for marine mammals. Additionally, this individual showed equally high concentrations in *p,p'*-DDE [226 mg/kg lw], PBDEs [5 mg/kg lw] and liver mercury levels [1.1 µg/g dry weight dw]. These results suggest a high placental transfer of pollutants from mother to foetus. Consequently, blubber and plasma PCB concentrations and calf mortality rates are both high in primiparous females. With such high pollutant levels, this neonate had poor prerequisites for survival. The neonate belonged to Ecotype I (generalist feeder) and carried the mitochondrial haplotype 35 present in about 16% of the North Atlantic killer whale from or close to the North Sea. The relevance of this data becomes apparent in the UK West Coast Community, the UK's only resident orca population, which is currently composed of only eight individuals (each four males and females) and no calves have been reported over the last 19 years. Despite worldwide regulations, PCBs persist in the environment and remain a severe concern for killer whale populations, placing calves at high risk due to the mother-offspring PCB-transfer resulting in a high toxicological burden of the neonates.

In a recent Science report, [Desforges et al. \(2018\)](#) predicted the global killer whale population to collapse due to pollution with polychlorinated biphenyls. Using an individual-based model framework and globally available data on PCB concentrations in killer whale tissues, they showed that PCB-mediated effects on reproduction and immune function threaten the long-term viability of > 50% of the world's killer whale populations. Here, we present empirical evidence, which supports and extends the reports' statement.

On 8 February 2016, a neonate male killer whale stranded on the beach of Rantum, German island of Sylt ([Fig. 1](#)). Killer whale strandings are rare and post-mortem examinations on fresh individuals are scarce.

This stranding probably marks the ninth recorded killer whale stranding along German coasts since the mid-1800s ([Reckendorf et al., 2018](#)). The carcass was necropsied approximately 10 h after collection and was examined according to standard protocol ([Siebert et al., 2001](#)). Total length, body weight, different body measurements, girths and blubber thickness (at nine locations along the left flank: dorsal, medial and ventral behind the flipper, in front of as well as behind the dorsal fin) were recorded. The stomach contained ~20 mL milk residue and no pathologies explaining the cause of death could be detected ([Reckendorf et al., 2018](#)). Blubber samples (n = 3) presenting low lipid concentrations [16–22%] were analysed for a suite of persistent organic

* Corresponding author.

E-mail address: joseph.schnitzler@tiho-hannover.de (J.G. Schnitzler).



Fig. 1. Picture of the neonate male killer whale stranded on 8 February 2016 on the beach of Rantum, on the German island of Sylt.

pollutants, such as PCBs (28 congeners), polybrominated diphenyl ethers (PBDEs) and Dichlorodiphenyltrichloroethane and metabolites (DDTs). The method used for sample extraction and clean-up has been previously described by Das et al., 2017. Liver samples were analysed for mercury following the method described by Cransveld et al., 2017. Skin samples were collected caudal to the dorsal fin for genotyping of the mitochondrial control region using established standard odontocete primers ProL and DLH (Tiedemann et al., 1996; Wiemann et al., 2010). For the analysed part of the mtDNA, many sequences from killer whales around the world are available in genetic databases. Hence, such genetic analyses can provide insight into the putative origin of the individuals, enabling them to be assigned to a certain population/ecotype (Foote et al., 2009; Morin et al., 2010).

Neonatal attributes including colouration, freshly healing umbilicus, distinct neonatal papillae on the tongues and partly erupted teeth were found and indicated an age of at least 3 days (Reckendorf et al., 2018). The blubber PCB concentrations were very high [Σ PCBs, average 225 mg/kg lipid weight (lw)], largely exceeding the PCB toxicity thresholds reported for the onset of immunosuppression [9 mg/kg lw Σ PCB] (Kannan et al., 2000) and for severe reproductive impairment [41 mg/kg lw Σ PCB] (Helle et al., 1976) in marine mammals (Jepson et al., 2016) (Fig. 2).

The study of Desforges et al. focuses on a single chemical class (PCBs), but a large number of additional contaminants are present in killer whale tissues. This individual showed equally high concentrations in p,p' -DDE [average 226 mg/kg lw]. In a previous study, adult killer whales from British and Irish coastal waters showed lower p,p' -DDE concentrations [ranging from 7 to 179 mg/kg lw], except one individual that presented a higher p,p' -DDE concentration of 567 mg/kg

lw (McHugh et al., 2007). A number of contaminant studies on marine mammals have determined the p,p' -DDE/ Σ DDT ratio in order to assess the chronology of DDT inputs and state that a ratio > 0.6 is indicative of a stable system with no new DDT inputs (Aguilar, 1984; Tanabe et al., 1997). In the current study, our neonate presented a p,p' -DDE/ Σ DDT ratio of 0.98, suggesting that DDT residues are derived from historic environmental contamination. However, several studies in delphinid species have shown higher transfer rates of p,p' -DDE from mother to calf during lactation than for p,p' -DDT or Σ DDT, which might potentially alter the observed ratio (Aguilar, 1984; Borrell et al., 1995; McKenzie et al., 1997).

Polybrominated diphenyl ethers (PBDEs) are a group of flame-retardant compounds which have been widely used in recent years. Compared to other orcas from UK waters, elevated levels of Σ PBDEs [average 5 mg/kg lw] were found in this individual. In a previous study, four adult killer whales showed lower Σ PBDE concentrations, [ranging from 0.69 to 4.37 mg/kg lw], while one individual presented a higher Σ PBDE concentration [of 23.8 mg/kg lw] (Law et al., 2005).

Environmental mercury (Hg) concentrations have increased over the past 150 years, resulting in over 92% of the mercury body burden in higher trophic level species being of man-made origin (Dietz et al., 2013). Total mercury (THg) analysed in liver samples of the neonate [1.1 μ g/g dry weight dw] revealed concentrations below the threshold [20 μ g/g dw] defined for liver lesion in adult marine mammals (Dietz et al., 2013; Rawson et al., 1993), but close to the mercury-associated neurochemical effect threshold in the 3–5 μ g/g dw range (Basu et al., 2007, 2006; Dietz et al., 2013). More subtle changes were observed at lower Hg concentrations in the brain of toothed whales with associated signs of neurochemical effects. The meaning of such a high Hg concentration in the tissue of a neonate is difficult to interpret, but methyl-Hg is immunotoxic and neurotoxic even at low doses (Basu et al., 2007, 2006; Dietz et al., 2013).

All these compounds may contribute to reproductive and immune failure or other health endpoints not included here, and raise concerns about the potential for other persistent contaminants to generate additional toxicological effects in long-lived, high-trophic level aquatic species. These results suggest a high placental transfer of pollutants from mother to foetus. Consequently, blubber and plasma PCB concentrations and calf mortality rates are both high in primiparous females (Schwacke et al., 2002; Wells et al., 2005). With such high pollutant levels, this neonate had poor prerequisites for survival. The




Fig. 2. Mean Σ PCBs concentrations in neonate and adult female killer whales (from Jepson et al., 2016). The blue bar represents the neonate and the grey bar are adult females. The red dashed line is the equivalent Σ PCBs concentrations threshold (9.0 mg/kg lw) for onset of physiological effects in experimental marine mammal studies (Kannan et al., 2000). The red solid line is the equivalent Σ PCBs concentrations threshold (41.0 mg/kg lw) for the highest PCB toxicity threshold published for marine mammals based on marked reproductive impairment in ringed seals in the Baltic Sea (Helle et al., 1976). Error bars = LowerCI and UpperCI.

neonate belonged to Ecotype I (generalist feeder) and carried the mitochondrial haplotype 35 present in about 16% of the North Atlantic killer whales from or close to the North Sea (*i.e.*, the UK, the Netherlands, Denmark) (Foote et al., 2009; Reckendorf et al., 2018). The relevance of this data becomes apparent in the UK West Coast Community, the UK's only resident orca population, which is currently composed of only eight individuals (each four males and females) and no calves have been reported over the last 19 years (Beck et al., 2014).

Despite worldwide regulations, marine pollutants, including mercury, organochlorine pesticides and PCBs, persist in the environment and remain a severe concern for killer whale populations, placing calves at high risk due to the vertical mother-offspring pollutant-transfer resulting in a high toxicological burden of the neonates.

Acknowledgements

The authors thank all people involved in the reporting, transport and imaging, as well as all ITAW colleagues who helped with necropsy and sampling. K. Das is a Senior F.R.S.-FNRS Research Associate and M. Pinzone received grants from FRIA (Fonds pour la formation à la recherche dans l'industrie et dans l'agriculture). We thank Claus Dethlefs for the picture of the stranded animal. The investigations were partly funded by the Ministry of Energy, Agriculture Environment and Rural Affairs, and the National Park Service of Schleswig-Holstein. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

References

- Aguilar, A., 1984. Relationship of DDE/ΣDDT in marine mammals to the chronology of DDT input into the ecosystem. *Can. J. Fish. Aquat. Sci.* 41, 840–844.
- Basu, N., Kwan, M., Man Chan, H., 2006. Mercury but not organochlorines inhibits muscarinic cholinergic receptor binding in the cerebrum of ringed seals (*Phoca hispida*). *J. Toxicol. Environ. Health Part A* 69, 1133–1143.
- Basu, N., Scheuhammer, A.M., Rouvinen-Watt, K., Grochowina, N., Evans, R.D., O'Brien, M., Chan, H.M., 2007. Decreased N-methyl-D-aspartic acid (NMDA) receptor levels are associated with mercury exposure in wild and captive mink. *Neurotoxicology* 28, 587–593.
- Beck, S., Foote, A.D., Koetter, S., Harries, O., Mandleberg, L., Stevick, P.T., Whooley, P., Durban, J.W., 2014. Using opportunistic photo-identifications to detect a population decline of killer whales (*Orcinus orca*) in British and Irish waters. *J. Mar. Biol. Assoc. U.K.* 94, 1327–1333.
- Borrell, A., Bloch, D., Desportes, G., 1995. Age trends and reproductive transfer of organochlorine compounds in long-finned pilot whales from the Faroe Islands. *Environ. Pollut.* 88, 283–292.
- Cransveld, A., Amouroux, D., Tessier, E., Koutrakis, E., Ozturk, A.A., Bettoso, N., Mieiro, C.L., Bérail, S., Barre, J.P.G., Sturaro, N., Schnitzler, J., Das, K., 2017. Mercury stable isotopes discriminate different populations of European seabass and trace potential Hg sources around Europe. *Environ. Sci. Technol.* 51, 12219–12228. <https://doi.org/10.1021/acs.est.7b01307>.
- Das, K., Malarvannan, G., Dirtu, A., Dulau, V., Dumont, M., Lepoint, G., Mongin, P., Covaci, A., 2017. Linking pollutant exposure of humpback whales breeding in the Indian Ocean to their feeding habits and feeding areas off Antarctica. *Environ. Pollut.* 220, 1090–1099.
- Desforges, J.-P., Hall, A., McConnell, B., Rosing-Asvid, A., Barber, J.L., Brownlow, A., De Guise, S., Eulaers, I., Jepson, P.D., Letcher, R.J., Levin, M., Ross, P.S., Samarra, F., Vikiingson, G., Sonne, C., Dietz, R., 2018. Predicting global killer whale population collapse from PCB pollution. *Science* 361, 1373–1376.
- Dietz, R., Sonne, C., Basu, N., Braune, B., O'Hara, T., Letcher, R.J., Scheuhammer, T., Andersen, M., Andreasen, C., Andriashek, D., 2013. What are the toxicological effects of mercury in Arctic biota? *Sci. Total Environ.* 443, 775–790.
- Foote, A.D., Newton, J., Piertney, S.B., Willerslev, E., Gilbert, M.T.P., 2009. Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Mol. Ecol.* 18, 5207–5217. <https://doi.org/10.1111/j.1365-294X.2009.04407.x>.
- Helle, E., Olsson, M., Jensen, S., 1976. PCB levels correlated with pathological changes in seal uteri. *Ambio* 261–262.
- Jepson, P.D., Deaville, R., Barber, J.L., Aguilar, À., Borrell, A., Murphy, S., Barry, J., Brownlow, A., Barnett, J., Berrow, S., Cunningham, A.A., Davison, N.J., ten Doeschate, M., Esteban, R., Ferreira, M., Foote, A.D., Genov, T., Giménez, J., Loveridge, J., Llavona, À., Martin, V., Maxwell, D.L., Papachlimitzou, A., Penrose, R., Perkins, M.W., Smith, B., de Stephanis, R., Tregenza, N., Verborgh, P., Fernandez, A., Law, R.J., 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Sci. Rep.* 6, 18573.
- Kannan, K., Blankenship, A.L., Jones, P.D., Giesy, J.P., 2000. Toxicity reference values for the toxic effects of polychlorinated biphenyls to aquatic mammals. *Hum. Ecol. Risk Assess.* 6, 181–201.
- Law, R.J., Allchin, C.R., Mead, L.K., 2005. Brominated diphenyl ethers in the blubber of twelve species of marine mammals stranded in the UK. *Mar. Pollut. Bull.* 50, 356–359.
- McHugh, B., Law, R.J., Allchin, C.R., Rogan, E., Murphy, S., Foley, M.B., Glynn, D., McGovern, E., 2007. Bioaccumulation and enantiomeric profiling of organochlorine pesticides and persistent organic pollutants in the killer whale (*Orcinus orca*) from British and Irish waters. *Mar. Pollut. Bull.* 54, 1724–1731.
- McKenzie, C., Rogan, E., Reid, R.J., Wells, D.E., 1997. Concentrations and patterns of organic contaminants in Atlantic white-sided dolphins (*Lagenorhynchus acutus*) from Irish and Scottish coastal waters. *Environ. Pollut.* 98, 15–27.
- Morin, P.A., Archer, F.I., Foote, A.D., Vilstrup, J., Allen, E.E., Wade, P., Durban, J., Parsons, K., Pitman, R., Li, L., 2010. Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Res.* 20, 908–916.
- Rawson, J., Patton, W., Hofman, S., Pietra, G., Johns, L., 1993. Liver abnormalities associated with chronic mercury accumulation in stranded Atlantic bottlenose dolphins. *Ecotoxicol. Environ. Saf.* 25, 41–47.
- Reckendorf, A., Ludes-Wehrmeister, E., Wohlsein, P., Tiedemann, R., Siebert, U., Lehnert, K., 2018. First record of *Halocercus* sp. (Pseudaliidae) lungworm infections in two stranded neonatal orcas (*Orcinus orca*). *Parasitology* 1–5.
- Schwacke, L.H., Voit, E.O., Hansen, L.J., Wells, R.S., Mitchum, G.B., Hohn, A.A., Fair, P. A., 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the Southeast United States Coast. *Environmental Toxicology And Chemistry / SETAC* 21, 2752–2764.
- Siebert, U., Wünschmann, A., Weiss, R., Frank, H., Benke, H., Frese, K., 2001. Post-mortem findings in harbour porpoises (*Phocoena phocoena*) from the German North and Baltic Seas. *J. Comp. Pathol.* 124, 102–114.
- Tanabe, S., Madhusree, B., Öztürk, A.A., Tatsukawa, R., Miyazaki, N., Özdamar, E., Aral, O., Samsun, O., Öztürk, B., 1997. Persistent organochlorine residues in harbour porpoise (*Phocoena phocoena*) from the Black Sea. *Mar. Pollut. Bull.* 34, 338–347.
- Tiedemann, R., Harder, J., Gmeiner, C., Haase, E., 1996. Mitochondrial DNA sequence pattern of harbour porpoises (*Phocoena phocoena*) from the North and the Baltic Sea. *Zeitschrift für Säugetierkunde* 61, 104–111.
- Wells, R.S., Tornero, V., Borrell, A., Aguilar, A., Rowles, T.K., Rhinehart, H.L., Hofmann, S., Jarman, W.M., Hohn, A.A., Sweeney, J.C., 2005. Integrating life-history and reproductive success data to examine potential relationships with organochlorine compounds for bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Sci. Total Environ.* 349, 106–119.
- Wiemann, A., Andersen, L.W., Berggren, P., Siebert, U., Benke, H., Teilmann, J., Lockyer, C., Pawliczka, I., Skóra, K., Roos, A., 2010. Mitochondrial control region and microsatellite analyses on harbour porpoise (*Phocoena phocoena*) unravel population differentiation in the Baltic Sea and adjacent waters. *Conserv. Genet.* 11, 195–211.