




# SARS-CoV-2 in carotid body

Bernard Lambermont<sup>1\*</sup> , Eric Davenne<sup>2</sup>, François Maclot<sup>3</sup> and Philippe Delvenne<sup>2</sup>

© 2021 Springer-Verlag GmbH Germany, part of Springer Nature

A 42-year-old man was admitted to the intensive care unit in October 2020 for coronavirus disease-19 (COVID-19)-related acute respiratory distress syndrome requiring mechanical ventilation. He presented with multi-organ failure and died 41 days later. Comorbidities included a sleeve gastrectomy in 2016 and cholecystectomy in April 2020.

Autopsy revealed diffuse alveolar damage (Fig. 1A) and presence of severe acute respiratory syndrome-coronavirus 2 (SARS-CoV-2) in the carotid body (Fig. 1B, C). Detection of SARS-CoV-2 in the carotid body was performed by real-time reverse transcription polymerase chain reaction (Fig. 1D).

The carotid body plays a role in peripheral arterial chemoreception, in metabolic and immune sensing, and could also be a route of nervous system invasion by SARS-CoV-2. Involvement of the carotid body by SARS-CoV-2 may explain silent hypoxemia and thus could also contribute to increased morbidity and mortality in COVID-19 patients.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 29 December 2020 Accepted: 8 January 2021  
Published online: 22 January 2021

## Author details

<sup>1</sup> Department of Intensive Care, University Hospital of Liege, Liège, Belgium.

<sup>2</sup> Department of Pathology, University Hospital of Liege, Liège, Belgium.

<sup>3</sup> Department of Microbiology, University Hospital of Liege, Liège, Belgium.

## Funding

None.

## Compliance with ethical standards


## Conflicts of interest

The authors declare that they have no conflict of interest.

\*Correspondence: b.lambermont@chuliege.be

<sup>1</sup> Department of Intensive Care, University Hospital of Liege, Liège, Belgium

Full author information is available at the end of the article



**Fig. 1** **A** Post-mortem tissue specimen showing a diffuse alveolar damage and a partially occluded microvascular structure in the lung (arrow) (hematoxylin eosin;  $\times 100$ ); **B** lobules of the left carotid body (open arrows) separated by thin connective tissue septa and located in close proximity of a muscular artery wall structure (asterisk) (hematoxylin eosin;  $\times 100$ ); **C** immunohistochemistry with an antibody against S-100 protein demonstrating many immunostained sustentacular and Schwann cells surrounding neuroendocrine chief cells within the lobules (open arrows) (immunoperoxidase;  $\times 100$ ); **D** real-time reverse transcription polymerase chain reaction using the Taqman<sup>®</sup> technology on LC480 thermocycler (Roche, Switzerland) and showing detection curves (blue curve for patient tissue; red and green curves for positive and negative controls, respectively) of severe acute respiratory syndrome-coronavirus 2 N1/N2 gene transcripts (cycle threshold: 36) after glomus tissue microdissection and RNA extraction from paraffin-embedded sections (tested in triplicate)