



Analysis of Autozygosity Using Whole-Genome Sequence Data of Full-Sib Families in Pikeperch (*Sander lucioperca*)

Lidia De los Ríos-Pérez¹, Tom Druet², Tom Goldammer^{3,4} and Dörte Wittenburg^{1*}

¹Institute of Genetics and Biometry, Research Institute for Farm Animal Biology (FBN), Dummerstorf, Germany, ²Unit of Animal Genomics, GIGA-R & Faculty of Veterinary Medicine, University of Liège, Liège, Belgium, ³Institute of Genome Biology, Research Institute for Farm Animal Biology (FBN), Dummerstorf, Germany, ⁴Molecular Biology and Fish Genetics, Faculty of Agriculture and Environmental Sciences, University of Rostock, Rostock, Germany

OPEN ACCESS

Edited by:

Paul Joseph Wilson,
Trent University, Canada

Reviewed by:

Hans Magnus Gjoen,
Norwegian University of Life Sciences,
Norway
Ole Madsen,
Wageningen University and Research,
Netherlands

*Correspondence:

Dörte Wittenburg
wittenburg@fhn-dummerstorf.de

Specialty section:

This article was submitted to
Evolutionary and Population Genetics,
a section of the journal
Frontiers in Genetics

Received: 30 September 2021

Accepted: 23 December 2021

Published: 17 January 2022

Citation:

De los Ríos-Pérez L, Druet T,
Goldammer T and Wittenburg D (2022)
Analysis of Autozygosity Using Whole-
Genome Sequence Data of Full-Sib
Families in Pikeperch
(*Sander lucioperca*).
Front. Genet. 12:786934.
doi: 10.3389/fgene.2021.786934

Pikeperch (*Sander lucioperca*) has emerged as a high value species to the aquaculture industry. However, its farming techniques are at an early stage and its production is often performed without a selective breeding program, potentially leading to high levels of inbreeding. In this study, we identified and characterized autozygosity based on genome-wide runs of homozygosity (ROH) on a sample of parental and offspring individuals, determined effective population size (N_e), and assessed relatedness among parental individuals. A mean of $2,235 \pm 526$ and $1,841 \pm 363$ ROH segments per individual, resulting in a mean inbreeding coefficient of 0.33 ± 0.06 and 0.25 ± 0.06 were estimated for the progeny and parents, respectively. N_e was about 12 until four generations ago and at most 106 for 63 generations in the past, with varying genetic relatedness amongst the parents. This study shows the importance of genomic information when family relationships are unknown and the need of selective breeding programs for reproductive management decisions in the aquaculture industry.

Keywords: autozygosity, genomic inbreeding, FROH, pikeperch, selection, animal breeding

INTRODUCTION

Pikeperch (*Sander lucioperca*) is one of the five species of the genus *Sander* from the Percidae family. It is a fresh and brackish water fish with a native distribution in Eastern Europe and Western Asia, inhabiting the drainages basins of the Caspian, Baltic, Black, Aral, North, and Aegean Sea basins (Kottelat and Freyhof, 2007; Kestemont et al., 2015) and has been introduced to other countries from Europe, Asia, and North America. The origins of pikeperch production date back to the 19th century in Central and Eastern Europe, where pikeperch was farmed in small quantities with carp (*Cyprinus carpio*) in earthen ponds. In the early 20th century, the production of pikeperch as restocking material for open waters began, being produced as monoculture or polyculture with carp. In the second half of the 20th century, pikeperch production in extensive systems started to develop in Western Europe, and by the beginning of the 21st century, its production in recirculating aquaculture systems (RAS) was established (FAO, 2012). Although pikeperch farming techniques are at an early stage in Europe, its growing consumer demand has placed it as a candidate for aquaculture diversification, initiating over the last decades research projects to accelerate its production in intensive inland aquaculture systems (Zakęś et al., 2013; Kristan et al., 2018; Nguinkal et al., 2019;

Thomas et al., 2020). Particularly in Germany, where pikeperch is also a native species, fish farmers and politics push to establish pikeperch as a new aquaculture species and thus to reduce the import from wild catches. However, to improve its production and economic profit, the use of domesticated stocks and the development of systematic and efficient breeding strategies are necessary.

The use of breeding programs implies the use of selective mating, where only some individuals will be chosen to transmit their genes to the next generation, which in the long term may lead to a change in the gene frequencies and a reduction of the effective population size (N_e). This reduction implies an increased probability of mating between relatives. Populations undergoing selective mating experience some degree of inbreeding over time. The inbreeding coefficient (F) is the probability that two alleles at a randomly chosen locus are identical by descent (IBD), and it refers to the amount of inbreeding accumulated from a specific point in ancestry of the population (e.g., Lewontin et al., 1968). Inbreeding is commonly associated with the reduction of mean phenotypic values of fitness traits, known as inbreeding depression (Falconer and Mackay, 1996; Lynch and Walsch, 1998). Thus, measuring individual inbreeding is critical in populations undergoing selection programs (Chavanne et al., 2016).

Prior to genomic information becoming available, pedigree-based relationships were used for estimating inbreeding. Pedigree inbreeding (F_{PED}) is based on Mendelian sampling and does not consider concrete recombination events during meiosis. Furthermore, the assumption that the founder individuals are unrelated, parentage errors, and incomplete information lead to biased inbreeding estimates (Purfield et al., 2012; Forutan et al., 2018). Nowadays, genotyping technologies allow for genomic-based or combined approaches to more accurately estimate inbreeding (Zhang et al., 2015a). One genomic approach is through runs of homozygosity (ROH). ROH are continuous homozygous segments of the genome which can be identical by state or identical by descent. The latter are mostly interpreted as homozygous-by-descent (HBD) or autozygous segments, and can, for instance, be identified through the analysis of high-density single nucleotide polymorphism (SNP) panels (McQuillan et al., 2008; Ferenčaković et al., 2013b). ROH allow differentiating between recent and more ancient inbreeding. Long ROH segments indicate recent inbreeding, where recombination has had fewer opportunities to break up the ROH, whereas short segments trace back to more remote ancestors with a higher number of recombination events (Thompson, 2013). The proportion of the genome lying in HBD segments provides an estimate of individual inbreeding (Kardos et al., 2015), termed F_{ROH} below.

In the present study, we used an ultra-high density SNP panel of a pikeperch sample of two generations to identify and characterize autozygosity based on genome-wide ROH, to determine N_e from linkage disequilibrium in the parental generation, and to assess relatedness among parental individuals. Since, to the best of our knowledge, neither a genome-based nor pedigree-based breeding program has been established yet for pikeperch in Germany, hence we expected to

TABLE 1 | Matings and number of individuals randomly sampled from a common tank.

Family	Sire Id	Dam Id	Number of progeny
1	1	2	28
2	3	4	89
3	5	6	3
4	7	8	223
5	9	10	14
6	9	11	6

observe an increased level of individual autozygosity and relatedness among individuals. The application of genomic tools helps discovering non-optimal breeding decisions in the past. Our results will contribute to setting the basis for the design of breeding strategies towards the improvement of the aquaculture production in pikeperch.

METHODS

Data

We analyzed genotype data of pikeperch families initially produced for a linkage analysis (De los Ríos-Pérez et al., 2020). The production was based on F0 individuals without information about pedigree and former breeding practices. About 20 matings were performed. Out of the entire pool of progeny, about 2000 individuals were chosen according to positive mass selection (excluding the very early growing individuals). Following the same strategy, about 200 breeding candidates were selected from this resource, constituting the parent (F1) generation. Eventually, 18 individuals were selected on visual inspection, and 7 matings were performed with a male:female ratio of 2:1 and 1:1 (one male was used twice); one nest needed to be discarded. In tanks with a 2:1 male:female ratio, only one male was expected to fertilize the eggs. The 18 selected parents (11 males and 7 females) and 375 progeny were tissue sampled for DNA extraction and whole genome paired-end sequencing, followed by genotyping for SNP identification. Pedigree reconstruction was performed to identify the successful male in 2:1 matings and the progeny belonging to each mating. Further processing of the data yielded a final panel of 992,340 genome-wide SNPs from 11 parents (5 males and 6 females) and 363 progeny of 6 families. The number of progeny corresponding to each mating is shown in **Table 1**.

Our analyses considered only markers within the 24 chromosomes of pikeperch, excluding those positioned in unplaced scaffolds, yielding a total of 992,313 SNPs on 2,709.64 centiMorgan (cM) or 896.05 Mega base pairs (Mb) on a sex-averaged map. Since the data came from an inbred population, no Hardy-Weinberg equilibrium or minor allele frequency filtering was performed.

Runs of Homozygosity and Autozygosity

Runs of homozygosity analysis was performed with the RZooRoH package version 0.2.3 in R (Druet and Gautier, 2017; Bertrand et al., 2019). This package identifies HBD segments and estimates

individual autozygosity. The model used by RZooRoH is a hidden Markov model (HMM) that partitions the genome-wide individual autozygosity into different HBD classes with pre-defined rates of ancestry change (R_k) that are related to the age of inbreeding. In RZooRoH, the marker positions should ideally be provided in genetic distances in cM units. Alternatively, physical distances in base pairs (bp) can be used, assuming 1 Mb = 1 cM. In pikeperch, this assumption fails, as shown in the study performed by De los Ríos-Pérez et al. (2020). Therefore, we transformed the marker positions from Mb to cM by multiplying by 3.024 (= 2,709.64 cM/896.05 Mb). We analyzed progeny and parents separately, both based on the allele depth of the markers. To avoid bias due to the large difference of family sizes, only the allele frequencies of the 18 individuals used as putative parents were considered in the model parameters. We applied a “mixKR model” with 14 HBD classes with rate $R_k = 2^k$ (i.e., $2^1, 2^2, 2^3, \dots, 2^{14}$), related to the age of inbreeding, and one non-HBD class. Hence, HBD segments can have different lengths depending on HBD (age) class. The number of heterozygous SNPs allowed in a segment depends on a small probability justifying genotyping errors; as recommended we considered a small $seqerror = 0.005$. Consequently, too many heterozygous loci will cut a segment into shorter ROH. Eventually, total autozygosity was estimated for progeny and parents as accumulated autozygosity over all HBD classes, i.e.,

$$F_{ROH} = \sum_{k=1}^{14} F_{Rk}$$

where F_{Rk} is the proportion of genome attributed to ROH segments in the k th HBD class.

Effective Population Size

N_e was derived from linkage disequilibrium between SNP pairs in parents according to Santiago et al. (2020), as implemented in the software package GONE. This approach enables the estimation of temporal N_e and accounts for accumulated drift effects but it circumvents the restriction that N_e follows a linear trend back in time as, for instance, in Corbin et al. (2012). For this analysis, genotype data of the parents and progeny were phased with ShapeIT version 2 (Delaneau et al., 2013). In GONE, we neglected the selection of a mapping function because only recombination rates less than 0.05 (default setting) were taken into account.

Additionally, we performed nucleotide diversity analysis with HBD segments excluded similarly to “diversity outside ROH” used in Bosse et al. (2012), reflecting more ancient population size. Nucleotide diversity (π) was calculated in windows of 10 kilobases (kb) using VCFtools version 0.1.16 (Danecek et al., 2011) and averaged over the genome.

Genetic Relatedness

Estimation of genetic relatedness between pairs of parental individuals was performed with ngsRelate version 2 (Hanghøj et al., 2019). This procedure utilizes identity-by-descent of SNP alleles and provides a scaled measure of relatedness in $[0,1]$ (Hedrick et al., 2015).

RESULTS

Runs of Homozygosity

A total of 811,440 ROH segments were identified in the progeny generation, with a mean of $2,235.37 \pm 525.55$ segments per individual and ranging from 741 to 3,279. The mean ROH length was 0.38 ± 1.59 cM, with the longest segment being identified on chromosome 15 with 75.78 cM and 25,207 SNPs. A total of 3,085 ROH segments were longer than 10 cM.

In the parental generation, 33,146 ROH segments were identified with a mean of $1,841.44 \pm 362.92$ segments per individual, and ranging from 1,261 to 2,369. The mean ROH length was 0.34 ± 1.01 cM, with the longest segment identified in chromosome 17 with 25.31 cM and 10,996 SNPs. The ROH analysis for different age-related inbreeding classes showed that in both generations the highest number of segments corresponded to short segments ($R_k = 2,048$) from ancient inbreeding events 1,024 generations ago ($\approx R_k/2$) (Druet and Gautier, 2017; Bertrand et al., 2019). No segments were found for the oldest classes (i.e., $R_k \geq 8,192$ and $R_k \geq 4,096$ for the progeny and parental generations, respectively). The distribution of ROH over HBD classes in parents and progeny is summarized in **Table 2**. The partitioning of the genome in the different HBD classes is shown in **Figure 1**.

The chromosome-wise analysis of the average number of ROH per individual showed a dependence of ROH segments assigned to the different HBD classes on chromosome length in both generations, see **Figure 2**. As expected, the shorter a chromosome was, the less ROH were observed.


Autozygosity

Mean of estimated F_{ROH} was 0.33 ± 0.06 (0.25 ± 0.06) with a range from 0.09 to 0.52 (0.18–0.35) for the progeny (parent) generation. Mean of accumulated inbreeding coefficients over HBD classes for progeny and parental generations are shown in **Figure 3**. HBD classes $R_k = 8$ to 64 contributed most to total autozygosity in progeny and $R_k = 32$ to 512 in parents (also see **Table 2**). We considered HBD classes $R_k \leq 4$ as “recent” inbreeding. $F_{Rk \leq 4}$ ranged from 0 to 0.16 with a mean of $7.82 \times 10^{-3} \pm 2.15 \times 10^{-2}$ for the progeny generation. With progeny of families one and two excluded, recent inbreeding ranged only from 0 to 0.10. The increased occurrence of ROH segments in the most recent HBD classes for the progeny of families one and two suggest that the parents of each family were highly related (**Figure 1**). The parents had a mean $F_{Rk \leq 4}$ of $2.25 \times 10^{-4} \pm 4.59 \times 10^{-4}$ ranging from 0 to 1.57×10^{-3} . For parents, we found that inbreeding was relatively old: F_{ROH} was about 0.15 with respect to 128 generations in the past but moderately low at 0.05 if 16 generations were considered. However, since the parent generation is rather small, this outcome may be influenced by sampling variation and shall not be overvalued.

The progeny with the lowest $F_{ROH} = 0.09$ belonged to family 4 which had the largest sample size. With the single outlier excluded, the average inbreeding of its siblings was 0.34 ± 0.05 , ranging between 0.25 and 0.52. No concrete reason was

TABLE 2 | Descriptive statistics of runs of homozygosity (ROH) and autozygosity contributions for the different age-related homozygous-by-descent (HBD) classes with pre-defined rates of ancestry change (R_k) for progeny and parents. cM, centiMorgan; CUM, cumulative autozygosity.

Generation	HBD class (R_k)	Age of inbreeding event ($\approx R_k/2$)	Number of ROH	No. of individuals with ≥ 1 ROH	Number of SNPs			Length (cM)			Proportion of genome in HBD class			
					Min	Max	Mean	Min	Max	Mean	Min	Max	Mean	CUM
Progeny	2	1	10	7	14,147	27,344	20,019	34.87	71.16	61.13	1.81×10^{-19}	3.56×10^{-2}	9.55×10^{-4}	9.55×10^{-4}
	4	2	135	33	2,396	27,344	13,910	7.87	75.78	42.79	5.12×10^{-18}	1.61×10^{-1}	6.87×10^{-3}	7.82×10^{-3}
	8	4	2,095	86	49	28,236	5,672	0.29	74.91	16.77	3.04×10^{-15}	2.77×10^{-1}	3.02×10^{-2}	3.80×10^{-2}
	16	8	2,971	69	33	16,642	2,852	0.38	45.81	8.18	7.69×10^{-10}	2.52×10^{-1}	2.45×10^{-2}	6.25×10^{-2}
	32	16	18,094	293	27	7,332	1,502	0.29	19.06	4.41	1.66×10^{-6}	1.76×10^{-1}	7.15×10^{-2}	1.34×10^{-1}
	64	32	24,427	248	9	3,869	757	0.23	9.69	2.07	1.08×10^{-4}	1.49×10^{-1}	5.42×10^{-2}	1.88×10^{-1}
	128	64	5,124	38	20	2,861	400	0.16	3.40	1.07	5.06×10^{-11}	9.73×10^{-2}	8.20×10^{-3}	1.96×10^{-1}
	256	128	78,885	305	9	1,754	224	0.08	1.91	0.58	1.33×10^{-7}	9.30×10^{-2}	4.04×10^{-3}	2.37×10^{-1}
	512	256	57,370	231	10	1,635	129	0.08	0.90	0.32	2.95×10^{-6}	7.31×10^{-2}	2.70×10^{-2}	2.64×10^{-1}
	1,024	512	7,319	28	1	657	79	1.00×10^{-6}	0.48	0.18	1.03×10^{-15}	4.21×10^{-2}	2.82×10^{-3}	2.67×10^{-1}
	2,048	1,024	606,219	362	1	488	36	1.00×10^{-6}	0.46	0.07	9.66×10^{-3}	1.05×10^{-1}	6.07×10^{-2}	3.27×10^{-1}
	4,096	2,048	8,791	35	1	155	17	1.00×10^{-6}	0.04	0.01	9.13×10^{-14}	3.86×10^{-2}	6.58×10^{-3}	3.34×10^{-1}
Parents	2	1	0	0	0	0	0	0	0	0	8.38×10^{-15}	4.26×10^{-4}	6.42×10^{-5}	6.42×10^{-5}
	4	2	0	0	0	0	0	0	0	0	1.18×10^{-15}	1.14×10^{-3}	1.61×10^{-4}	2.25×10^{-4}
	8	4	0	0	0	0	0	0	0	0	2.80×10^{-14}	3.49×10^{-3}	5.08×10^{-4}	7.34×10^{-4}
	16	8	3	3	4,977	10,186	7,041	17.09	24.14	21.24	9.20×10^{-8}	1.34×10^{-2}	2.72×10^{-3}	3.45×10^{-3}
	32	16	441	18	35	10,996	2,044	0.76	25.31	6.41	1.78×10^{-2}	1.48×10^{-1}	5.03×10^{-2}	5.38×10^{-2}
	64	32	953	12	16	3,821	888	0.42	9.48	2.60	5.46×10^{-23}	1.10×10^{-1}	4.18×10^{-2}	9.59×10^{-2}
	128	64	818	8	24	1,931	492	0.20	4.01	1.35	2.82×10^{-10}	7.81×10^{-2}	2.32×10^{-2}	1.19×10^{-1}
	256	128	2,660	13	11	1,331	251	0.12	1.88	0.66	6.95×10^{-5}	8.49×10^{-2}	3.52×10^{-2}	1.54×10^{-1}
	512	256	3,060	12	13	932	137	0.05	0.80	0.35	2.39×10^{-8}	5.25×10^{-2}	2.69×10^{-2}	1.81×10^{-1}
	1,024	512	1,227	6	12	324	88	0.02	0.40	0.21	2.99×10^{-11}	3.22×10^{-2}	8.40×10^{-3}	1.89×10^{-1}
	2,048	1,024	23,984	18	2	466	39	0.00	0.43	0.07	2.87×10^{-2}	7.58×10^{-2}	5.69×10^{-2}	2.46×10^{-1}
	4,096	2,048	0	0	0	0	0	0	0	0	8.48×10^{-37}	1.05×10^{-2}	9.86×10^{-4}	2.47×10^{-1}



found to explain the difference in inbreeding coefficients between the outlier and its siblings.

Effective Population Size

Estimates of historical N_e are shown in **Figure 4**. Until four generations in the past, estimated N_e was extremely low at about 12 and increased steadily towards $N_e = 106$ for 63 generations ago. For more than 150 generations ago, N_e stabilizes at about 12. The particularly low N_e for the recent history agreed with our expectation on reproduction practices discarding relatedness.

Nucleotide diversity was on average $2.44 \times 10^{-4} \pm 2.13 \times 10^{-4}$ per 10 kb window among all individuals.

Genetic Relatedness



Most parental individuals were almost unrelated with regard to genotype data (**Figure 5**). A suspiciously high coefficient of relatedness was observed for two individuals, one of them was father of family three, with no obvious explanation. A group of four individuals, being parents of the progeny generation, revealed medium relatedness of about 0.5 among each other indicating common ancestor(s). Against our assumption from

autozygosity analysis, a closer relationship between parents within family one and two was not confirmed; relatedness was 0.14 and 0.08, respectively.

DISCUSSION

We assessed genome-wide autozygosity in parents and progeny of a pikeperch population based on an ultra-high density SNPs panel. Our study represents the first investigation of inbreeding based on ROH in this species. Considering progeny and parents, our results showed that the majority of autozygosity is associated with ancestors over 16 generations ago ($R_k \geq 32$). According to pikeperch history, the origins of its production dates back to the 19th century (FAO, 2012). Assuming a generation interval of 3–5 years, pikeperch production origins took place approximately 43–73 generations ago.

Genome-wide ROH analysis has been performed in multiple livestock species, such as cattle (Ferenčaković et al., 2013a; Peripolli et al., 2020), chicken (Dementieva et al., 2020), and pig (Bosse et al., 2015; Shi et al., 2020). However, only few studies are available on aquaculture



species. ROH reported in rainbow trout (*Oncorhynchus mykiss*) showed a moderate level of autozygosity in several lines, with F_{ROH} ranging from 10.0% in an unselected experimental line to 19.5% in a commercial line (D'Ambrosio et al., 2019). In turbot (*Scophthalmus maximus*), large differences were found in ROH distribution between wild and domestic populations, with wild populations showing only short length segments (<2 Mb) (Aramburu et al., 2020). In coho salmon (*Oncorhynchus kisutch*), an analysis of ROH patterns between two pure lines and one admixed line showed a larger number and greater mean length of ROH in the pure lines (Yoshida et al., 2020). More precisely, mean \pm SD F_{ROH} for two pure lines and one admixed line were of $14.2 \pm 3.8\%$, $15.2 \pm 4.4\%$ and $0.4 \pm 0.3\%$, respectively. As the specification of length of ROH varied among literature, a thorough comparison between studies is actually not possible. Nevertheless, in our study, F_{ROH} estimates for the progeny

generation were found to range between 25 and 52%, which is considerably higher than the F_{ROH} estimates observed in aquaculture species undergoing a selection program. Unlike SNP array data used in previous studies, sequence data can also capture ancient inbreeding (through small HBD segments), consequently leading to higher and more precise estimates of F_{ROH} (Zhang et al., 2015a). Restricted to HBD classes with $R_k \leq 128$ (mean segment length was ≥ 1 cM comparable to array data), mean autozygosity was then 19% among progeny. Evaluations and comparisons between ROH (in terms of IBS) and HBD methods showed that at high marker density, the estimated inbreeding coefficients are highly correlated and that the longest ROH are captured by both methods (Druet and Gautier, 2017; Solé et al., 2017; Alemu et al., 2021). The main difference is related to the capacity of the methods to measure autozygosity associated with the shortest HBD segments which capture background LD.




FIGURE 4 | Estimates of historical N_e .




FIGURE 5 | Relatedness matrix for the parent population. Column and row names refer to family assignment. In case of “noFam,” an individual was not used for mating; extension -f indicates females and -m males.

The observed nucleotide diversity was very low compared to mammals, such as cattle (Zhang et al., 2015b) or pigs (Bosse et al., 2012). Such low values have also been observed in other fish species, for instance, salmon (Kijas et al., 2018) and catfish (Jensen et al., 2021). Based on the relationship $\pi = 4 N_e \nu$ with a mutation rate ν per sequence (Nei and Takahata, 1993), this outcome also strengthens the indication that past N_e size must have been rather small in pikeperch.

Estimates of N_e for recent generations (≤ 4 generations ago) were found well below the minimal recommended value of 50 individuals which are necessary to avoid inbreeding depression in the short term (FAO, 1998), and also lower than those estimated for other aquaculture species (e.g., Saura et al., 2021). In rainbow trout, N_e estimates in four commercial

lines mainly selected for growth ranged from 37 to 48 for one generation in the past (D’Ambrosio et al., 2019). In a study performed in channel catfish (*Ictalurus punctatus*), N_e was calculated based on mean linkage disequilibrium between adjacent markers and on inbreeding F_{PED} , where both methods gave similar results, i.e., 27 and 28, respectively (Garcia et al., 2018). A study performed in Atlantic salmon (*Salmo salar*) showed N_e estimates ranging from 15 to 72 few generations ago of three populations under study (Barria et al., 2018). Comparison between our results and those obtained in other aquaculture species following breeding programs makes doubtlessly clear that pedigree information and inbreeding levels should be closely monitored. Only then, breeders can ensure sufficient genetic diversity so that a population can adapt to future breeding goals and the accumulation of detrimental effects associated with inbreeding is avoided.

The results presented were based on RZooRoH’s mixKR model with parameters $K = 15$ and $R = 2$. We additionally evaluated a variety of model parameters to identify a statistically “optimal” number of HBD classes. To this end, models with $K = 2$ to $K = 18$ and $R = 2$ to $R = 10$ were compared using the Bayesian information criterion on individual likelihoods (Druet and Gautier, 2017). With the optimal parameter setting $K = 5$ and $R = 7$, estimates of total autozygosity were almost unaffected (Supplementary Table S1). We decided to present results allowing for a finer retrospect on demographical events.

Implications of the Study

In the aquaculture industry, the large number of progeny per female might suggest the need of keeping only few broodstock to satisfy production requirements. Inadequate numbers of broodstock lead to a population susceptible to inbreeding. In the present study, the data was obtained from an aquaculture facility without a breeding program or an animal record of ancestry. This is reflected in the high levels of inbreeding obtained in the population, making evident the importance of knowing the genetic diversity and kinship relationships for reproductive management decisions. Therefore, the implementation of a breeding program is highly recommended to increase the genetic values of the population while controlling inbreeding. Additional measures might be required, such as importing pikeperch from other populations that contribute additional genetic variation. Furthermore, the integration of genomic tools in the design of a breeding program, such as SNP panels for parentage assignment and estimation of breeding values, will allow to better estimate genetic parameters, particularly inbreeding, in a population (Dekkers, 2012; Yáñez et al., 2015).

DATA AVAILABILITY STATEMENT

Publicly available datasets were analyzed in this study. This data can be found here: The raw sequencing data are

available at the NCBI Sequence Read Archive (SRA) under Accession Number PRJNA626522. The genome assembly of Sander lucioperca (SLUC_FBN_1.2) is available at the NCBI GenBank under the Accession Number GCA_008315115.2.

ETHICS STATEMENT

The animal study was reviewed and approved by the Committee on the Ethics of Animal Experiments of Mecklenburg-Western Pomerania (Landesamt für Gesundheit und Soziales LAGuS). Approval ID: 7221.3-1-009/19.

AUTHOR CONTRIBUTIONS

LdlR-P performed data analysis and wrote the manuscript. TD developed the software used in this study and contributed to discussions of the manuscript. TG developed the idea for the collaborative project and raised project funding. DW conceived and supervised the study. All authors have read and approved the manuscript.

REFERENCES

- Alemu, S. W., Kadri, N. K., Harland, C., Faux, P., Charlier, C., Caballero, A., et al. (2021). An Evaluation of Inbreeding Measures Using a Whole-Genome Sequenced Cattle Pedigree. *Heredity* 126, 410–423. doi:10.1038/s41437-020-00383-9
- Aramburu, O., Ceballos, F., Casanova, A., Le Moan, A., Hemmer-Hansen, J., Bekkevold, D., et al. (2020). Genomic Signatures after Five Generations of Intensive Selective Breeding: Runs of Homozygosity and Genetic Diversity in Representative Domestic and Wild Populations of Turbot (*Scophthalmus maximus*). *Front. Genet.* 11, 1–14. doi:10.3389/fgene.2020.00296
- Barria, A., López, M. E., Yoshida, G., Carvalheiro, R., Lhorente, J. P., and Yáñez, J. M. (2018). Population Genomic Structure and Genome-wide Linkage Disequilibrium in Farmed Atlantic Salmon (*Salmo salar* L.) Using Dense SNP Genotypes. *Front. Genet.* 9, 649. doi:10.3389/fgene.2018.00649
- Bertrand, A. R., Kadri, N. K., Flori, L., Gautier, M., and Druet, T. (2019). RZooRoH: An R Package to Characterize Individual Genomic Autozygosity and Identify Homozygous-by-descent Segments. *Methods Ecol. Evol.* 10, 860–866. doi:10.1111/2041-210X.13167
- Bosse, M., Megens, H.-J., Madsen, O., Crooijmans, R. P. M. A., Ryder, O. A., Austerlitz, F., et al. (2015). Using Genome-wide Measures of Coancestry to Maintain Diversity and Fitness in Endangered and Domestic Pig Populations. *Genome Res.* 25, 970–981. doi:10.1101/gr.187039.114
- Bosse, M., Megens, H.-J., Madsen, O., Paudel, Y., Frantz, L. A. F., Schook, L. B., et al. (2012). Regions of Homozygosity in the Porcine Genome: Consequence of Demography and the Recombination Landscape. *Plos Genet.* 8, e1003100. doi:10.1371/journal.pgen.1003100
- Chavanne, H., Janssen, K., Janssen, K., Hofherr, J., Contini, F., Haffray, P., et al. (2016). A Comprehensive Survey on Selective Breeding Programs and Seed Market in the European Aquaculture Fish Industry. *Aquacult Int.* 24, 1287–1307. doi:10.1007/s10499-016-9985-0
- Corbin, L. J., Liu, A. Y. H., Bishop, S. C., and Woolliams, J. A. (2012). Estimation of Historical Effective Population Size Using Linkage Disequilibria with Marker Data. *J. Anim. Breed. Genet.* 129, 257–270. doi:10.1111/j.1439-0388.2012.01003.x

FUNDING

The project was funded by the European Maritime and Fisheries Fund (EMFF) and the Ministry of Agriculture and Environment of Mecklenburg-Western Pomerania, Germany; grant MV-II.1-RN-001. The publication of this article was funded by the Open Access Fund of the FBN.

ACKNOWLEDGMENTS

We thank Marcus Stüeken from the state's aquaculture facility (State Research Institute for Agriculture and Fisheries in Hohen Wangelin, Mecklenburg-Western Pomerania, Germany) for proving the samples and for assistance beyond. We also thank the reviewers for their helpful comments. TD is Senior Research Associate from the F.R.S.-FNRS.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2021.786934/full#supplementary-material>

- D'Ambrosio, J., Phocas, F., Haffray, P., Bestin, A., Brard-Fudulea, S., Poncet, C., et al. (2019). Genome-wide Estimates of Genetic Diversity, Inbreeding and Effective Size of Experimental and Commercial Rainbow trout Lines Undergoing Selective Breeding. *Genet. Sel. Evol.* 51, 1–15. doi:10.1186/s12711-019-0468-4
- Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., et al. (2011). The Variant Call Format and VCFtools. *Bioinformatics* 27, 2156–2158. doi:10.1093/bioinformatics/btr330
- De los Ríos-Pérez, L., Nguinkal, J. A., Verleih, M., Rebl, A., Brunner, R. M., Klosa, J., et al. (2020). An Ultra-high Density SNP-Based Linkage Map for Enhancing the Pikeperch (*Sander lucioperca*) Genome Assembly to Chromosome-Scale. *Sci. Rep.* 10, 22335. doi:10.1038/s41598-020-79358-z
- Dekkers, J. C. (2012). Application of Genomics Tools to Animal Breeding. *Cg* 13, 207–212. doi:10.2174/138920212800543057
- Delaneau, O., Howie, B., Cox, A. J., Zagury, J.-F., and Marchini, J. (2013). Haplotype Estimation Using Sequencing Reads. *Am. J. Hum. Genet.* 93, 687–696. doi:10.1016/j.ajhg.2013.09.002
- Dementieva, N. V., Kudinov, A. A., Larkina, T. A., Mitrofanova, O. V., Dysin, A. P., Terletsky, V. P., et al. (2020). Genetic Variability in Local and Imported Germplasm Chicken Populations as Revealed by Analyzing Runs of Homozygosity. *Animals* 10, 1887–1889. doi:10.3390/ani10101887
- Druet, T., and Gautier, M. (2017). A Model-Based Approach to Characterize Individual Inbreeding at Both Global and Local Genomic Scales. *Mol. Ecol.* 26, 5820–5841. doi:10.1111/mec.14324
- Falconer, D. S., and Mackay, T. F. C. (1996). *Introduction to Quantitative Genetics*. 4th ed. Essex, UK: Longman Group.
- FAO (2012). "Sander lucioperca. Cultured Aquatic Species Information Programme," in *FAO Fisheries and Aquaculture Department [online]*. Editor Z. Zakęš (Rome, Italy). Available at: https://www.fao.org/fishery/en/culturedspecies/sander_lucioperca/en (Accessed December 29, 2020).
- FAO (1998). *Secondary Guidelines for Development of National Farm Animal Genetic Resources Management Plans: Management of Small Populations at Risk*. Rome, Italy: Food and Agriculture Organization of the United Nations. Available at: <http://www.fao.org/3/w9361e/w9361e.pdf>.
- Ferenčaković, M., Hamzić, E., Gredler, B., Solberg, T. R., Klemetsdal, G., Curik, I., et al. (2013a). Estimates of Autozygosity Derived from Runs of Homozygosity: Empirical Evidence from Selected Cattle Populations. *J. Anim. Breed. Genet.* 130, 286–293. doi:10.1111/jbg.12012

- Ferenčaković, M., Sölkner, J., and Curik, I. (2013b). Estimating Autozygosity from High-Throughput Information: Effects of SNP Density and Genotyping Errors. *Genet. Sel. Evol.* 45, 1–9. doi:10.1186/1297-9686-45-42
- Forutan, M., Ansari Mahyari, S., Baes, C., Melzer, N., Schenkel, F. S., and Sargolzaei, M. (2018). Inbreeding and Runs of Homozygosity before and after Genomic Selection in North American Holstein Cattle. *BMC Genomics* 19, 98. doi:10.1186/s12864-018-4453-z
- García, A. L. S., Bosworth, B., Waldbieser, G., Misztal, I., Tsuruta, S., and Lourenco, D. A. L. (2018). Development of Genomic Predictions for Harvest and Carcass Weight in Channel Catfish. *Genet. Sel. Evol.* 50, 66. doi:10.1186/s12711-018-0435-5
- Hanghøj, K., Moltke, I., Andersen, P. A., Manica, A., and Korneliussen, T. S. (2019). Fast and Accurate Relatedness Estimation from High-Throughput Sequencing Data in the Presence of Inbreeding. *Gigascience* 8, 1–9. doi:10.1093/gigascience/giz034
- Hedrick, P. W., Lacy, R. C., and Baker, C. S. (2015). Measuring Relatedness between Inbred Individuals. *J. Hered.* 106, 20–25. doi:10.1093/jhered/esu072
- Jensen, A., Lillie, M., Bergström, K., Larsson, P., and Höglund, J. (2021). Whole Genome Sequencing Reveals High Differentiation, Low Levels of Genetic Diversity and Short Runs of Homozygosity Among Swedish Wels Catfish. *Heredity* 127, 79–91. doi:10.1038/s41437-021-00438-5
- Kardos, M., Luikart, G., and Allendorf, F. W. (2015). Measuring Individual Inbreeding in the Age of Genomics: Marker-Based Measures Are Better Than Pedigrees. *Heredity* 115, 63–72. doi:10.1038/hdy.2015.17
- Kestemont, P., Dabrowski, K., and Summerfelt, R. C. (2015). *Biology and Culture of Percid Fishes*. Dordrecht: Springer. doi:10.1007/978-94-017-7227-3
- Kijas, J., McWilliam, S., Naval Sanchez, M., Kube, P., King, H., Evans, B., et al. (2018). Evolution of Sex Determination Loci in Atlantic Salmon. *Sci. Rep.* 8, 1–11. doi:10.1038/s41598-018-23984-1
- Kottelat, M., and Freyhof, J. (2007). *Handbook of European Freshwater Fishes*. Cornol, Switzerland and Freyhof, Berlin, Germany: Kottelat.
- Kristan, J., Zarski, D., Blecha, M., Policar, T., Malinovsky, O., Samarina, A. M., et al. (2018). Fertilizing Ability of Gametes at Different post-activation Times and the Sperm-Oocyte Ratio in the Artificial Reproduction of Pikeperch *Sander lucioperca*. *Aquac. Res.* 49, 1383–1388. doi:10.1111/are.13570
- Lewontin, R., Kirk, D., and Crow, J. (1968). Selective Mating, Assortative Mating, and Inbreeding: Definitions and Implications. *Eugenics Q.* 15, 141–143. doi:10.1080/19485565.1968.9987764
- Lynch, M., and Walsh, B. (1998). *Genetics and Analysis of Quantitative Traits*. Sunderland, Massachusetts, USA: Sinauer Associates.
- McQuillan, R., Leutenegger, A.-L., Abdel-Rahman, R., Franklin, C. S., Pericic, M., Barac-Lauc, L., et al. (2008). Runs of Homozygosity in European Populations. *Am. J. Hum. Genet.* 83, 359–372. doi:10.1016/j.ajhg.2008.08.007
- Nei, M., and Takahata, N. (1993). Effective Population Size, Genetic Diversity, and Coalescence Time in Subdivided Populations. *J. Mol. Evol.* 37, 240–244. doi:10.1007/BF00175500
- Nguinkal, J. A., Brunner, R. M., Verleih, M., Rebl, A., de los Ríos-Pérez, L., Schäfer, N., et al. (2019). The First Highly Contiguous Genome Assembly of Pikeperch (*Sander lucioperca*), an Emerging Aquaculture Species in Europe. *Genes* 10, 708. doi:10.3390/genes10090708
- Peripolli, E., Stafuzza, N. B., Amorim, S. T., Lemos, M. V. A., Grigoletto, L., Kluska, S., et al. (2020). Genome-wide Scan for Runs of Homozygosity in the Composite Montana Tropical Beef Cattle. *J. Anim. Breed. Genet.* 137, 155–165. doi:10.1111/jbg.12428
- Purfield, D. C., Berry, D. P., McParland, S., and Bradley, D. G. (2012). Runs of Homozygosity and Population History in Cattle. *BMC Genet.* 13, 70. doi:10.1186/1471-2156-13-70
- Santiago, E., Novo, I., Pardiñas, A. F., Saura, M., Wang, J., and Caballero, A. (2020). Recent Demographic History Inferred by High-Resolution Analysis of Linkage Disequilibrium. *Mol. Biol. Evol.* 37, 3642–3653. doi:10.1093/molbev/msaa169
- Saura, M., Caballero, A., Santiago, E., Fernández, A., Morales-González, E., Fernández, J., et al. (2021). Estimates of Recent and Historical Effective Population Size in Turbot, Seabream, Seabass and Carp Selective Breeding Programmes. *Genet. Sel. Evol.* 53, 1–8. doi:10.1186/s12711-021-00680-9
- Shi, L., Wang, L., Liu, J., Deng, T., Yan, H., Zhang, L., et al. (2020). Estimation of Inbreeding and Identification of Regions under Heavy Selection Based on Runs of Homozygosity in a Large White Pig Population. *J. Anim. Sci. Biotechnol.* 11, 1–10. doi:10.1186/s40104-020-00447-0
- Solé, M., Gori, A.-S., Faux, P., Bertrand, A., Farnir, F., Gautier, M., et al. (2017). Age-based Partitioning of Individual Genomic Inbreeding Levels in Belgian Blue Cattle. *Genet. Sel. Evol.* 49, 1–18. doi:10.1186/s12711-017-0370-x
- Thomas, M., Lecocq, T., Abregal, C., Nahon, S., Aubin, J., Jaeger, C., et al. (2020). The Effects of Polyculture on Behaviour and Production of Pikeperch in Recirculation Systems. *Aquac. Rep.* 17, 100333. doi:10.1016/j.aqrep.2020.100333
- Thompson, E. A. (2013). Identity by Descent: Variation in Meiosis, across Genomes, and in Populations. *Genetics* 194, 301–326. doi:10.1534/genetics.112.148825
- Yáñez, J. M., Newman, S., and Houston, R. D. (2015). Genomics in Aquaculture to Better Understand Species Biology and Accelerate Genetic Progress. *Front. Genet.* 6, 128. doi:10.3389/fgene.2015.00128
- Yoshida, G. M., Cáceres, P., Marín-Nahuelpi, R., Koop, B. F., and Yáñez, J. M. (2020). Estimates of Autozygosity through Runs of Homozygosity in Farmed Coho salmon. *Genes* 11, 490. doi:10.3390/genes11050490
- Zakęś, Z., Szczepkowski, M., Partyka, K., and Wunderlich, K. (2013). Effect of Gonadotropin Hormonal Stimulation on Out-Of-Season Propagation success of Different Year Classes of Indoor-Reared Pikeperch (*Sander lucioperca* (L.)). *Aquacult. Int.* 21, 801–810. doi:10.1007/s10499-012-9562-0
- Zhang, Q., Calus, M. P., Gulbrandtsen, B., Lund, M. S., and Sahana, G. (2015a). Estimation of Inbreeding Using Pedigree, 50k SNP Chip Genotypes and Full Sequence Data in Three Cattle Breeds. *BMC Genet.* 16, 88. doi:10.1186/s12863-015-0227-7
- Zhang, Q., Gulbrandtsen, B., Bosse, M., Lund, M. S., and Sahana, G. (2015b). Runs of Homozygosity and Distribution of Functional Variants in the Cattle Genome. *BMC Genomics* 16, 1–16. doi:10.1186/s12864-015-1715-x

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 De los Ríos-Pérez, Druet, Goldammer and Wittenburg. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.