

The molecular basis of zinc homeostasis in cereals

Sahand Amini¹ | Borjana Arsova²  | Marc Hanikenne¹ 

¹InBioS-PhytoSystems, Translational Plant Biology, University of Liège, Liège, Belgium

²Root Dynamics Group, IBG-2 - Plant Sciences, Institut für Bio- und Geowissenschaften (IBG), Forschungszentrum, Jülich, Germany

Correspondence

Marc Hanikenne, Department of Life Sciences, InBioS-PhytoSystems, Translational Plant Biology, University of Liège, Quartier de la Vallée, 1, Chemin de la Vallée, 4 - Bât B22, Liège B4000, Belgium.
Email: marc.hanikenne@uliege.be

Funding information

Fonds De La Recherche Scientifique - FNRS, Grant/Award Numbers: MIS F.4511.16, CDR J.0009.17, PDR T0120.18, and CDR J.0082.21; Forschungszentrum Jülich

Abstract

Plants require zinc (Zn) as an essential cofactor for diverse molecular, cellular and physiological functions. Zn is crucial for crop yield, but is one of the most limiting micronutrients in soils. Grasses like rice, wheat, maize and barley are crucial sources of food and nutrients for humans. Zn deficiency in these species therefore not only reduces annual yield but also directly results in Zn malnutrition of more than two billion people in the world. There has been good progress in understanding Zn homeostasis and Zn deficiency mechanisms in plants. However, our current knowledge of monocots, including grasses, remains insufficient. In this review, we provide a summary of our knowledge of molecular Zn homeostasis mechanisms in monocots, with a focus on important cereal crops. We additionally highlight divergences in Zn homeostasis of monocots and the dicot model *Arabidopsis thaliana*, as well as important gaps in our knowledge that need to be addressed in future research on Zn homeostasis in cereal monocots.

KEYWORDS

barley, chelation, dicot, Fe uptake strategy, monocot, rice, transport, uptake, wheat, zinc

1 | INTRODUCTION

Gramineae (Grasses), like all life forms, demand optimal zinc (Zn) supply to maintain growth and reproduction (Broadley et al., 2007). Zn is a key cofactor for enzymes involved in various cellular pathways such as gene expression, DNA replication, cell development, and protein synthesis (Gupta et al., 2016). These enzymes contribute to several physiological processes including hormone regulation, chlorophyll synthesis, lipid metabolism, carbohydrate metabolism, pollen formation and seed development (Hafeez et al., 2013; Vallee & Falchuk, 1993; J. Zhang et al., 2019). Zn requirement for plants was first discovered more than a century ago in maize (*Zea mays*, Zm) and then in barley (*Hordeum vulgare*, Hv) (Mazé, 1915; Sommer & Lipman, 1926). Zn deficiency has since been recognized as one of the most common micronutrient deficiencies in crops (Alloway, 2008; Cakmak, 2004). Zn malnutrition has similarly the largest impact among all micronutrients on human health (Black et al., 2008). There is mild to high risk of Zn deficiency on all continents, accounting for half of the world population (Gregory et al., 2017). The annual death of ~800 000 children worldwide is, for instance, attributed to acute

Zn deficiency (Akhtar, 2013). Populations that rely primarily on cereal-based diets suffer the most from this so-called hidden hunger (Rawat et al., 2013). Zn uptake from the soil is influenced by a number of characteristics including soil concentration of water-soluble Zn, soil pH, type, moisture, and mineral blend (Alloway, 2009; Broadley et al., 2007; Shahzad et al., 2014). Unfortunately, a considerable portion of Zn, that is, two-thirds in wheat (*Triticum aestivum*, Ta), one-third in rice (*Oryza sativa*, Os) and three-quarters in barley, is polished off the seeds before being consumed, while Zn retained in the endosperm is partially bound to phytate and thus poorly bioavailable to humans (Balk et al., 2019; Lombi et al., 2011; Swamy et al., 2016). Moreover, forage grasses constitute a significant share of livestock and poultry feed and are equally critical for meat-rich diets.

Grasses are the second largest and economically most important family of monocotyledons, where cereal grasses, in particular rice, wheat, maize, and barley, represent the majority of the biomass produced by agriculture (Eckardt, 2000; <http://www.fao.org/publications/sofa/2016/en/>). However, compared to the dicotyledonous model *Arabidopsis* (*Arabidopsis thaliana*, At), the molecular

basis of Zn homeostasis, unlike iron (Fe) (Kobayashi, 2019), remains poorly known in monocots. Here, we review the current knowledge of the Zn pathway from roots to the seeds in cereals (Figures 1–3, Box 1), focusing on potential differences within monocots as well as with dicots (Box 2 and 3). Whenever possible, we place characterized molecular actors along this Zn pathway (Figures 1–3). We shed light on existing gaps in our current knowledge of Zn homeostasis in staple grain crops and provide a perspective for basic research and future development of Zn biofortification strategies.

2 | Zn CHELATION IN MONOCOTS

Metal chelators such as nicotianamine (NA), free histidine, phytochelatin, glutathione, phosphate and organic acids (malate, citrate, and oxalate), increase metal solubility inside and outside the cells, buffer intracellular metal ion concentrations and control unwanted interactions with cellular components, thus contributing to metal movement and alleviating metal toxicity (Graham & Stangoulis, 2018; Olsen & Palmgren, 2014; Salt et al., 1999; Sinclair & Krämer, 2012). Among those, NA is very important for Zn homeostasis and detoxification, while citrate and histidine also play their part (Clemens, 2019). For many of the others, information is scarce. Unlike dicots, grasses produce an additional type of metal chelators: phytosiderophores (PSs, Figure 1a) (Caldelas & Weiss, 2017). Synthesized from NA (Shojima et al., 1990), mugineic acids (MAs) are a group of PSs that fulfill two roles: (i) upon secretion from roots into the rhizosphere, mobilize, chelate and stabilize Fe(III) and Zn¹ before uptake and (ii) as internal metal chelators, enable metal movement across membranes via specific transporters (Figure 1b). NA can similarly serve both roles (Clemens, 2019).

The dual role of NA, both as metal chelator and PS precursor, leads to different physiological functions for NA Synthase (NAS) in Arabidopsis and grasses (Figure 1c,d). This is reflected by differences between NAS subcellular localization in Arabidopsis and rice/maize/barley. AtNAS1 to AtNAS4 and ZmNAS1 are distributed uniformly in the cytoplasm when expressed in tobacco cells, while OsNAS2, ZmNAS2, ZmNAS3 and HvNAS1, together with two intermediate MA biosynthesis enzymes, NA aminotransferase-A (HvNAAT-A) and deoxymugineic acid (DMA) synthase 1 (HvDMAS1), localize at rough endoplasmic reticulum (rER)-derived vesicles (Mizuno et al., 2003; Nozoye et al., 2014). Similar vesicles appear in rice and barley root cells just before DMA secretion into the rhizosphere (Negishi et al., 2002; Nishizawa & Mori, 1987). It is suggested that in grasses, NA and MAs are synthesized and stored in these vesicles, before being translocated to the required locations (Figure 1b). This may prevent Fe and Zn deficiency responses triggered by metal chelation in the cytoplasm (Nozoye et al., 2014).

PSs, especially DMA, play a key role in monocots as part of a chelation-based Fe-uptake system (Strategy II). In contrast, dicots use a reduction-based Fe strategy (Strategy I), including Fe(III) to Fe(II)

BOX 1 The Zn pathway from roots to seeds in monocots

Once taken up into the root, Zn uses two different but interconnected pathways to move from root epidermal cells to the xylem (Figure 1b,d). In the symplastic pathway, Zn is transported via the cytoplasmic continuum of cells, linked by plasmodesmata, and passes through cortical and endodermal cells. This pathway is initiated at the root epidermis by plasma membrane transporters, and further influenced by plasma and vacuolar membrane transporters, as well as chelators, which altogether determine the balance between radial Zn transport to the xylem and Zn storage (Figure 1). In the apoplastic pathway, Zn is pumped extracellularly into the stellar apoplast in regions where the Casparian strip is not fully formed (Sinclair & Krämer, 2012; White, 2001). As a layer of lignin that surrounds the cells of the root endodermis (Figure 1), the Casparian strip forms an impermeable diffusion barrier dividing in two the root apoplast. Zn movement towards the stele thus requires passage to the symplast. The transport into and out of the apoplast is considered a major bottleneck in micronutrient homeostasis in plants (Olsen & Palmgren, 2014).

Once it reaches the central cylinder, Zn flux into above-ground tissues occurs via active loading of the root xylem, where Zn moves as free cation due to the acidic pH (Olsen & Palmgren, 2014). In wheat and barley, root-to-shoot translocation is a bottleneck, especially for grain Zn filling (Palmgren et al., 2008). However, this is not the case for rice as excess Zn is stored at relatively similar levels in stems and roots (Stomph et al., 2009). After being transported into the shoot xylem, Zn is taken up into living xylem parenchyma and mesophyll cells, where it is necessary for photosynthesis, and is subsequently exported into the leaf apoplast for phloem loading. In grasses, xylem-phloem exchange of Zn solutes occurs at nodes (Figure 2). Due to low solubility at the alkaline pH, Zn has to be chelated in the phloem.

Finally, Zn is provided to developing seeds via long-distance transport in the phloem (Figure 3), which has two distinct sources in grasses: (i) phloem loading after remobilization from vegetative tissues, especially flag leaves, and/or (ii) direct and continuous xylem flow from roots followed by xylem-to-phloem transfer in the stem and nodes (Waters et al., 2009). Depending on the species, this transport occurs either symplastically or from the apoplast through cellular uptake (Olsen & Palmgren, 2014; Sinclair & Krämer, 2012; Turgeon & Wolf, 2009).

¹Zn is meant for Zn(II). The oxidation state of Fe will be specified in the text, as it can occur in two valences.

BOX 2 Plant anatomy and Zn homeostasis

Zn homeostasis has distinct features in different species, stemming from anatomical differences. First, while both monocots and dicots have an apoplastic barrier in the endodermis, monocots have another layer of Casparian strip at the exodermis; the outermost layer of the cortex (Figure 1) (Olsen & Palmgren, 2014). Second, in grasses, nodes act as hubs to facilitate the distribution of Zn to young vegetative organs and reproductive tissues (Figure 2) (Yamaguchi et al., 2012), which demands intervascular Zn transfers at nodes, including from the xylem of enlarged vascular bundles (that goes toward flag leaves) to the xylem of diffuse vascular bundles (that goes toward productive tissues) (Kawakami & Bhullar, 2018). Third, distinct grain anatomy is responsible for major differences in Zn homeostasis among monocot species: at the base of rice grains, the xylem of vascular bundles is continuous and Zn can be loaded directly to the nucellar projection and the aleurone layer (Figure 3b) (Stomph et al., 2009). In contrast, xylem-to-phloem Zn transport is required at the base of seeds in wheat and barley. In wheat, several apoplastic barriers need to be crossed for Zn to reach the grain: (i) between the stem xylem and the grain base phloem (Olsen & Palmgren, 2014), (ii) between the vascular parenchyma and nucellar projections, and (iii) between maternal transfer cells and the filial endosperm (Wang et al., 1994). Active transport of free and chelated Zn ions, between symplast and apoplast, is therefore required to efflux Zn from these bottlenecks until it reaches the grain aleurone layer, endosperm, and embryo (Figure 3f) (Palmgren et al., 2008). In wheat grain, Zn is more concentrated in the embryo, while the aleurone layer contains a higher Zn concentration than the endosperm, and Fe is the highest in aleurone layer (Ajiboye et al., 2015; Balk et al., 2019). The rice seed, like the wheat grain, has an apoplastic barrier between maternal tissues and filial endosperm and embryo tissues, and Zn transporters have a critical role in these symplastic pathways (Kawakami & Bhullar, 2018). The rice endosperm also has the lowest Zn concentration in grain, and it seems that differences in nutrient transport pathways at the grain base of rice and wheat are not leading to distinct distribution of Zn in the seeds (Stomph et al., 2009).

reduction and Fe(II) uptake (Figure 1c) (Kobayashi et al., 2012; Marschner et al., 1986). The use of two strategies, ending up with the uptake of two distinct Fe forms, determines distinct interactions between Zn and Fe homeostasis (Hanikenne et al., 2021). Rice takes advantage of both strategies (Figure 3, Box 3) (Ishimaru et al., 2006).

In addition to Fe uptake, PSs have a critical role in Zn uptake from the rhizosphere in monocots (Kawakami & Bhullar, 2018). Zn deficiency

induces root PS secretion in barley (Suzuki et al., 2006), and Zn-PS complexes are taken up by maize roots (Figure 1a) (Von Wirén et al., 1996). Consistently, Zn and Fe deficiencies result in the upregulation of maize MA pathway genes (e.g., *ZmNAS1* and *ZmNAS3*) (Mallikarjuna et al., 2020). Differences in Zn uptake efficiency among rice and wheat genotypes are attributed to their PS secretion rate into the rhizosphere (Arnold et al., 2010; Cakmak et al., 1996; Neeraja et al., 2018). However, decreased secretion of DMA into the rice rhizosphere under Zn deficiency is in contrast with the elevated MA secretion observed in wheat and barley in these conditions (Ishimaru et al., 2008). Consistently, increasing DMA alone (by overexpression of *OsNAS1*) in rice does not increase Zn accumulation in the embryo and endosperm, while coupled increase in DMA and NA (by overexpressing *HvNAATb* and *OsNAS1* together) results in Zn enrichment in both tissues (Díaz-Benito et al., 2018). Fe concentration in contrast increases in seeds in both cases. Fe(III)-DMA and Zn-NA may thus be the dominant forms transported and stored in rice grains (Díaz-Benito et al., 2018).

Consistent with their role in Zn uptake, NA and DMA contribute to Zn excess tolerance in rice, particularly in intracellular and intercellular Zn mobility in roots toward pericycle cells, and therefore in root-to-shoot metal translocation. Under Zn excess, rice genes involved in NA and DMA synthesis (*OsNAS1*, *OsNAS2*, *OsNAAT1* and *OsDMAS1*), are upregulated in roots, but not in shoot (Ishimaru et al., 2008). NA and DMA as well as their transporters (Table 1) consequently play an important role in controlling the balance between Zn sequestration in root vacuoles and Zn radial transport towards the xylem, and therefore, with other players (see Zn transport Section), regulate shoot Zn accumulation (Haydon et al., 2012; Ishimaru et al., 2008; Sinclair & Krämer, 2012).

NA is also crucial for Zn long-distance transport in phloem and xylem (Figure 1a,c) (Olsen & Palmgren, 2014). Overexpression of *OsNAS1*, 2 and 3 as well as *HvNAS1* increases Zn content of polished and whole grains in transgenic lines of the respective species (Banakar et al., 2017; Johnson et al., 2011; Lee et al., 2009, 2011; Masuda et al., 2009; Singh et al., 2017; Wirth et al., 2009). In many monocotyledonous and dicotyledonous species, Zn can be transported as free Zn ions and mostly as bound to histidine and NA in the xylem sap (Caldelas & Weiss, 2017; Flis et al., 2016; White & Broadley, 2009), while citrate can also be a dominant Zn ligand in the xylem in other species (Cornu et al., 2015; Flis et al., 2016). *OsNAS3*, expressed in cells adjacent to the xylem (Inoue et al., 2003), is upregulated by Zn deficiency and downregulated by Zn excess (Ishimaru et al., 2008; Suzuki et al., 2008). *OsNAS3* transcript levels, as well as NA and DMA contents, are higher in developing panicles of high-Zn-grain rice strains than low-Zn strains (Neeraja et al., 2018).

For translocation to developing grains, NA is the dominant Zn chelator in rice phloem (Nishiyama et al., 2012). NAS and NA-metal transporter genes are highly expressed in barley seed transfer cells (Tauris et al., 2009): Zn remobilization from the phloem to transfer cells may occur as NA-Zn, whereas NA- and PS-Zn complexes would be present in the transfer cell cytosol. TaNAAT2 and TaNAS3 similarly have an active role in chelating metals during senescence in wheat, when nutrients are (re-)allocated to reproductive organs (Figure 1a,c) (Pearce et al., 2014).

BOX 3 Rice and Zn homeostasis in monocots

As it feeds a large part of the human population, rice is extensively used as a model plant for grasses (Sasaki & Burr, 2000). The number of rice (52), barley (27), wheat (16), and maize (8) genes cited in this manuscript to be linked with Zn homeostasis/tolerance (Table 1), reflects that a substantial amount of studies on the molecular basis of Zn homeostasis in cereals has been dedicated to rice. However, rice is not a perfect model for wheat and barley, especially for uncovering metal homeostasis mechanisms. It was indeed suggested that bottlenecks for Zn transport from the rhizosphere to grain are different in rice and other cereals (Stomph et al., 2009).

First, rice is primarily cultivated in paddy fields, in conditions that prevent Fe oxidation and precipitation (Grillet & Schmidt, 2019), and in contrast to other cereals, it uses both Fe uptake strategies (Figure 3a) (Ishimaru et al., 2006). It can, therefore, be speculated that a number of metal homeostasis genes play dissimilar roles in rice and other grasses. Differential DMA-secretion patterns in rice compared to wheat and barley in response to Zn deficiency (Suzuki et al., 2008), may also reflect the presence of a dual Fe uptake strategy in rice.

Second, rice has different Zn allocation strategies to grains. Unlike wheat, rice leaves are not a Zn source for grains and rice plants dominantly use Zn uptake from soil and xylem transport postflowering to provide Zn to grains. In general, differential mechanisms of Zn translocation to rice grains under sufficiency and deficiency are reported (Sperotto, 2013; Wu et al., 2010). For instance, when Zn is not supplied sufficiently, stored Zn in roots, stem, and sheath, but not leaves, is reported to remobilize to rice grains (Jiang et al., 2007; Sperotto et al., 2009; Stomph et al., 2009). In contrast, in field conditions, pre-anthesis remobilization of deposited Zn from vegetative tissues via phloem transport has been reported as the main Zn source for grains in wheat, while maize grains accumulate Zn post-silking from the soil (Xue et al., 2019). The contribution of Zn sources, soil, or remobilization from other tissues, in grain Zn accumulation, can also depend on factors like water availability, timing of senescence, length of grain-filling period, and nitrogen status (Cakmak & Kutman, 2018).

Third, Zn root-to-shoot transport is not (or is less of) a bottleneck in rice compared to other cereals. The continuous Zn uptake from soil in rice might also be explained by a highly effective root-to-shoot Zn transport system, compared to wheat and barley where Zn excess is stored in root vacuoles (Stomph et al., 2009). OSHMA2 may be the key for this feature in rice. The control of HMAs on Zn xylem loading is indeed crucial for Zn root-to-shoot partitioning (Claus et al., 2013; Hanikenne et al., 2008). However, differences in Zn radial transport in rice roots remain to be investigated.

Finally, rice has a different grain anatomy than wheat, barley, and maize, which modify the Zn pathway to the grain (Figure 3b,f, see Box 2 for details) (Hands and Drea, 2012). The genes responsible for this transport in wheat and barley are not identified yet.

In addition to organic metal chelators, a class of proteins, metallothioneins (MT) plays a role in metal chelation (Sinclair & Krämer, 2012). HvMT3 and HvMT4 can bind Zn, and HvMT4 may contribute to Zn storage in barley grains (Hegelund et al., 2012). HvMT2b2 is strongly induced in the embryo and the aleurone of barley grains upon foliar Zn application and may be a chelator under excess Zn. HvMT2a was also proposed as a key Zn chelator in the endosperm (Tauris et al., 2009). In rice, *OsMT1a* is induced by Zn and its overexpression increases Zn concentration in both vegetative tissues and grains (Z. Yang, Wu, et al., 2009), whereas *OsMTI-1b* was shown to bind Zn (Malekzadeh et al., 2020).

3 | Zn TRANSPORT IN MONOCOTS

Zn cellular import/export and intracellular trafficking in plants require a passage through numerous biological membranes (Figures 1–3, Box 1), ensured by many proteins active transport of Zn and other divalent cations. The function of several of these proteins in Zn homeostasis is tightly linked to Fe homeostasis and contributes to interactions between these two nutrients (Hanikenne et al., 2021).

3.1 | ZIP family

Zn-regulated transporter (ZRT), iron-regulated transporter (IRT)-like protein (ZIP) transporters are generally mediating cellular uptake of Zn and other divalent metals like manganese (Mn), Fe or even cadmium (Cd), with a transport direction from outside the cell or from an organelle to the cytoplasm (Guerinot, 2000). Several rice ZIPs are linked to Zn movement throughout the plant toward grains. For instance, *OsZIP3* and *OsZIP1* are among rice marker genes for Zn status and show a good association between gene expression level and grain Zn concentration (Swamy et al., 2016). *OsZIP1* is expressed in the root exodermis and endodermis (Figure 1b) and is upregulated in roots and shoot under Zn deficiency (Ramesh et al., 2003). Higher *OsZIP1* expression is associated with higher tolerance to Zn deficiency among cultivars (Lu et al., 2021). *OsZIP1*, and *ZmZIP5*, display a dual localization in the ER and plasma membrane (PM) (Figure 1b) (Li et al., 2013, 2019; Liu et al., 2019; Mallikarjuna et al., 2020). Interestingly, a role of the ER in facilitating root-to-shoot Zn movement has been proposed in Arabidopsis (Figure 1d) (Sinclair et al., 2018). *OsZIP1* and *ZmZIP5* may therefore not only detoxify excess Zn out of cytosol but also contribute to Zn root-to-shoot translocation.




FIGURE 1 Schematic of root and molecules involved in Zn homeostasis in monocots and dicots. Whole-plant and root layers of monocots (a and b) and the dicot *Arabidopsis thaliana* (c and d) are shown with the most important Zn transporters and chelators that play a role in Zn uptake from rhizosphere into roots and Zn movement to the root stele. Differences in Fe-uptake strategies between monocots and dicots are illustrated as well as the additional cell layer in monocots, the exodermis. Black boxes in the exo- and endodermis represent apoplastic barriers. At, *Arabidopsis thaliana*; DMA, deoxymugineic acid; ER, endoplasmic reticulum; FRO, ferric reductase oxidase; HMA, heavy metal ATPases; Hv, *Hordeum vulgare*; MTP, metal transport protein; NA, nicotianamine; NAS, nicotianamine synthase; Os, *Oryza sativa*; PS, phytosiderophore; rER, rough ER; YS, yellow stripe; YSL, yellow stripe-like; ZIP, Zn-regulated transporter (ZRT); iron-regulated transporter (IRT)-like protein; Zm, *Zea mays*

OsZIP8 is highly induced in roots and shoot by Zn deficiency and complements a Zn-deficient yeast strain (Lee, Kim, et al., 2010; X. Yang, Huang, et al., 2009). The OsZIP8 protein is localized at the PM in root cells. OsZIP8 overexpression in rice results in root Zn accumulation, while shoot and seeds are deficient (Lee, Kim, et al., 2010). Similar results were reported for OsZIP5 (Lee, Jeong, et al., 2010). OsZIP5 and OsZIP9 are PM localized and involved in Zn and Cd influx into root cells, with the corresponding genes being induced by Zn deficiency (L. Tan et al., 2020). *oszip5*, *oszip9* and *oszip5oszip9* mutants accumulate lower Zn concentrations and display growth retardation under Zn deficiency. The *oszip9* and *oszip5oszip9* mutants have the most severe phenotypes (L. Tan et al., 2020). OsZIP9 is expressed in root epidermal and exodermal cells. Its root expression level is highly associated with Zn content of milled grains (Swamy et al., 2016; Yang et al., 2020). In contrast to OsZIP5, OsZIP9 is controlled by shoot Zn status (L. Tan et al., 2020), suggesting a long-distance systemic regulation as reported for AtMTP2 and AtHMA2 in Arabidopsis (Sinclair et al., 2018). OsZIP8, OsZIP5 and OsZIP9 are therefore involved in Zn uptake into root cells (Figure 1b).

OsZIP3 is expressed in the xylem transfer cells in enlarged vascular bundles and transports Zn (Figure 2) (Sasaki et al., 2015). The OsZIP3 gene has higher expression in flag leaves than roots and is slightly induced by Zn deficiency in shoot (Ramesh et al., 2003).

The OsZIP4 gene functions in root-to-shoot Zn translocation (Bashir, Takahashi, Nakanishi, et al., 2013). It is regulated by Zn availability (Suzuki et al., 2012). OsZIP4 is localized at the PM (Ishimaru et al., 2005, 2008) and is likely involved in transporting Zn from the enlarged to the diffuse vascular bundle in the basal node and to panicles (Box 1, Figure 2) (Mu et al., 2021). Consistently, the expression level of OsZIP4 in flag leaves correlates with rice grain Zn content (Swamy et al., 2016).

OsZIP7 rescues the Zn uptake-defective *zrt1zrt2* yeast mutant under Zn deficiency, but not under Fe deficiency (Ricachenevsky et al., 2018; L. Tan et al., 2019). OsZIP7 localizes at the PM in rice protoplasts (L. Tan et al., 2019). The OsZIP7 gene is highly expressed in rice roots and less in flag leaves (Chandel et al., 2010), while it is upregulated by Zn deficiency in root stele and parenchyma cells of

TABLE 1 List and function of genes linked with Zn homeostasis/tolerance in monocots

	Rice	Barley	Wheat	Maize
ZIP	Chelator synthesis OsNAS1	NA/MA synthesis, Zn enrichment in grain HvNAS1	NA/MA synthesis, Zn enrichment in grain TaNAS3	NA/MA synthesis ZmNAS1
	OsNAS2	NA/MA synthesis, Zn enrichment in grain HvNAAT-A	NA/MA synthesis TaNAAT2	NA/MA synthesis ZmNAS2
	OsNAS3	NA/MA synthesis, Zn enrichment in grain HvNAAT-b	NA/MA synthesis ZMNAS3	NA/MA synthesis ZMNAS3
	OsNAAT1	NA/MA synthesis HvDMAS1	NA/MA synthesis HvDMAS1	
	OsDMAS1	NA/MA synthesis HvMT2a	Zn chelator in grain endosperm HvMT2a	
	OsMT1a	Zn enrichment HvMT2b2	Zn chelator in grain under excess Zn HvMT2b2	
	OsMT1-b	Zn/Cd chelator HvMT3	- HvMT3	
HMA	OsZIP1	Root Zn uptake, Zn transport into endodermis/out of exodermis, Zn detoxification into ER HvZIP2	TaZIP3	ZmZIP5
	OsZIP3	Zn translocation in nodes HvZIP3	TaZIP6	ZmZIP7
	OsZIP4	Zn translocation in nodes HvZIP5	TaZIP7	ZmIRT1
	OsZIP5	Root Zn/Cd uptake HvZIP6	TaZIP9	
	OsZIP6	- HvZIP7	TaZIP10-like1	
	OsZIP7	Zn/Cd xylem loading in root, Zn/Cd translocation in nodes HvZIP8	TaZIP13	
	OsZIP8	Root Zn uptake HvZIP10	TaZIP13-like2	
	OsZIP9	Root Zn/Cd/Co uptake HvZIP13		
	OsZIP10	- HvIRT1		
	OsIRT1	Fe/Zn uptake		
HMA	OsHMA2	Root-to-shoot Zn translocation, xylem-to-phloem transport of Zn at nodes, Zn translocation to the seed endosperm HvHMA1	TaHMA2	Zn root-to-shoot translocation at nodes
	OsHMA3		TaHMA2-like	

TABLE 1 (Continued)

	Rice	Barley	Wheat	Maize
		Zn/Cd sequestration into root vacuoles, Zn detoxification in root	Root-to-shoot Zn translocation, Zn translocation to the seed endosperm	
	OshMA9	Zn/Cu efflux from root cells		
MTP	OsMTP1 (OsOZT1)	Vacuolar Zn sequestration	HvMTP1	Vacuolar Zn sequestration in seeds
	OsMTP5-like	Involved in Zn tolerance	HvMTPc4	Vacuolar Zn sequestration in seeds
	OsMTP6	-	HvMTP5	-
	OsMTP7	-		
	OsMTP7-like	Involved in Zn tolerance		
	OsMTP8.1	-		
	OsMTP9	-		
	OsMTP11	-		
	OsMTP12	-		
MFS (ZIF/ZIFL)	OsZIFL2	-	HvZIF1	Zn sequestration in grains
	OsZIFL4 (OsTOM1)	DMA secretion from root		
	OsZIFL5 (OsTOM2)			
	OsZIFL7 (OsTOM3)			
	OsZIFL10	-		
	OsZIFL12 (OsVMT)	DMA sequestration into vacuole		
	OsENA1	Transports NA out of cells into the apoplasm		
YSL	OsYSL2	-	HvYSL1	Transport DMA-Zn/Fe/Mn complexes
	OsYSL6	-	HvYSL2	Transport DMA-Zn/Fe/Mn complexes in the
			ZmYSL1	transport PS-Zn complex

(Continues)

TABLE 1 (Continued)

	Rice	Barley	Wheat	Maize
	OsYSL13-like OsYSL15		endodermis, xylem DMA-metal loading	
VIT	- DMA transport			
	OsVIT1 OsVIT2 OsVIT5	Zn/Fe sequestration in vacuoles of flag leaves Zn/Fe sequestration in vacuoles of flag leaves grain Zn accumulation	TdVIT1 TdVIT2	- -
NRAMP	OsNRAMP1 OsNRAMP5	- -		ZmNRAMP1 -
Transcription Factors	OsZIP48 OsZIP50 OsNAC5 OsIDEF1 OsIRO2	Regulation of Zn homeostasis Regulation of Zn homeostasis Regulation of Zn/Fe remobilization to developing grains regulation of ZIFL genes	HvbZIP56 HvbZIP62 TaNAM-B1 (TaGPC1)	Regulation of Zn homeostasis Regulation of Zn homeostasis Regulation of Zn/Fe remobilization to developing grains
Other genes	OsFRDL1 OsPHO1;1	Zn/Fe homeostasis, citrate transport into the xylem Zn/Fe/P cross-homeostasis		

Note: The gene inventory is limited to genes discussed in the document. The genes are categorized by gene family and species. Genes may have other functions that are not Zn-related and not noted in the table. Dash (-), no reported/proposed function. Blank cells, no available information. Genes on each row are not orthologs.

Abbreviations: bZIP, basic leucine zipper; DMA, deoxymugineic acid; DMAS, deoxymugineic acid synthase; ENA, efflux transporter of NA; ER, endoplasmic reticulum; FRDL, ferric reductase defective-like; GPC, grain protein content; HMA, heavy metal ATPases; Hv, *Hordeum vulgare*; IDEF, iron deficiency-responsive element-binding factor; IRO, iron-related transcription factor; IRT, iron-regulated transporter; MA, mugineic acid; MFS, major facilitator superfamily; MT, metallothioneins; MTP, metal transport protein; NA, nicotianamine; NAAT, NA aminotransferase; NAC, NAM; ATAF and CUC; NAM, no apical meristem; NAS, nicotianamine synthase; NRAMP, natural resistance-associated macrophage protein; Os, *Oryza sativa*; OZT, *Oryza sativa* Zn transporter; PHO, phosphate; PS, phytosiderophore; Ta, *Triticum aestivum*; TOM, transporter of mugineic acid; VIT, vacuolar iron transporter; VMT, vacuolar mugineic acid transporter; YSL, yellow stripe; YSL, yellow stripe-like; ZIF, Zn-induced facilitator; ZIFL, Zn-induced facilitator-like; ZIP, Zn-regulated transporter (ZRT) iron-regulated transporter (IRT)-like protein; Zm, *Zea mays*.

enlarged vascular bundle in the basal node (Figures 1b and 2) (L. Tan et al., 2019). An *oszip7* mutant displays Zn accumulation in roots and basal nodes and reduced Zn delivery to upper plant parts (Gindri et al., 2020; L. Tan et al., 2019). Overexpression of *OsZIP7* in Arabidopsis significantly increases leaf, but decreases root, Zn concentration (Ricachenevsky et al., 2018). Moreover, *OsZIP7*-overexpressing Arabidopsis lines are hypersensitive to high Zn and display significantly increased Zn seed concentrations (Ricachenevsky et al., 2018). Accordingly, *OsZIP7* plays an important role in Zn xylem loading in roots, inter-vascular transport in the basal node, and consequently Zn translocation toward leaves and rice grains (L. Tan et al., 2019). The barley ortholog of *OsZIP7*, *HvZIP7*, is similarly localized at the PM and was implicated in root-to-shoot Zn translocation (Figure 1) (Tiong et al., 2014). *TaZIP7*, another orthologous gene in wheat, is highly expressed during senescence in leaves (Pearce et al., 2014), suggesting a role in Zn translocation toward the phloem and reproductive organs. Finally, and similar to *OsZIP7*, *ZmZIP7* increases Zn concentration in seeds when expressed in Arabidopsis (Li et al., 2016). Unlike rice/wheat, *ZmZIP7* is more expressed in flag leaves than in baby kernels, and is among low-expressed ZIPs in maize (Mondal et al., 2014). Their Arabidopsis homologs, *AtIRT3* and *AtZIP4* (Evens et al., 2017; Tiong et al., 2015), are respectively, suggested to function in Zn xylem unloading (Lin et al., 2009), and Zn uptake into root endodermis and pericycle cells to mediate Zn xylem loading (Lin et al., 2016). This subgroup of plant ZIPs, including *AtIRT3*, *AtZIP4* as well as the rice, barley, wheat and maize ZIP7 proteins (Evens et al., 2017), thus appear to share a role in mediating root-to-shoot Zn translocation.

Other ZIPs were shown to be Zn-responsive in monocots, but their contribution to Zn uptake and distribution remains unclear. For instance, *OsZIP6* is expressed in roots and shoot, and is highly upregulated in shoot under Zn, Fe and Mn deficiency (Kavitha et al., 2015). Its orthologs *HvZIP6* and *TaZIP6* are only induced in roots by Zn deficiency (Evens et al., 2017; Tiong et al., 2015). In this respect, the monocotyledonous ZIP6 genes differ from their Arabidopsis ortholog, *AtZIP6*, which is unresponsive to the Zn status (Talke et al., 2006; Spielmann et al., 2020). The *HvZIP2*, *HvZIP3*, *HvZIP5*, *HvZIP8*, *HvZIP10* and *HvZIP13* genes are upregulated by Zn starvation, and *HvZIP3*, *HvZIP5* and *HvZIP8* transport Zn in yeast (Nazri et al., 2017; Pedas et al., 2009; Tiong et al., 2015). Reports on wheat ZIP genes are rare and with a few exceptions (see *TaZIP6* and *TaZIP7* above), their role remains to be studied. However, Zn transport activity in yeast was shown for *TaZIP3*, *TaZIP6*, *TaZIP7*, *TaZIP9* and *TaZIP13* proteins (Evens et al., 2017).

Finally, in rice, *OsIRT1* and *OsIRT2* are upregulated under Fe deficiency and the corresponding proteins are known as Fe transporters, contributing to the dual Fe uptake strategy in rice (Che et al., 2019; Ishimaru et al., 2006). Their transcript levels exhibit good association with grain Zn content (Swamy et al., 2016). Rice plants overexpressing *OsIRT1* accumulate higher Fe and Zn in roots, shoot and seeds, and are sensitive to excess Zn (Lee & An, 2009), which suggest that in addition to Fe, *OsIRT1* can also transport Zn (Figure 1b). *ZmIRT1* complements yeast strains

defective in Zn or Fe uptake under low Zn or Fe supply, respectively, and does so more efficiently than any other ZmZIP transporter (*ZmZIP1–8*) and *OsIRT1* (S. Li et al., 2013). Overexpression of *ZmIRT1* increases Zn and Fe content in Arabidopsis roots and shoot (Li et al., 2015). Several barley IRTs also rescue a yeast strain defective in Zn transport (Pedas et al., 2008, 2009). Although it is involved in Mn uptake in barley (Pedas et al., 2008), expression of *HvIRT1* is induced by excess Zn (Kaznina et al., 2019). *AtIRT1–3* are the best-characterized ZIPs in Arabidopsis. They are mainly known for Fe homeostasis but they have broad substrate range and were reported to transport Zn as well (Korshunova et al., 1999). *AtIRT1* is key for Strategy I Fe(II) uptake in Arabidopsis (Kobayashi et al., 2012), and it takes up Zn, Cd, Mn, Co and nickel (Ni) from soil (Figure 1d) (Fukada et al., 2011; Rogers et al., 2000; Vert et al., 2002). Its transcript level is upregulated by Fe deficiency and excess Zn (Becher et al., 2004; Connolly et al., 2002). In dicots, IRT1-like proteins play an important role in the interactions between Zn and Fe homeostasis (Hanikenne et al., 2021). In contrast to its Arabidopsis homolog, the *OsIRT1* transcript level is not induced by Zn excess (Ishimaru et al., 2008), indicating a distinct interaction between Zn and Fe in the two species.




FIGURE 2 Schematic of a rice node and molecules involved in the distribution of Zn to the leaf and the panicle. The Zn pathway from the root vasculature to the node where Zn is to be distributed between leaf and panicle is illustrated. Black boxes in Bundle sheath represent the apoplastic barrier. DMA, deoxymugineic acid; HMA, heavy metal ATPases; Os, *Oryza sativa*; VMT, vacuolar mugineic acid transporter (also called *OsZIFL12*); ZIP, Zn-regulated transporter (ZRT), iron-regulated transporter (IRT)-like protein

3.2 | HMA family

A subset of heavy metal ATPases (HMAs) of the P_{1B}-type ATPases has a Zn transport function (Hanikenne & Baurain, 2014; Williams & Mills, 2005). These pumps are unique in plants among eukaryotes for Zn cellular export (Olsen et al., 2016). Hence, *OsHMA9* is expressed in root vascular bundles and anthers and is induced by excess Zn, Cu and Cd (Lee et al., 2007). *oshma9* mutants are hypersensitive to excess Zn and Cu. Localized at the PM, *OsHMA9* is probably involved in Zn and Cu efflux from root cells (Lee et al., 2007). A similar function was recently proposed for *OsZIP1* under elevated metal levels (Liu et al., 2019).

OsHMA3 was identified as a major determinant of Cd accumulation variation in leaves and grains among rice cultivars (Miyadate et al., 2011; Ueno et al., 2010). Its homolog, *AtHMA3*, has a similar function in Arabidopsis (Chao et al., 2012; Morel et al., 2009). Mostly expressed in roots, HMA3 proteins mediate Cd and Zn sequestration into vacuoles (Figure 1b) (Cai et al., 2019; Morel et al., 2009; Ueno et al., 2010). In addition to Cd, *OsHMA3* may be involved in Zn homeostasis: it transports Zn when expressed in Arabidopsis (Miyadate et al., 2011), and *OsHMA3* overexpression in rice leads to an enhanced Zn concentration, as well as induction of the *OsZIP4*, *OsZIP5*, *OsZIP8*, *OsZIP9* and *OsZIP10* genes, in roots (Sasaki et al., 2014). As a higher Zn sequestration in root vacuoles by *OsHMA3* results in cytosolic Zn depletion, this *ZIP* upregulation corresponds to a Zn-deficiency response, which enables unchanged Zn concentration in aerial parts of the overexpressor lines (Sasaki et al., 2014). Conversely, *OsHMA3* loss of function results in enhanced sensitivity to excess Zn, and reduced Zn accumulation in root and shoot vacuoles. Triggered by the higher Zn availability in root cell cytosol, *OsZIP4*, *OsZIP5*, *OsZIP8* and *OsZIP10* in roots, as well as *OsZIP5* and *OsZIP8* in shoot, are downregulated in the *oshma3* mutant, which may be responsible for the lower Zn concentration in shoot (Cai et al., 2019). Altogether, *OsHMA3* is involved in Zn detoxification in rice roots, and in Zn sequestration into root vacuole as a reserve for Zn-deficient conditions (Cai et al., 2019).

The *AtHMA2* and *AtHMA4* pumps play a key role in Zn root-to-shoot translocation (Figure 1d) (Hussain et al., 2004). It is interesting to note that monocots, in most cases, have only one such pump whereas this function is duplicated in Brassicaceae with *AtHMA2/4*-like proteins (Baxter et al., 2003; Hanikenne & Baurain, 2014). As its Arabidopsis homologs which are involved in active Zn export from the symplast to the xylem by pumping it out of adjacent cells in the roots (Hussain et al., 2004), *OsHMA2* is expressed in the root vasculature and is responsible for root-to-shoot Zn translocation (Figure 1b) (Satoh-Nagasawa et al., 2012). Rice *hma2* mutants accordingly display decreased root-to-shoot Zn transfer (Satoh-Nagasawa et al., 2012). *OsHMA2* is also involved in xylem-to-phloem transport of Zn at nodes: it is expressed in the phloem of enlarged and diffuse vascular bundles (Figure 2) (Takahashi et al., 2012; Yamaji et al., 2013). A wheat homolog of *OsHMA2*, *TaHMA2*, is also localized at the PM. The *TaHMA2* gene is highly expressed in nodes, and its overexpression increases Zn root-to-shoot translocation in wheat, rice and tobacco (J. Tan et al., 2013). *TaHMA2* and its paralog, *TaHMA2-like*, are significantly upregulated during senescence

(Pearce et al., 2014), suggesting a role in source-to-sink remobilization of Zn and/or other metals, which coincides with senescence (Figure 3e). Moreover, expression of *HvHMA2* complements the shoot Zn-deficiency phenotype of the *hma2hma4* Arabidopsis double-mutant (Mills et al., 2012).

In addition to a defect in Zn root-to-shoot translocation, the Arabidopsis *hma2hma4* double mutant fails to direct Zn toward the grain endosperm and embryo, and Zn accumulates in maternal seed coat, where *AtHMA4* was proposed to play a central role in mediating postphloem translocation of Zn to the seed endosperm (Olsen et al., 2016). A similar role may be fulfilled by *OsHMA2* (Østerberg & Palmgren, 2018) and *HvHMA2*, which are expressed in the transfer cells of developing seeds (Figure 3d,h) (Olsen & Palmgren, 2014; Tauris et al., 2009). In contrast, *TaHMA2*-overexpressing wheat lines show decreased Zn concentration in the endosperm combined to increase Zn in the embryo and the aleurone layer (J. Tan et al., 2013). This discrepancy might result from ectopic overexpression of *TaHMA2* with a ubiquitin promoter in the transgenic lines. Alternatively, *TaHMA2* may play a different role than its homologs in Arabidopsis and rice. It is also possible that Zn transport in rice grains is different from wheat and barley due to anatomical differences (Box 2).

HvHMA1 was reported to mediate export of Zn and other metals, especially copper (Cu), out of chloroplasts in barley (Figure 1) (Mikkelsen et al., 2012). Its homolog in Arabidopsis, *AtHMA1*, has been described both as a Zn exporter out of chloroplast (Y. Y. Kim et al., 2009) and a Cu importer into the chloroplast (Seigneurin-Berny et al., 2006). *HvHMA1* expression is induced in leaves by Zn deficiency and suppressed by excess Zn. *HvHMA1*, besides leaf plastids, localizes to intracellular compartments of grain aleurone cells, and was suggested to be involved in Zn and Cu homeostasis during grain filling (Mikkelsen et al., 2012).

3.3 | MTP family

Cation diffusion facilitators (CDF), called metal tolerance/transport protein (MTP) in plants, are involved in metal transport out the cytoplasm. A MTP subfamily mediates Zn vacuolar sequestration (Montanini et al., 2007). *Oryza sativa Zn Transporter 1 (OZT1)*, also called *OsMTP1*, is induced by Zn excess in roots and shoot. *OsMTP1* localizes to the tonoplast *in planta* and confers Zn resistance when expressed in yeast (Lan et al., 2013; Menguer et al., 2013). *OsMTP1* RNAi lines are sensitive to Zn (Yuan et al., 2012). Similar to its Arabidopsis homologs, *AtMTP1* and *AtMTP3*, *OsMTP1* has a role in vacuolar Zn sequestration (Figure 1b). The *AtMTP1*, *AtMTP2* and *AtMTP3* proteins are respectively involved in Zn vacuolar sequestration (Desbrosses-Fonrouge et al., 2005), in Zn transport into the ER to facilitate cell-to-cell Zn movement in roots, consequently facilitating root-to-shoot translocation (Sinclair et al., 2018) (Figure 1d) and in vacuolar storage enabling Zn removal from the root-to-shoot pathway under excess Zn (Arrivault et al., 2006). The *AtMTP2* function is possibly conserved in monocots via *OsZIP1* and *ZmZIP5* for instance (see above). *HvMTP1*, which transports Zn in yeast,




FIGURE 3 Schematic of seed and molecules involved in Zn homeostasis in monocot grains. Whole-plant and seed anatomy of rice (a–d) and other grasses like wheat and barley (e–h) are shown with the most important players Zn homeostasis in the seed. Differences in the anatomy of rice and barley seeds are illustrated at the base of grains. Different growth environments between the two species and their impact of Fe/Zn uptake are also shown. For each rice and other grasses, the middle/top figure shows whole seeds (b, f), transverse section (c, g) as indicated at the top, and a zoom on the indicated boxes on the transverse section (d, h). HMA, heavy Metal ATPases; Hv, *Hordeum vulgare*; MTP, metal transport protein; NA, nicotianamine; Os, *Oryza sativa*; PS, phytosiderophore; Ta, *Triticum aestivum*; TOM, transporter of mugineic acid (also called ZIFL); YSL, yellow stripe-like; ZIF, Zn-induced facilitator

localizes to the tonoplast and is expressed in roots, shoot and grains, especially in the aleurone layer (Podar et al., 2012; Tauris et al., 2009). In developing seeds, HvMTP1 may sequester Zn in vacuoles and thus control the amount of Zn transported across postphloem apoplastic barriers (Figure 3g) (Olsen & Palmgren, 2014). A similar role was proposed for HvMTPc4 in barley grains (Figure 3g). Overexpression of HvMTP1 under an endosperm-specific promoter in barley increases Zn, Fe, Cu and Mn concentrations in grains under control growth conditions (Menguer et al., 2018). Moreover, HvMTP5 transcripts are significantly less abundant in barley lines naturally accumulating high Zn in seeds compared to low-Zn lines. It might thus sequester Zn in maternal tissues and prevent Zn loading to filial tissues (Detterbeck et al., 2019).

In rice, Zn starvation induces the expression of multiple MTP genes: *OsMTP6*, *OsMTP7*, *OsMTP8.1* and *OsMTP12* in roots, and *OsMTP9* and *OsMTP12* in shoot (Ram et al., 2019). *OsMTP1*, *OsMTP5-like* and *OsMTP7-like* are in addition upregulated in rice cultivars with higher tolerance to Zn deficiency (Lu et al., 2021). Although their corresponding genes are regulated by Zn, *OsMTP9* and *OsMTP8.1* are involved in Mn homeostasis. *OsMTP9* localizes at the PM of root cells and has a critical role in Mn translocation out of the exodermis and endodermis to the root stele and thus to shoot (Ueno

et al., 2015). *OsMTP8.1* is reported to be an Mn-specific transporter involved in Mn tolerance in rice shoot (Chen et al., 2013). A similar Mn-specific role was previously reported for *AtMTP11* in Arabidopsis (Delhaize et al., 2007; Peiter et al., 2007). These rice MTP proteins in fact belong to distinct CDF subfamilies, defined by phylogenetic analyses, with different substrate specificities: *OsMTP9* and *OsMTP8.1* belong to an Mn sub-family, *OsMTP6* and *OsMTP7* to a Zn/Fe sub-family and *OsMTP12* to a Zn subfamily (Ram et al., 2019). Taken together, these Zn-induced *OsMTP* genes may act directly or indirectly (i.e., by contributing to the homeostasis of other metals) in Zn homeostasis in rice.

3.4 | ZIF family

The Zn-Induced facilitator 1 (ZIF1)-like (ZIFL) protein family, belonging to major facilitator superfamily (MFS) of transporters, is involved in metal homeostasis, especially Zn in Arabidopsis (Sinclair & Krämer, 2012). *AtZIF1* localizes to the tonoplast and transports NA into the vacuole. *AtZIF1* expression is increased by high Zn or Fe deficiency. An *atzif1* mutant is hypersensitive to Zn, while *AtZIF1* overexpression enhances vacuolar Zn and NA accumulation and leads

to Zn- and Fe-deficiency symptoms (Haydon & Cobbett, 2007; Haydon et al., 2012). Based on sequence similarities to Arabidopsis ZIFL proteins, 10–15 members of this family are identified in rice, maize and wheat (Ricachenevsky et al., 2011; Sharma et al., 2019). The higher number of monocot ZIFL genes compared to dicots may result from early segmental and tandem genomic duplications in the monocot lineage (Tang et al., 2010; Ricachenevsky et al., 2011). Six rice ZIFLs, *OsZIFL2*, *OsZIFL4* (or Transporter Of Mugineic acid 1, *OsTOM1*), *OsZIFL5* (or *OsTOM2*), *OsZIFL7* (or *OsTOM3*), *OsZIFL10* and *OsZIFL12*, are significantly upregulated in roots upon Zn excess. Fe deficiency also induces *OsTOM1* to -3 and *OsZIFL12* expression in roots, but not in shoot (Che et al., 2019; Ricachenevsky et al., 2011). *OsTOM1* is involved in root secretion of DMA and Fe(III) stabilization in the rhizosphere (Figure 3a) (Nozoye et al., 2011). Compared to WT plants, *ostom2* RNAi rice lines accumulate slightly higher Zn concentration in leaves and roots (Nozoye et al., 2015). Eight out of 15 wheat ZIFL genes are similarly induced by excess Zn, five of which are also induced by Fe-deficient conditions (Sharma et al., 2019). The PM Efflux transporter of NA 1 (*OsENA1*), another MFS/TOM protein, transports NA out of cells into the apoplast (Nozoye et al., 2019), where it may be important for Fe mobilization. As Zn supplementation complements the shorter and hairy root phenotypes of *OsENA1*-overexpressing rice plants, *OsENA1*, on top of Fe, may also contribute to Zn homeostasis (Nozoye et al., 2019).

OsZIFL12, also called *OsVMT* (vacuolar mugineic acid transporter), is induced by Fe deficiency in roots. It is highly expressed in roots and in parenchyma cell bridges of node I (Che et al., 2019), which are part of the nutrient pathway between the enlarged and diffuse vascular bundle xylem (Yamaji et al., 2015). The corresponding protein localizes at the tonoplast and is proposed to sequester DMA into vacuoles, enabling the formation of Fe(III)-DMA complexes in this compartment (Figure 2) (Che et al., 2019). The Zn concentration is higher in node I of brown rice and polished grains in *vmt* mutant lines compared to WT plants. A higher DMA concentration is also observed in polished grains of *vmt* mutants, and, as DMA can chelate Zn, this was proposed to be responsible for the higher Zn concentration in nodes and grains (Che et al., 2019).

Finally, using laser capture microdissection, different parts of barley grains, grown in soil under control condition and upon foliar Zn application, were dissected and transcription of potential Zn homeostasis-related genes was analyzed (Tauris et al., 2009). *HvZIF1* was the only Zn homeostasis-related transporter gene upregulated by foliar Zn application in the embryo (Tauris et al., 2009), and may contribute to Zn sequestration in barley grains (Figure 3f).

3.5 | Yellow stripe-like (YSL) family

YSL transporters are members of the oligopeptide transporter (OPT) superfamily, and are known to transport PS/NA-metal complexes into the cytosol by proton-coupled symport (Sinclair & Krämer, 2012). *ZmYS1* was the first identified YSL and, as part of Strategy II, has a central role in PS-Fe(III) uptake into the maize root

epidermis (Figure 1b) (Curie et al., 2001). In addition to this crucial role in Fe nutrition, it was later shown to transport DMA complexes with several metals including Zn (Schaaf et al., 2004). Furthermore, *YS1* mutation in maize significantly reduces Zn uptake and the root-to-shoot translocation rate of PS-Zn complexes (Von Wirén et al., 1996).

HvYSL2, and to a lower extent *HvYS1*, are transporting DMA-Zn/Fe/Mn complexes (Araki et al., 2011). The corresponding genes, especially *HvYSL2*, are highly induced by Zn and/or Fe deficiency in roots of barley seedlings: *HvYSL2* is more induced by Zn than Fe deficiency, and vice-versa for *HvYS1*, and combined Fe and Zn deficiencies highly induce both. These genes are not regulated significantly by Zn and/or Fe starvation in shoot. *HvYSL2* appears to be involved in PS-metal transport in the endodermis and in xylem metal loading (Figure 1b). *HvYS1* rather plays a role in Fe uptake from the rhizosphere (Araki et al., 2011). YSL transporters were suggested as potential candidates for a role in Zn accumulation in barley grains (Detterbeck et al., 2019).

Rice YSLs are also linked to Zn homeostasis. Upon Zn excess, *OsYSL2* and *OsYSL15* are upregulated in roots (Ishimaru et al., 2008). *OsYSL15* is upregulated in shoot under Fe deficiency, as well (Anuradha et al., 2012; Che et al., 2019). Moreover, *OsYSL15*, *OsYSL13-like* and *OsYSL6* are more highly expressed in rice cultivars with higher tolerance to Zn deficiency (Lu et al., 2020). If the Arabidopsis *AtYSL1* to 3 are NA-metal (including Zn) transporters (Sinclair & Krämer, 2012), it is still not clear whether monocotyledonous YSL transporters can transport NA-bound Zn.

3.6 | Vacuolar iron transporter (VIT) family

VIT proteins can transport Fe, Mn and Zn into the vacuole for storage and detoxification (Brear et al., 2013). *AtVIT1* is important for Fe accumulation and partitioning in Arabidopsis seeds (S. A. Kim et al., 2006). Two rice proteins, *OsVIT1* and *OsVIT2*, localize to the tonoplast and although they have the ability to transport Zn in yeast, both appear to be mainly involved in Fe vacuolar sequestration in flag leaves (Figure 3a) (Y. Zhang et al., 2012). Note however that (i) the *OsVIT1* and *OsVIT2* expression is regulated by Zn and Fe supply (Y. Zhang et al., 2012); (ii) *OsVIT2* is among highly expressed genes in rice cultivars tolerant to Zn deficiency (Lu et al., 2021) and (iii) the corresponding mutants display decreased and increased Zn accumulation in flag leaves and grain embryo, respectively (Bashir, Takahashi, Akhtar, et al., 2013; Y. Zhang et al., 2012). Wheat homologs of *OsVITs*, *TaVIT1*, and *TaVIT2*, are downregulated during senescence (Pearce et al., 2014). Reduced vacuolar storage can facilitate nutrient remobilization from senescing tissues to reproductive organs. *TaVIT2* overexpression enhances Zn levels in barley grain endosperm but has no effect in wheat (Connorton et al., 2017). It is proposed to be mainly involved in Fe and Mn transport.

Finally, another rice VIT gene, *OsVIT5*, is highly expressed in panicles of cultivars with high Zn in grains and may contribute to grain Zn accumulation (Figure 3c) (Neeraja et al., 2018).

3.7 | NRAMP family

The Natural Resistance-Associated Macrophage Protein (NRAMP) family members act in proton/transition metal cation cotransport or antiport (Bozzi & Gaudet, 2021). *OsNRAMP5*, like *OsNAS3* and *OsVIT5*, is highly expressed in developing panicles of cultivars containing high Zn in grains (Neeraja et al., 2018). This gene however contributes mainly to Mn homeostasis and Cd accumulation (Sasaki et al., 2012).

OsNRAMP1 is highly induced in both roots and shoot under Zn excess and Fe deficiency. The Zn excess response is likely not a Zn-induced Fe-deficiency secondary effect (Ishimaru et al., 2008). However, *OsNRAMP1* also significantly contributes to Mn and Cd uptake in rice (Chang et al., 2020). In maize, *ZmNRAMP1* was recently shown to be induced by both Zn and Fe deficiencies in roots and shoot (Mallikarjuna et al., 2020).

4 | REGULATION OF Zn HOMEOSTASIS

Several transcription factors (TFs) belonging to basic Leucine Zipper (bZIP) and basic Helix-Loop-Helix (bHLH) widely found in Eukaryotes, as well as the plant-specific NAM, ATAF and CUC (NAC) and Abscisic acid Insensitive 3/Viviparous 1 (ABI3/VP1), families contribute to the regulation of Zn homeostasis in monocots.

The Arabidopsis bZIP19 and bZIP23 represent the main regulatory mechanism of the Zn homeostasis network in plants known to date (Assunção, Herrero, et al., 2010; Assunção, Schat, et al., 2010). Through Zinc-Deficiency Response Element (ZDRE) cis motifs, *AtbZIP19* and *AtbZIP23* control the upregulation of the Arabidopsis *IRT3*, *ZIP1*, *ZIP3*, *ZIP4*, *ZIP5*, *ZIP9*, *ZIP10*, *ZIP12*, *NAS2* and *NAS4* genes in response to Zn deficiency (Assunção, Herrero, et al., 2010; Assunção, Schat, et al., 2010; Castro et al., 2017). The two TFs have partially overlapping functions in controlling Zn homeostasis (Inaba et al., 2015; Lilay et al., 2019) and act directly as Zn sensors, via Zn binding by their Cys/His-rich motif that inactivates the TFs under sufficient Zn concentrations (Lilay et al., 2021).

Several F-group bZIPs, as *AtbZIP19* and *AtbZIP23*, are encoded in monocotyledonous plant genomes. Their expression in the Arabidopsis *bzip19bzip23* double mutant background has been extensively used to examine their function: the expression of *HvbZIP56* and *HvbZIP62* (Nazri et al., 2017), *TabZIPF1* and *TabZIPF4* (Evens et al., 2017), and *OsbZIP48* and *OsbZIP50* (Lilay et al., 2020) genes rescue the severe growth defect of the mutant upon Zn deficiency. *HvbZIP56*-overexpressing Arabidopsis lines also showed that *AtbZIP19/23* target genes are efficiently regulated by the ectopic expression of monocot bZIP genes (Nazri et al., 2017). These evidence are indicative of a conserved Zn deficiency regulatory network in plants (Castro et al., 2017).

A wheat NAC TF, Grain Protein Content 1 (GPC1) or NAM-B1, named after its Arabidopsis homolog No Apical Meristem (NAM), was suggested to be a central regulator of multiple processes during leaf senescence, including nutrient (Zn, Fe) and protein source-to-sink remobilization during grain filling (Uauy et al., 2006). The *OsNAC5* TF, highly expressed in flag leaves during grain filling, was suggested to have a

similar role in rice (Sperotto et al., 2009). RNAi or mutant lines for *GPC1* display delayed senescence and a 36% reduction of Zn concentration in wheat grains in lab and field conditions (Pearce et al., 2014). In *gpc1* plants, downregulation of the *TaHMA2-like* (ortholog of *OshMA2*), *TaZIP3* (ortholog of *OsZIP3*), *TaZIP7* (ortholog of *OsZIP7*), *TaZIP10-like1* (ortholog of *OsZIP10*), *TaZIP13-like2* (ortholog of *OsZIP8*) and *TaNAAT2* (ortholog of *OsNAAT1*) genes is observed, while they are all highly expressed during senescence (Pearce et al., 2014). These genes are possible downstream targets of *TaGPC1*.

Several bHLH TFs including FIT, bHLH38, 39, 100, 101, and PYE are known to mediate Fe deficiency responses through regulating Fe acquisition-related genes such as *AtFIT*, *AtNAS4*, *AtIRT1*, or the ferric chelate reductase *AtFRO2/AtFRO3* in Arabidopsis (Brumbarova et al., 2015; Gao et al., 2019; Kobayashi, Ozu, et al., 2019). Similarly, *OsiRO2*, a bHLH TF, is a regulator of Fe deficiency-inducible genes in rice, including NA and DMA synthesis genes such as *OsNAS1*, *OsNAS2*, *OsNAAT1* and *OsDMAS1*, and a DMA transporter gene *OsYSL15* (Ogo et al., 2007). Zn excess induces *OsiRO2*, its downstream genes, and consequently NA and DMA production in rice roots. In contrast, *OsiRT1* is not a target of *OsiRO2* and is not induced by Zn excess (Ishimaru et al., 2008).

The Iron Deficiency-responsive Element-binding Factor 1 (IDEF1) TF, belonging to the ABI3/VP1 family, is a positive regulator of *OsiRO2* (Kobayashi et al., 2007). *OsiDEF1* is a Fe deficiency-responsive TF that binds to Fe and other divalent metals, including Zn, and is important for Fe deficiency-responses and Fe tolerance in rice (Kobayashi et al., 2007, 2012). Some of these ZIFL genes are induced by both Fe deficiency and Zn excess (Ricachenevsky et al., 2011), and thus *IDEF1* may be involved in regulation of ZIFL family genes and Zn homeostasis in rice.

Besides transregulatory factors, epigenetic and posttranscriptional regulation, as well as hormones, contribute to the control of metal homeostasis in monocots. For instance, *OsMTP11* is highly induced in roots and shoot by Zn and Mn excess as well as Cd and Ni. This induction by metals may be regulated by changes in DNA methylation status of the *OsMTP11* promoter (M. Zhang & Liu, 2017). Modifications of histone methylation at the *OsZIP1* locus are suggested to control Cd stress response in rice (Liu et al., 2019). In *Sorghum bicolor*, 19 microRNAs (miRNAs) respond to Zn deficiency in roots or shoot. Interestingly, bZIP-binding ZDRE motifs were predicted in the promoter region of some of these miRNAs (Y. Li et al., 2013), suggesting a potentially important role for miRNAs in regulating Zn deficiency responses in Sorghum. Similarly, involvement of several miRNAs has been reported in regulation of Fe (Buhtz et al., 2010; Waters et al., 2012), Mn (Valdés-López et al., 2010) and Cu (Abdel-Ghany & Pilon, 2008; Pilon, 2017; Sunkar et al., 2006) homeostasis in Arabidopsis (Paul et al., 2015).

Recently, hormone biosynthetic and metabolic processes were shown to be enriched in the transcriptomic responses of Zn-deficient rice (Lu et al., 2020) and maize (Mallikarjuna et al., 2020) plants, as well as Zn-resupplied Zn-deficient *Brachypodium distachyon* plants (Amini et al., 2021). In these reports, jasmonic acid, auxin or gibberellic acid signaling pathways are suggested to be involved in the regulation of Zn homeostasis. Previously, auxin

signaling was linked to Zn uptake and transport under Zn deficiency in rice (Begum et al., 2016).

5 | INTERDEPENDENCE OF Zn HOMEOSTASIS AND OTHER NUTRIENTS

Plant nutrition, due to a sophisticated interaction network, requires a fine-tuned balance for the acquisition and distribution of micro- and macronutrients. In rice, for instance, Fe deficiency results in highly increased Zn content in roots and especially in shoot (Saenchai et al., 2016). Zn excess increases Fe concentration in roots, but results in slightly lower Fe concentration in shoot, where the expression of the Fe-responsive *OsFER1* is inhibited, indicating Fe deficiency in rice shoot (Ishimaru et al., 2008). However, the Fe excess-responsive marker gene, *OsFER1*, is not induced in rice roots, suggesting that Fe accumulates in the apoplast rather than in the cell cytoplasm (Ishimaru et al., 2008). On the other hand, the Fe-deficiency-induced *OsIRO2* gene is upregulated by Zn excess (Ishimaru et al., 2008), indicating that the *OsIRO2* TF is a potential mediator of Fe-Zn crosstalk in rice (Ricachenevsky et al., 2011). The presence of predicted *IRO2*-binding sites upstream of some *TaZIFL* genes was also suggested as a potential link between Fe and Zn responses (Ricachenevsky et al., 2011; Sharma et al., 2019). However, *IRO2*-binding sites were not found in the promoter region of rice *ZIFL* genes, and *IDEF1* was alternatively proposed as regulator of these genes (Ricachenevsky et al., 2011).

Fe-Zn interactions are well documented in dicots (Hanikenne et al., 2021), including a complex regulation of *AtIRT1* at the transcript and protein level (Connolly et al., 2002; Dubeaux et al., 2018; Fukao et al., 2011; Shanmugam et al., 2012). The transcript levels of Zn detoxification genes are on the other hand, upregulated in a FIT-dependent manner under Fe deficiency when *AtIRT1* is strongly upregulated (Arrivault et al., 2006; Colangelo & Gueriot, 2004; Sinclair & Krämer, 2012). Conversely, Fe supplementation attenuates Zn toxicity caused by Zn excess (Fukao et al., 2011; Lešková et al., 2017; Shanmugam et al., 2011; Zargar et al., 2015). Ferric Reductase Defective 3 (*AtFRD3*), which transports citrate into the xylem where it is required for Fe translocation to shoot (Durrett et al., 2007), is also involved in Fe-Zn interaction (Charlier et al., 2015; Pineau et al., 2012; Scheepers et al., 2020). In contrast, *OsIRT1* is not regulated by Zn excess (Ishimaru et al., 2008) and the functional homolog of *FRD3*, *OsFRDL1* (FRD-Like 1), seems to be less crucial in Fe/Zn homeostasis (Inoue et al., 2004; Yokosho et al., 2009, 2016; Yoneyama et al., 2015), suggesting a distinct impact of Zn on Fe homeostasis in *Arabidopsis* and rice.

Indeed, in monocots that use a chelation-based Fe uptake strategy, the competition between Zn and Fe(III) uptake from the soil, due to their different oxidation state, is expected to be less prevalent, or at least differ from the Zn/Fe(II) competition in reduction-based systems (Fukao et al., 2011). However, as evidenced by the response of Fe homeostasis genes to Zn excess in several species described above, it is clear that Zn excess nevertheless perturbs Fe in monocots. This is particularly true on calcareous soils, where grasses are

more sensitive to Zn toxicity than dicots, as Fe deficiency-induced PSs produced by roots strongly solubilize Zn (Chaney, 1993). In an investigation for Zn/Fe QTLs in wheat, only four QTL-rich clusters (QRC) were specific for Zn or Fe, and 10 out of 14 QRC were common for Zn and Fe, indicating a high interdependence between the homeostasis of these metal ions in wheat (Tong et al., 2020).

Similar to Fe, Mn concentration is increased and decreased respectively in rice roots and shoot upon Zn excess (Ishimaru et al., 2008). Conversely, the Mn concentration is reduced in roots of Zn-deficient wheat plants (Evens et al., 2017). Therefore, it is necessary to consider the interplay between Zn and other metals, in particular Fe, Mn and Cu when investigating metal homeostasis networks and regulatory pathways (Antosiewicz et al., 2014; Hanikenne et al., 2021).

Zn and phosphate (P) interaction is another well-studied nutrient cross-homeostasis in *Arabidopsis* (Bouain et al., 2014; Hanikenne et al., 2021; Khan et al., 2014; Kisko et al., 2018). For instance, the *AtPHR1* TF is involved in coordination of both P and Zn homeostasis. Zn deficiency leads on the other hand to upregulation of the *AtPHT1;1* gene, encoding a phosphate transporter (Jain et al., 2013), and the consequent over-accumulation of P in *Arabidopsis* shoot (Khan et al., 2014). The *Arabidopsis* *bZIP19/23* TFs were hypothesized, besides Zn, to be involved in Mn, Cu and P homeostasis (Lilay et al., 2019). Zn deficiency was previously demonstrated to induce P transporters in barley (Huang et al., 2000). Zn-deficient Sorghum plants accumulate a higher concentration of P and this is gradually reversed by Zn resupply (Y. Li et al., 2013). In rice, P deficiency leads to a reduced Zn concentration in shoot (Saenchai et al., 2016). In contrast, P excess substantially decreases Zn concentration in wheat shoot and grains (Ova et al., 2015). This P-Zn interaction in wheat is only observed in soil and was proposed to be mycorrhiza-dependent (Ova et al., 2015). On the other hand, synergistic interactions between Fe and P homeostasis were also reported (Saenchai et al., 2016), and therefore a complex crosstalk network between Zn, Fe and P homeostasis and signalling likely takes place and needs further examination. The *OsPHO1;1* gene, encoding a protein with a putative role in P transport through the endomembrane system, seems to be involved in such tripartite signalling in rice (Saenchai et al., 2016).

Finally, there is a strong interaction between Zn/Fe and nitrogen (N), especially for Zn and Fe storage in cereal grains (Guttieri et al., 2013; Persson et al., 2016). Zn concentration of wheat and maize grains increases considerably when N is supplied adequately (Cakmak et al., 2010; Xia et al., 2020, 2019). This interaction might be, at least partly, due to the role of N in the synthesis of transporter proteins and NA, but there is no report on a clear role of N in Zn homeostasis (Persson et al., 2016).

6 | CONCLUSIONS

Although comprehensive data are lacking to compare gene expression profiles during Zn deficiency responses in multiple species, it appears that key players and gene families, i.e. chelator synthesis

proteins, transporters and TFs, are conserved between Arabidopsis and grasses. Based on transcriptomic profiling of Fe-deficient rice and Arabidopsis plants, co-expression gene networks were built and compared (Grillet & Schmidt, 2019): the Fe uptake network is strikingly conserved in the two species and rice and Arabidopsis share more commonalities than differences. It will be relevant to conduct similar analyses with monocot species that specifically use the chelation-based strategy for Fe uptake and to extend it to other nutrients such as Zn or Cu. Yet, there are two notable differences between monocots and dicots concerning their Zn homeostasis, as noted above: the presence of PSs and an exodermis cell layer with its Casparian strip in grasses (Figure 1b,d).

While the gene families and their function remain similar, the main sources of differences in Zn homeostasis within monocots, and between monocots and dicots, stem from different growth environments, anatomical differences, and the use of NA and/or PS molecules.

7 | OUTLOOK

Two major obstacles hindering investigations of the Zn homeostasis molecular network in grasses are: (i) the lack of a proper model plant that represents wheat and barley, and (ii) most of our knowledge in rice, barley and wheat directly or indirectly originates from Arabidopsis. In particular, intracellular Zn trafficking transporters, to the mitochondria, vesicles or the Golgi complex, sensing, signalling and regulatory pathways behind Zn uptake, long-distance transport and redistribution toward seeds remain poorly known in both monocots and dicots. Application of comprehensive omics studies combined with targeted genetic approaches would help uncover distinct features of monocotyledonous Zn homeostasis, like the discovery of a novel NAC family TF in wheat (Uauy et al., 2006). Furthermore, using laser capture microdissection technology (Che et al., 2019; Tauris et al., 2009) or state-of-the-art single-cell tools (Denyer et al., 2019; Rodriguez-Villalon & Brady, 2019; Ryu et al., 2019; T. Q. Zhang et al., 2019) in studying the ionome and genetics of distinct cell and tissue types, will help to discover transporters that contribute to Zn transport across the biological barriers towards grains. As a consequence of these distinctions between rice and other grasses, it seems indispensable to perform independent studies in wheat and barley rather than translating the knowledge from rice (Box 2 and Box 3). Alternatively, Brachypodium may be used as a promising model system for wheat and barley plants (Brkljacic et al., 2011; Vogel et al., 2010), and was proposed as a model for Fe and Cu homeostasis studies in grasses (Jung et al., 2014; Yordem et al., 2011). This can be extended to Zn homeostasis (Amini et al., 2021). Improving Zn content of polished grains to tackle the hidden hunger in the world necessitates an even better understanding of molecular Zn players in cereal grasses.

ACKNOWLEDGEMENTS

Research funding in the laboratory is provided by F.R.S.-FNRS grants MIS F.4511.16, CDR J.0009.17, PDR T0120.18, and CDR-J.0082.21

(M. H.). S. A. is funded by a PhD grant from FZJ. M.H. was a Senior Research Associate of F.R.S.-FNRS.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

DATA AVAILABILITY STATEMENT

Data sharing not applicable—no new data generated.

ORCID

Borjana Arsova  <http://orcid.org/0000-0002-0566-2009>

Marc Hanikenne  <https://orcid.org/0000-0002-8964-9601>

REFERENCES

- Abdel-Ghany, S.E. & Pilon, M. (2008) MicroRNA-mediated systemic down-regulation of copper protein expression in response to low copper availability in Arabidopsis. *Journal of Biological Chemistry*, 283(23), 15932–15945. Available from: <https://doi.org/10.1074/jbc.M801406200>
- Ajiboye, B., Cakmak, I., Paterson, D., de Jonge, M.D., Howard, D.L., Stacey, S.P., et al. (2015) X-ray fluorescence microscopy of zinc localization in wheat grains biofortified through foliar zinc applications at different growth stages under field conditions. *Plant and Soil*, 392(1–2), 357–370. Available from: <https://doi.org/10.1007/s11104-015-2467-8>
- Akhtar, S. (2013) Zinc status in South Asian populations—an update. *Journal of Health, Population, and Nutrition*, 31(2), 139–149. Available from: <https://doi.org/10.3329/jhpn.v31i2.16378>
- Alloway, B.J. (2008) *Zinc in soils and crop nutrition*. Belgium: International Zinc Association Brussels.
- Alloway, B.J. (2009) 'Soil factors associated with zinc deficiency in crops and humans. *Environmental Geochemistry and Health*, 31(5), 537–548. Available from: <https://doi.org/10.1007/s10653-009-9255-4>
- Amini, S., Arsova, B., Gobert, S., Carnol, M., Bosman, B., Motte, P., et al. (2021) Transcriptional regulation of ZIP genes is independent of local zinc status in Brachypodium shoots upon zinc deficiency and resupply. *Plant, Cell and Environment*, 44, 3376–3397.
- Antosiewicz, D.M., Barabasz, A. & Siemianowski, O. (2014) Phenotypic and molecular consequences of overexpression of metal-homeostasis genes. *Frontiers in Plant Science*, 5, 80. Available from: <https://doi.org/10.3389/fpls.2014.00080>
- Anuradha, K., Agarwal, S., Rao, Y.V., Rao, K.V., Viraktamath, B.C. & Sarla, N. (2012) Mapping QTLs and candidate genes for iron and zinc concentrations in unpolished rice of Madhukar×Swarna RILs. *Gene*, 508(2), 233–240. Available from: <https://doi.org/10.1016/j.gene.2012.07.054>
- Araki, R., Murata, J. & Murata, Y. (2011) A novel barley yellow stripe 1-like transporter (HvYSL2) localized to the root endodermis transports metal-phytosiderophore complexes. *Plant and Cell Physiology*, 52(11), 1931–1940. Available from: <https://doi.org/10.1093/pcp/pcr126>
- Arnold, T., Kirk, G.J., Wissuwa, M., Frei, M., Zhao, F.J., Mason, T.F. et al. (2010) Evidence for the mechanisms of zinc uptake by rice using isotope fractionation. *Plant, Cell and Environment*, 33(3), 370–381. Available from: <https://doi.org/10.1111/j.1365-3040.2009.02085.x>
- Arrivault, S., Senger, T. & Krämer, U. (2006) The Arabidopsis metal tolerance protein AtMTP3 maintains metal homeostasis by mediating Zn exclusion from the shoot under Fe deficiency and Zn oversupply. *Plant Journal*, 46(5), 861–879. Available from: <https://doi.org/10.1111/j.1365-3113X.2006.02746.x>
- Assunção, A.G.L., Herrero, E., Lin, Y.F., Huettel, B., Talukdar, S., Smaczniak, C., et al. (2010) Arabidopsis thaliana transcription factors

- bZIP19 and bZIP23 regulate the adaptation to zinc deficiency. *Proceedings of the National Academy of Sciences*, 107, 10296–10301.
- Assunção, A.G.L., Schat, H. & Aarts, M.G.M. (2010) Regulation of the adaptation to zinc deficiency in plants. *Plant Signaling & Behavior*, 5(12), 1553–1555. Available from: <https://doi.org/10.4161/psb.5.12.13469>
- Balk, J., Connorton, J.M., Wan, Y., Lovegrove, A., Moore, K.L., Uauy, C., et al. (2019) Improving wheat as a source of iron and zinc for global nutrition. *Nutrition Bulletin*, 44(1), 53–59. Available from: <https://doi.org/10.1111/nbu.12361>
- Banakar, R., Alvarez Fernandez, A., Díaz-Benito, P., Abadia, J., Capell, T. & Christou, P. (2017) Phytosiderophores determine thresholds for iron and zinc accumulation in biofortified rice endosperm while inhibiting the accumulation of cadmium. *Journal of Experimental Botany*, 68(17), 4983–4995. Available from: <https://doi.org/10.1093/jxb/erx304>
- Bashir, K., Takahashi, R., Akhtar, S., Ishimaru, Y., Nakanishi, H. & Nishizawa, N.K. (2013) The knockdown of OsVIT2 and MIT affects iron localization in rice seed. *Rice*, 6(1), 1–6. Available from: <https://doi.org/10.1186/1939-8433-6-31>
- Bashir, K., Takahashi, R., Nakanishi, H. & Nishizawa, N.K. (2013) The road to micronutrient biofortification of rice: progress and prospects. *Frontiers of Plant Science*, 4, 15. Available from: <https://doi.org/10.3389/fpls.2013.00015>
- Baxter, I., Tchiew, J., Sussman, M.R., Boutry, M., Palmgren, M.G., Gribskov, M., et al. (2003) Genomic comparison of P-type ATPase ion pumps in arabidopsis and rice. *Plant Physiology*, 132(2), 618–628. Available from: <https://doi.org/10.1104/pp.103.021923>
- Becher, M., Talke, I.N., Krall, L. & Krämer, U. (2004) Cross-species microarray transcript profiling reveals high constitutive expression of metal homeostasis genes in shoots of the zinc hyperaccumulator *Arabidopsis halleri*. *The Plant Journal*, 37, 251–268. Available from: <https://doi.org/10.1046/j.1365-313X.2003.01959.x>
- Begum, M.C., Islam, M., Sarkar, M.R., Azad, M.A.S., Huda, A.K.M.N. & Kabir, A.H. (2016) Auxin signaling is closely associated with Zn-efficiency in rice (*Oryza sativa* L.). *Journal of Plant Interactions*, 11(1), 124–129. Available from: <https://doi.org/10.1080/17429145.2016.1220026>
- Black, R.E., Allen, L.H., Bhutta, Z.A., Caulfield, L.E., de Onis, M., Ezzati, M., et al. (2008) Maternal and child undernutrition: global and regional exposures and health consequences. *The Lancet*, 371(9608), 243–260. Available from: [https://doi.org/10.1016/S0140-6736\(07\)61690-0](https://doi.org/10.1016/S0140-6736(07)61690-0)
- Bouain, N., Shahzad, Z., Rouached, A., Khan, G.A., Berthomieu, P., Abdelly, C., et al. (2014) 'Phosphate and zinc transport and signalling in plants: toward a better understanding of their homeostasis interaction'. *Journal of Experimental Botany*, 65(20), 5725–5741. Available from: <https://doi.org/10.1093/jxb/eru314>
- Bozzi, A.T. & Gaudet, R. (2021) Molecular mechanism of nramp-family transition metal transport. *Journal of Molecular Biology*, 433, 166991. Available from: <https://doi.org/10.1016/j.jmb.2021.166991>
- Brear, E.M., Day, D.A. & Smith, P.M.C. (2013) Iron: an essential micronutrient for the legume-rhizobium symbiosis'. *Frontiers in Plant Science*, 4, 359. Available from: <https://doi.org/10.3389/fpls.2013.00359>
- Brkljajic, J., Grotewold, E., Scholl, R., Mockler, T., Garvin, D.F., Vain, P., et al. (2011) Brachypodium as a model for the grasses: today and the future. *Plant Physiology*, 157(1), 3–13. Available from: <https://doi.org/10.1104/pp.111.179531>
- Broadley, M.R., White, P.J., Hammond, J.P., Zelko, I. & Lux, A. (2007) Zinc in plants. *The New Phytologist*, 173(4), 677–702. Available from: <https://doi.org/10.1111/j.1469-8137.2007.01996.x>
- Brumbarova, T., Bauer, P. & Ivanov, R. (2015) Molecular mechanisms governing Arabidopsis iron uptake. *Trends in Plant Science*, 124–133. Available from: <https://doi.org/10.1016/j.tplants.2014.11.004>
- Buhtz, A., Pieritz, J., Springer, F. & Kehr, J. (2010) Phloem small RNAs, nutrient stress responses, and systemic mobility. *BMC Plant Biology*, 10(1), 1–13. Available from: <https://doi.org/10.1186/1471-2229-10-64>
- Cai, H., Huang, S., Che, J., Yamaji, N. & Ma, J.F. (2019) The tonoplast-localized transporter OsHMA3 plays an important role in maintaining Zn homeostasis in rice. *Journal of Experimental Botany*, 70(10), 2717–2725. Available from: <https://doi.org/10.1093/jxb/erz091>
- Cakmak, I. (2004) *Identification and correction of widespread zinc deficiency in Turkey: a success story; (a NATO-science for Stability Project)*. Paper Presented to The International Fertiliser Society at a Conference in Cambridge, on 10th December 2004, Fertiliser Society.
- Cakmak, I., Kalayci, M., Kaya, Y., Torun, A.A., Aydin, N., Wang, Y., et al. (2010) Biofortification and localization of zinc in wheat grain. *Journal of Agricultural and Food Chemistry*, 58(16), 9092–9102. Available from: <https://doi.org/10.1021/jf101197h>
- Cakmak, I. & Kutman, U.B. (2018) Agronomic biofortification of cereals with zinc: a review. *European Journal of Soil Science*, 69(1), 172–180. Available from: <https://doi.org/10.1111/ejss.12437>
- Cakmak, I., Sari, N., Marschner, H., Ekiz, H., Kalayci, M., Yilmaz, A. et al. (1996) Phytosiderophore release in bread and durum wheat genotypes differing in zinc efficiency. *Plant and Soil*, 180(2), 183–189. Available from: <https://doi.org/10.1007/BF00015301>
- Caldelas, C. & Weiss, D.J. (2017) Zinc homeostasis and isotopic fractionation in plants: a review. *Plant and Soil*, 411(1), 17–46. Available from: <https://doi.org/10.1007/s11104-016-3146-0>
- Castro, P.H., Lilay, G.H., Muñoz-Mérida, A., Schjoerring, J.K., Azevedo, H. & Assunção, A. (2017) Phylogenetic analysis of F-bZIP transcription factors indicates conservation of the zinc deficiency response across land plants. *Scientific Reports*, 7(1), 3806. Available from: <https://doi.org/10.1038/s41598-017-03903-6>
- Chandel, G., Banerjee, S., Vasconcelos, M. & Grusak, M.A. (2010) Characterization of the root transcriptome for iron and zinc homeostasis-related genes in indica rice (*Oryza sativa* L.). *Journal of Plant Biochemistry and Biotechnology*, 19(2), 145–152. Available from: <https://doi.org/10.1007/BF03263334>
- Chaney, R.L. (1993) Zinc phytotoxicity. A. D. Robson, *Zinc in soils and plants*. Netherlands: Springer, pp. Developments in Plant and Soil Sciences, 55, 135–150. https://doi.org/10.1007/978-94-011-0878-2_10
- Chang, J.D., Huang, S., Yamaji, N., Zhang, W., Ma, J.F. & Zhao, F.J. (2020) OsNRAMP1 transporter contributes to cadmium and manganese uptake in rice. *Plant, Cell and Environment*, 43(10), 2476–2491. Available from: <https://doi.org/10.1111/pce.13843>
- Chao, D.-Y., Silva, A., Baxter, I., Huang, Y.S., Nordborg, M. & Danku, J. et al. (2012) Genome-wide association studies identify heavy metal ATPase3 as the Primary determinant of natural variation in leaf cadmium in *Arabidopsis thaliana*. *PLOS Genetics*, 8(9), e1002923 <https://doi.org/10.1371/journal.pgen.1002923>
- Charlier, J.B., Polese, C., Nouet, C., Carnol, M., Bosman, B., Krämer, U., et al. (2015) Zinc triggers a complex transcriptional and post-transcriptional regulation of the metal homeostasis gene FRD3 in Arabidopsis relatives. *Journal of Experimental Botany*, 66(13), 3865–3878. Available from: <https://doi.org/10.1093/jxb/erv188>
- Che, J., Yokosho, K., Yamaji, N. & Ma, J.F. (2019) A vacuolar phytosiderophore transporter alters iron and zinc accumulation in polished rice grains. *Plant Physiology*, 181(1), 276–288. Available from: <https://doi.org/10.1104/pp.19.00598>
- Chen, Z., Fujii, Y., Yamaji, N., Masuda, S., Takemoto, Y., Kamiya, T., et al. (2013) Mn tolerance in rice is mediated by MTP8. 1, a member of the cation diffusion facilitator family. *Journal of Experimental Botany*, 64, 4375–4387.
- Claus, J., Bohmann, A. & Chavarría-Krauser, A. (2013) Zinc uptake and radial transport in roots of *Arabidopsis thaliana*: a modelling approach

- to understand accumulation. *Annals of Botany*, 112(2), 369–380. Available from: <https://doi.org/10.1093/aob/mcs263>
- Clemens, S. (2019) Metal ligands in micronutrient acquisition and homeostasis. *Plant Cell and Environment*. Blackwell Publishing Ltd, pp. 2902–2912. <https://doi.org/10.1111/pce.13627>
- Colangelo, E.P. & Guerinot, M.L. (2004) The essential basic helix-loop-helix protein FIT1 is required for the iron deficiency response. *The Plant Cell*, 16(12), 3400–3412. Available from: <https://doi.org/10.1105/tpc.104.024315>
- Connolly, E.L., Fett, J.P. & Guerinot, M.L. (2002) Expression of the IRT1 metal transporter is controlled by metals at the levels of transcript and protein accumulation. *The Plant Cell*, 14(6), 1347–1357. Available from: <https://doi.org/10.1105/tpc.001263>
- Connorton, J.M., Jones, E.R., Rodríguez-Ramiro, I., Fairweather-Tait, S., Uauy, C. & Balk, J. (2017) Wheat vacuolar iron transporter TaVIT2 transports Fe and Mn and is effective for biofortification. *Plant Physiology*, 174(4), 2434–2444. Available from: <https://doi.org/10.1104/pp.17.00672>
- Cornu, J.Y., Deinlein, U., Höreth, S., Braun, M., Schmidt, H., Weber, M., et al. (2015) Contrasting effects of nicotianamine synthase knockdown on zinc and nickel tolerance and accumulation in the zinc/cadmium hyperaccumulator *Arabidopsis halleri*. *New Phytologist*, 206(2), 738–750. Available from: <https://doi.org/10.1111/nph.13237>
- Curie, C., Panaviene, Z., Loulergue, C., Dellaporta, S.L., Briat, J.F. & Walker, E.L. (2001) Maize yellow stripe1 encodes a membrane protein directly involved in Fe(III) uptake. *Nature*, 409(6818), 346–349. Available from: <https://doi.org/10.1038/35053080>
- Delhaize, E., Gruber, B.D., Pittman, J.K., White, R.G., Leung, H., Miao, Y., et al. (2007) A role for the AtMTP11 gene of Arabidopsis in manganese transport and tolerance. *Plant Journal*, 51(2), 198–210. Available from: <https://doi.org/10.1111/j.1365-3113X.2007.03138.x>
- Denyer, T., Ma, X., Klesen, S., Scacchi, E., Nieselt, K. & Timmermans, M. (2019) Spatiotemporal developmental trajectories in the arabidopsis root revealed using high-throughput single-cell RNA sequencing. *Developmental Cell*, 48(6), 840–852.e5. Available from: <https://doi.org/10.1016/j.devcel.2019.02.022>
- Desbrosses-Fonrouge, A.-G., Voigt, K., Schröder, A., Arrivault, S., Thomine, S. & Krämer, U. (2005) Arabidopsis thaliana MTP1 is a Zn transporter in the vacuolar membrane which mediates Zn detoxification and drives leaf Zn accumulation. *FEBS Letters*, 579, 4165–4174. Available from: <https://doi.org/10.1016/j.febslet.2005.06.046>
- Detterbeck, A., Nagel, M., Rensch, S., Weber, M., Börner, A., Persson, D.P., et al. (2019) The search for candidate genes associated with natural variation of grain Zn accumulation in barley. *Biochemical Journal*, 476(13), 1889–1909. Available from: <https://doi.org/10.1042/bcj20190181>
- Díaz-Benito, P., Banakar, R., Rodríguez-Menéndez, S., Capell, T., Pereiro, R., Christou, P., et al. (2018) Iron and zinc in the embryo and endosperm of rice (*Oryza sativa* L.) Seeds in contrasting 2⁺-deoxymugineic acid/nicotianamine scenarios. *Frontiers of Plant Science*, 9, 1190. Available from: <https://doi.org/10.3389/fpls.2018.01190>
- Dubeaux, G., Neveu, J., Zelazny, E. & Vert, G. (2018) Metal sensing by the IRT1 Transporter-receptor orchestrates its own degradation and plant metal nutrition. *Molecular Cell*, 69(6), 953–964.e5. Available from: <https://doi.org/10.1016/j.molcel.2018.02.009>
- Durrett, T.P., Gassmann, W. & Rogers, E.E. (2007) The FRD3-mediated efflux of citrate into the root vasculature is necessary for efficient iron translocation. *Plant Physiology*, 144(1), 197–205. Available from: <https://doi.org/10.1104/pp.107.097162>
- Eckardt, N.A. (2000) Sequencing the rice genome. *The Plant Cell*, 12, 2011–2017. Available from: <https://doi.org/10.1105/tpc.12.11.2011>
- Evens, N.P., Buchner, P., Williams, L.E. & Hawkesford, M.J. (2017) The role of ZIP transporters and group F bZIP transcription factors in the Zn-deficiency response of wheat (*Triticum aestivum*). *Plant Journal*, 92(2), 291–304. Available from: <https://doi.org/10.1111/tpj.13655>
- Flis, P., Ouerdane, L., Grillet, L., Curie, C., Mari, S. & Lobinski, R. (2016) Inventory of metal complexes circulating in plant fluids: a reliable method based on HPLC coupled with dual elemental and high-resolution molecular mass spectrometric detection. *New Phytologist*, 211(3), 1129–1141. Available from: <https://doi.org/10.1111/nph.13964>
- Fukada, T., Yamasaki, S., Nishida, K., Murakami, M. & Hirano, T. (2011) Zinc homeostasis and signaling in health and diseases. *Journal of Biological Inorganic Chemistry*, 16(7), 1123–1134. Available from: <https://doi.org/10.1007/s00775-011-0797-4>
- Fukao, Y., Ferjani, A., Tomioka, R., Nagasaki, N., Kurata, R., Nishimori, Y., et al. (2011) iTRAQ analysis reveals mechanisms of growth defects due to excess zinc in Arabidopsis. *Plant Physiology*, 155(4), 1893–1907. Available from: <https://doi.org/10.1104/pp.110.169730>
- Gao, F., Robe, K., Gaymard, F., Izquierdo, E. & Dubos, C. (2019) The transcriptional control of iron homeostasis in plants: a tale of bHLH transcription factors?. *Frontiers in Plant Science*, 10, 6. Available from: <https://doi.org/10.3389/fpls.2019.00006>
- Gindri, R.G., Navarro, B.B., da Cruz Dias, P.V., Tarouco, C.P., Nicoloso, F.T., Brunetto, G., et al. (2020) Physiological responses of rice (*Oryza sativa* L.) *oszip7* loss-of-function plants exposed to varying Zn concentrations. *Physiology and Molecular Biology of Plants*, 26(7), 1349–1359. Available from: <https://doi.org/10.1007/s12298-020-00824-z>
- Graham, R.D. & Stangoulis, J.C.R. (2018) Trace element uptake and distribution in plants. *The Journal of Nutrition*, 133(5), 1502S–1505S. Available from: <https://doi.org/10.1093/jn/133.5.1502s>
- Gregory, P.J., Wahbi, A., Adu-Gyamfi, J., Heiling, M., Gruber, R., Joy, E.J.M., et al. (2017) Approaches to reduce zinc and iron deficits in food systems. *Global Food Security*, 15, 1–10. Available from: <https://doi.org/10.1016/j.gfs.2017.03.003>
- Grillet, L. & Schmidt, W. (2019) Iron acquisition strategies in land plants: not so different after all. *New Phytologist*, 224(1), 11–18. Available from: <https://doi.org/10.1111/nph.16005>
- Guerinot, M.L. (2000) The ZIP family of metal transporters. *Biochimica et Biophysica Acta-Biomembranes*, 1465(1–2), 190–198. Available from: [https://doi.org/10.1016/S0005-2736\(00\)00138-3](https://doi.org/10.1016/S0005-2736(00)00138-3)
- Gupta, N., Ram, H. & Kumar, B. (2016) Mechanism of Zinc absorption in plants: uptake, transport, translocation and accumulation. *Rev Environ Sci Biotechnol*, 15, 89–109. Available from: <https://doi.org/10.1007/s11157-016-9390-1>
- Guttieri, M.J., Stein, R.J. & Waters, B.M. (2013) Nutrient partitioning and grain yield of TaNAM-RNAi wheat under abiotic stress. *Plant and Soil*, 371(1–2), 573–591. Available from: <https://doi.org/10.1007/s11104-013-1713-1>
- Hafeez, B., Khanif, Y.M. & Saleem, M. (2013) Role of zinc in plant nutrition—a review. *American journal of experimental Agriculture*, 3(2), 374–391. Available from: <https://doi.org/10.9734/AJEA/2013/2746>
- Hands, P. & Drea, S. (2012) A comparative view of grain development in *Brachypodium distachyon*. *Journal of Cereal Science*, 56(1), 2–8. Available from: <https://doi.org/10.1016/j.jcs.2011.12.010>
- Hanikenne, M. & Baurain, D. (2014) Origin and evolution of metal P-type ATPases in Plantae (Archaeplastida). *Frontiers in Plant Science*, 4, 544. Available from: <https://doi.org/10.3389/fpls.2013.00544>
- Hanikenne, M., Esteves, S.M., Fanara, S. & Rouached, H. (2021) Coordinated homeostasis of essential mineral nutrients: a focus on iron. *Journal of Experimental Botany*, 72(6), 2136–2153. Available from: <https://doi.org/10.1093/jxb/eraa483>
- Hanikenne, M., Talke, I.N., Haydon, M.J., Lanz, C., Nolte, A., Motte, P., et al. (2008) Evolution of metal hyperaccumulation required

- cis-regulatory changes and triplication of HMA4. *Nature*, 453(7193), 391–395. Available from: <https://doi.org/10.1038/nature06877>
- Haydon, M.J. & Cobbett, C.S. (2007) A novel major facilitator superfamily protein at the tonoplast influences zinc tolerance and accumulation in Arabidopsis. *Plant Physiology*, 143(4), 1705–1719. Available from: <https://doi.org/10.1104/pp.106.092015>
- Haydon, M.J., Kawachi, M., Wirtz, M., Hillmer, S., Hell, R. & Krämer, U. (2012) Vacuolar nicotianamine has critical and distinct roles under iron deficiency and for zinc sequestration in Arabidopsis. *The Plant Cell*, 24(2), 724–737. Available from: <https://doi.org/10.1105/tpc.111.095042>
- Hegelund, J.N., Schiller, M., Kichey, T., Hansen, T.H., Pedas, P., Husted, S. et al. (2012) Barley metallothioneins: MT3 and MT4 are localized in the grain aleurone layer and show differential zinc binding. *Plant Physiology*, 159(3), 1125–1137. Available from: <https://doi.org/10.1104/pp.112.197798>
- Huang, C., Barker, S.J., Langridge, P., Smith, F.W. & Graham, R.D. (2000) Zinc deficiency up-regulates expression of high-affinity phosphate transporter genes in both phosphate-sufficient and -deficient barley roots. *Plant Physiology*, 124(1), 415–422. Available from: <https://doi.org/10.1104/pp.124.1.415>
- Hussain, D., Haydon, M.J., Wang, Y., Wong, E., Sherson, S.M., Young, J., et al. (2004) P-type ATPase heavy metal transporters with roles in essential zinc homeostasis in Arabidopsis. *The Plant Cell*, 16(5), 1327–1339. Available from: <https://doi.org/10.1105/tpc.020487>
- Inaba, S., Kurata, R., Kobayashi, M., Yamagishi, Y., Mori, I., Ogata, Y. et al. (2015) Identification of putative target genes of bZIP19, a transcription factor essential for Arabidopsis adaptation to Zn deficiency in roots. *Plant Journal*, 84(2), 323–334. Available from: <https://doi.org/10.1111/tbj.12996>
- Inoue, H., Higuchi, K., Takahashi, M., Nakanishi, H., Mori, S. & Nishizawa, N.K. (2003) Three rice nicotianamine synthase genes, OsNAS1, OsNAS2, and OsNAS3 are expressed in cells involved in long-distance transport of iron and differentially regulated by iron. *The Plant Journal*, 36(3), 366–381. Available from: <https://doi.org/10.1046/j.1365-313X.2003.01878.x>
- Inoue, H., Mizuno, D., Takahashi, M., Nakanishi, H., Mori, S. & Nishizawa, N.K. (2004) A rice FRD3-like (OsFRDL1) gene is expressed in the cells involved in long-distance transport. *Soil Science and Plant Nutrition*, 50(7), 1133–1140. Available from: <https://doi.org/10.1080/00380768.2004.10408586>
- International Brachypodium Initiative (2010) Genome sequencing and analysis of the model grass Brachypodium distachyon. *Nature*, 463(7282), 763–768. Available from: <https://doi.org/10.1038/nature08747>
- Ishimaru, Y., Suzuki, M., Kobayashi, T., Takahashi, M., Nakanishi, H., Mori, S. et al. (2005) OsZIP4, a novel zinc-regulated zinc transporter in rice. *Journal of Experimental Botany*, 56(422), 3207–3214. Available from: <https://doi.org/10.1093/jxb/eri317>
- Ishimaru, Y., Suzuki, M., Ogo, Y., Takahashi, M., Nakanishi, H., Mori, S. et al. (2008) Synthesis of nicotianamine and deoxymugineic acid is regulated by OsIRO2 in Zn excess rice plants. *Soil Science and Plant Nutrition*, 54(3), 417–423. Available from: <https://doi.org/10.1111/j.1747-0765.2008.00259.x>
- Ishimaru, Y., Suzuki, M., Tsukamoto, T., Suzuki, K., Nakazono, M., Kobayashi, T., et al. (2006) Rice plants take up iron as an Fe³⁺-phytosiderophore and as Fe²⁺. *Plant Journal*, 45(3), 335–346. Available from: <https://doi.org/10.1111/j.1365-313X.2005.02624.x>
- Jain, A., Sinilal, B., Dhandapani, G., Meagher, R.B. & Sahi, S.V. (2013) Effects of deficiency and excess of zinc on morphophysiological traits and spatiotemporal regulation of zinc-responsive genes reveal incidence of cross talk between micro- and macronutrients. *Environmental Science and Technology*, 47(10), 5327–5335. Available from: <https://doi.org/10.1021/es400113y>
- Jiang, W., Struik, P.C., Lingna, J., van Keulen, H., Ming, Z. & Stomph, T.J. (2007) Uptake and distribution of root-applied or foliar-applied 65 Zn after flowering in aerobic rice. *Annals of Applied Biology*, 150(3), 383–391. Available from: <https://doi.org/10.1111/j.1744-7348.2007.00138.x>
- Johnson, A.A., Kyriacou, B., Callahan, D.L., Carruthers, L., Stangoulis, J., Lombi, E. et al. (2011) Constitutive overexpression of the OsNAS gene family reveals single-gene strategies for effective iron- and zinc-biofortification of rice endosperm. *PLoS One*, 6(9), e24476. Available from: <https://doi.org/10.1371/journal.pone.0024476>
- Jung, H.I., Gayomba, S.R., Yan, J. & Vatamaniuk, O.K. (2014) Brachypodium distachyon as a model system for studies of copper transport in cereal crops. *Frontiers in Plant Science*, 5, 236. Available from: <https://doi.org/10.3389/fpls.2014.00236>
- Kavitha, P.G., Kuruvilla, S. & Mathew, M.K. (2015) Functional characterization of a transition metal ion transporter, OsZIP6 from rice (*Oryza sativa* L.). *Plant Physiology and Biochemistry*, 97, 165–174. Available from: <https://doi.org/10.1016/j.plaphy.2015.10.005>
- Kawakami, Y. & Bhullar, N.K. (2018) Molecular processes in iron and zinc homeostasis and their modulation for biofortification in rice. *Journal of Integrative Plant Biology*, 60(12), 1181–1198. Available from: <https://doi.org/10.1111/jipb.12751>
- Kaznina, N.M., Titov, A.F., Repkina, N.S. & Batova, Y.V. (2019) Effect of zinc excess and low temperature on the IRT1 gene expression in the roots and leaves of barley. *Doklady Biochemistry and Biophysics*, 487(1), 264–268. Available from: <https://doi.org/10.1134/S1607672919040057>
- Khan, G.A., Bouraine, S., Wege, S., Li, Y., de Carbonnel, M., Berthomieu, P., et al. (2014) Coordination between zinc and phosphate homeostasis involves the transcription factor PHR1, the phosphate exporter PHO1, and its homologue PHO1;H3 in Arabidopsis. *Journal of Experimental Botany*, 65(3), 871–884. Available from: <https://doi.org/10.1093/jxb/ert444>
- Kim, S.A., Punshon, T., Lanzirotti, A., Li, L., Alonso, J.M., et al. (2006) Localization of iron in Arabidopsis seed requires the vacuolar membrane transporter VIT1. *Science*, 314(5803), 1295–1298. Available from: <https://doi.org/10.1126/science.1132563>
- Kim, Y.Y., Choi, H., Segami, S., Cho, H.T., Martinoia, E., et al. (2009) AthHMA1 contributes to the detoxification of excess Zn(II) in Arabidopsis. *Plant Journal*, 58(5), 737–753. Available from: <https://doi.org/10.1111/j.1365-313X.2009.03818.x>
- Kisko, M., Bouain, N., Safi, A., Medici, A., Akkers, R.C., Secco, D., et al. (2018) LPCAT1 controls phosphate homeostasis in a zinc-dependent manner. *eLife*, 7, e32077. Available from: <https://doi.org/10.7554/eLife.32077>
- Kobayashi, T. (2019) Understanding the complexity of iron sensing and signaling cascades in plants. *Plant and Cell Physiology*, 60(7), 1440–1446. Available from: <https://doi.org/10.1093/pcp/pcz038>
- Kobayashi, T., Itai, R.N., Aung, M.S., Senoura, T., Nakanishi, H. & Nishizawa, N.K. (2012) The rice transcription factor IDEF1 directly binds to iron and other divalent metals for sensing cellular iron status. *The Plant Journal*, 69(1), 81–91. Available from: <https://doi.org/10.1111/j.1365-313X.2011.04772.x>
- Kobayashi, T., Ogo, Y., Itai, R.N., Nakanishi, H., Takahashi, M., Mori, S. et al. (2007) The transcription factor IDEF1 regulates the response to and tolerance of iron deficiency in plants. *Proceedings of the National Academy of Sciences of the United States of America*, 104(48), 19150–19155. Available from: <https://doi.org/10.1073/pnas.0707010104>
- Kobayashi, T., Ozu, A., Kobayashi, S., An, G., Jeon, J.S. & Nishizawa, N.K. (2019) OsbHLH058 and OsbHLH059 transcription factors positively regulate iron deficiency responses in rice. *Plant Molecular Biology*, 101(4–5), 471–486. Available from: <https://doi.org/10.1007/s11103-019-00917-8>

- Korshunova, Y.O., Eide, D., Clark, W.G., Guerinot, M.L. & Pakrasi, H.B. (1999) The IRT1 protein from *Arabidopsis thaliana* is a metal transporter with a broad substrate range. *Plant Molecular Biology*, 40(1), 37–44. Available from: <https://doi.org/10.1023/A:1026438615520>
- Lan, H.X., Wang, Z.F., Wang, Q.H., Wang, M.M., Bao, Y.M., et al. (2013) Characterization of a vacuolar zinc transporter OZT1 in rice (*Oryza sativa* L.). *Molecular Biology Reports*, 40(2), 1201–1210. Available from: <https://doi.org/10.1007/s11033-012-2162-2>
- Lee, S. & An, G. (2009) Over-expression of OsIRT1 leads to increased iron and zinc accumulations in rice. *Plant, Cell and Environment*, 32(4), 408–416. Available from: <https://doi.org/10.1111/j.1365-3040.2009.01935.x>
- Lee, S., Jeon, U.S., Lee, S.J., Kim, Y.K., Persson, D.P., et al. (2009) Iron fortification of rice seeds through activation of the nicotianamine synthase gene. *Proceedings of the National Academy of Sciences of the United States of America*, 106(51), 22014–22019. Available from: <https://doi.org/10.1073/pnas.0910950106>
- Lee, S., Jeong, H.J., Kim, S.A., Lee, J., Guerinot, M.L. et al. (2010) OsZIP5 is a plasma membrane zinc transporter in rice. *Plant Molecular Biology*, 73(4), 507–517. Available from: <https://doi.org/10.1007/s11103-010-9637-0>
- Lee, S., Kim, S.A., Lee, J., Guerinot, M.L. & An, G. (2010) Zinc deficiency-inducible OsZIP8 encodes a plasma membrane-localized zinc transporter in rice. *Molecules and Cells*, 29(6), 551–558. Available from: <https://doi.org/10.1007/s10059-010-0069-0>
- Lee, S., Kim, Y.Y., Lee, Y. & An, G. (2007) Rice P1B-type heavy-metal ATPase, OsHMA9, is a metal efflux protein. *Plant Physiology*, 145(3), 831–842. Available from: <https://doi.org/10.1104/pp.107.102236>
- Lee, S., Persson, D.P., Hansen, T.H., Husted, S., Schjoerring, J.K., et al. (2011) Bio-available zinc in rice seeds is increased by activation tagging of nicotianamine synthase. *Plant Biotechnology Journal*, 9(8), 865–873. Available from: <https://doi.org/10.1111/j.1467-7652.2011.00606.x>
- Lešková, A., Giehl, R., Hartmann, A., Fargašová, A. & von Wirén, N. (2017) Heavy metals induce iron deficiency responses at different hierarchic and regulatory levels. *Plant Physiology*, 174(3), 1648–1668. Available from: <https://doi.org/10.1104/pp.16.01916>
- Li, S., Liu, X., Zhou, X., Li, Y., Yang, W. & Chen, R. (2019) Improving zinc and iron accumulation in maize grains using the zinc and iron transporter ZmZIP5. *Plant and Cell Physiology*, 60(9), 2077–2085. Available from: <https://doi.org/10.1093/pcp/pcz104>
- Li, S., Zhou, X., Huang, Y., Zhu, L., Zhang, S., Zhao, Y., et al. (2013) Identification and characterization of the zinc-regulated transporters, iron-regulated transporter-like protein (ZIP) gene family in maize. *BMC Plant Biology*, 13(1), 114. Available from: <https://doi.org/10.1186/1471-2229-13-114>
- Li, S., Zhou, X., Li, H., Liu, Y., Zhu, L., Guo, J., et al. (2015) Overexpression of ZmIRT1 and ZmZIP3 enhances iron and zinc accumulation in transgenic *Arabidopsis*. *PLOS One*, 10(8), e0136647. Available from: <https://doi.org/10.1371/journal.pone.0136647>
- Li, S., Zhou, X., Zhao, Y., Li, H., Liu, Y., Zhu, L., et al. (2016) Constitutive expression of the ZmZIP7 in *Arabidopsis* alters metal homeostasis and increases Fe and Zn content. *Plant Physiology and Biochemistry*, 106, 1–10. Available from: <https://doi.org/10.1016/j.plaphy.2016.04.044>
- Li, Y., Zhang, Y., Shi, D., Liu, X., Qin, J., Ge, Q., et al. (2013) Spatial-temporal analysis of zinc homeostasis reveals the response mechanisms to acute zinc deficiency in *Sorghum bicolor*. *New Phytologist*, 200(4), 1102–1115. Available from: <https://doi.org/10.1111/nph.12434>
- Lilay, G.H., Castro, P.H., Campilho, A. & Assunção, A. (2019) The *Arabidopsis* bZIP19 and bZIP23 activity requires zinc deficiency—insight on regulation from complementation lines. *Frontiers in Plant Science*, 9, 1955. Available from: <https://doi.org/10.3389/fpls.2018.01955>
- Lilay, G.H., Castro, P.H., Guedes, J.G., Almeida, D.M., Campilho, A., Azevedo, H., et al. (2020) Rice F-bZIP transcription factors regulate the zinc deficiency response. *Journal of Experimental Botany*, 71(12), 3664–3677. Available from: <https://doi.org/10.1093/jxb/eraa115>
- Lilay, G.H., Persson, D.P., Castro, P.H., Liao, F., Alexander, R.D., Aarts, M. et al. (2021) *Arabidopsis* bZIP19 and bZIP23 act as zinc sensors to control plant zinc status. *Nature Plants*, 7(2), 137–143. Available from: <https://doi.org/10.1038/s41477-021-00856-7>
- Lin, Y.F., Hassan, Z., Talukdar, S., Schat, H. & Aarts, M.G. (2016) Expression of the Znt1 zinc transporter from the metal hyperaccumulator *Noccaea caerulea* confers enhanced zinc and cadmium tolerance and accumulation to *Arabidopsis thaliana*. *PLOS One*, 11(3), 0149750. Available from: <https://doi.org/10.1371/journal.pone.0149750>
- Lin, Y.F., Liang, H.M., Yang, S.Y., Boch, A., Clemens, S., et al. (2009) *Arabidopsis* IRT3 is a zinc-regulated and plasma membrane localized zinc/iron transporter. *New Phytologist*, 182(2), 392–404. Available from: <https://doi.org/10.1111/j.1469-8137.2009.02766.x>
- Liu, X.S., Feng, S.J., Zhang, B.Q., Wang, M.Q., Cao, H.W., et al. (2019) OsZIP1 functions as a metal efflux transporter limiting excess zinc, copper and cadmium accumulation in rice. *BMC Plant Biology*, 19(1), 283. Available from: <https://doi.org/10.1186/s12870-019-1899-3>
- Lombi, E., Smith, E., Hansen, T.H., Paterson, D., de Jonge, M.D., Howard, D.L., et al. (2011) Megapixel imaging of (micro)nutrients in mature barley grains. *Journal of Experimental Botany*, 62(1), 273–282. Available from: <https://doi.org/10.1093/jxb/erq270>
- Lu, X., Liu, S., Zhi, S., Chen, J. & Ye, G. (2021) Comparative transcriptome profile analysis of rice varieties with different tolerance to zinc deficiency. *Plant Biology*, 23(2), 375–390. Available from: <https://doi.org/10.1111/plb.13227>
- Malekzadeh, R., Shahpiri, A. & Siapoush, S. (2020) Metalation of a rice type 1 metallothionein isoform (OsMT1-1b). *Protein Expression and Purification*, 175, 105719. Available from: <https://doi.org/10.1016/j.pep.2020.105719>
- Mallikarjuna, M.G., Thirunavukkarasu, N., Sharma, R., Shiriga, K., Hossain, F., Bhat, J.S., et al. (2020) Comparative transcriptome analysis of iron and zinc deficiency in maize (*Zea mays* L.). *Plants*, 9(12), 1–31. Available from: <https://doi.org/10.3390/PLANTS9121812>
- Marschner, H., Romheld, V. & Kissel, M. (1986) Different strategies in higher plants in mobilization and uptake of iron. *Journal of Plant Nutrition*, 9(3–7), 695–713. Available from: <https://doi.org/10.1080/01904168609363475>
- Masuda, H., Usuda, K., Kobayashi, T., Ishimaru, Y., Kakei, Y., Takahashi, M., et al. (2009) Overexpression of the barley nicotianamine synthase gene HvNAS1 increases iron and zinc concentrations in rice grains. *Rice*, 2(4), 155–166. Available from: <https://doi.org/10.1007/s12284-009-9031-1>
- Mazé, P. (1915) Détermination des éléments minéraux rares nécessaires au développement du maïs. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*, 160, 211–214.
- Menguer, P.K., Farthing, E., Peaston, K.A., Ricachenevsky, F.K., Fett, J.P. & Williams, L.E. (2013) Functional analysis of the rice vacuolar zinc transporter OsMTP1. *Journal of Experimental Botany*, 64, 2871–2883. Available from: <https://doi.org/10.1093/jxb/ert136>
- Menguer, P.K., Vincent, T., Miller, A.J., Brown, J., Vincze, E., Borg, S., et al. (2018) Improving zinc accumulation in cereal endosperm using HvMTP1, a transition metal transporter. *Plant Biotechnology Journal*, 16(1), 63–71. Available from: <https://doi.org/10.1111/pbi.12749>
- Mikkelsen, M.D., Pedas, P., Schiller, M., Vincze, E., Mills, R.F., Borg, S., et al. (2012) Barley HvHMA1 is a heavy metal pump involved in mobilizing organellar Zn and Cu and plays a role in metal loading into

- grains. *PLOS One*, 7, e49027. Available from: <https://doi.org/10.1371/journal.pone.0049027>
- Mills, R.F., Peaston, K.A., Runions, J. & Williams, L.E. (2012) HvHMA2, a P1B-ATPase from barley, is highly conserved among cereals and functions in Zn and Cd transport. *PLOS One*, 7, e42640. Available from: <https://doi.org/10.1371/journal.pone.0042640>
- Miyadate, H., Adachi, S., Hiraizumi, A., Tezuka, K., Nakazawa, N., Kawamoto, T., et al. (2011) OsHMA3, a P1B-type of ATPase affects root-to-shoot cadmium translocation in rice by mediating efflux into vacuoles. *New Phytologist*, 189(1), 190–199. Available from: <https://doi.org/10.1111/j.1469-8137.2010.03459.x>
- Mizuno, D., Higuchi, K., Sakamoto, T., Nakanishi, H., Mori, S. & Nishizawa, N.K. (2003) Three nicotianamine synthase genes isolated from maize are differentially regulated by iron nutritional status. *Plant Physiology*, 132(4), 1989–1997. Available from: <https://doi.org/10.1104/pp.102.019869>
- Mondal, T.K., Ganie, S.A., Rana, M.K. & Sharma, T.R. (2014) Genome-wide analysis of zinc transporter genes of maize (*Zea mays*). *Plant Molecular Biology Reporter*, 32(2), 605–616. Available from: <https://doi.org/10.1007/s11105-013-0664-2>
- Montanini, B., Blaudez, D., Jeandroz, S., Sanders, D. & Chalot, M. (2007) Phylogenetic and functional analysis of the cation diffusion facilitator (CDF) family: improved signature and prediction of substrate specificity. *BMC Genomics*, 8(1), 1–16. Available from: <https://doi.org/10.1186/1471-2164-8-107>
- Morel, M., Crouzet, J., Gravot, A., Auroy, P., Leonhardt, N., Vavasseur, A. et al. (2009) AtHMA3, a P 1B -ATPase allowing Cd/Zn/Co/Pb vacuolar storage in Arabidopsis. *Plant Physiology*, 149(2), 894–904. Available from: <https://doi.org/10.1104/pp.108.130294>
- Mu, S., Yamaji, N., Sasaki, A., Luo, L., Du, B., Che, J., et al. (2021) A transporter for delivering zinc to the developing tiller bud and panicle in rice. *Plant Journal*, 105(3), 786–799. Available from: <https://doi.org/10.1111/tpj.15073>
- Nazri, A.Z., Griffin, J., Peaston, K.A., Alexander-Webber, D. & Williams, L.E. (2017) F-group bZIPs in barley—a role in Zn deficiency. *Plant, Cell and Environment*, 40(11), 2754–2770. Available from: <https://doi.org/10.1111/pce.13045>
- Neeraja, C.N., Kulkarni, K.S., Madhu Babu, P., Sanjeeva Rao, D., Surekha, K. & Ravindra Babu, V. (2018) Transporter genes identified in landraces associated with high zinc in polished rice through panicle transcriptome for biofortification. *PLOS One*, 13(2), e0192362. Available from: <https://doi.org/10.1371/journal.pone.0192362>
- Negishi, T., Nakanishi, H., Yazaki, J., Kishimoto, N., Fujii, F., Shimbo, K., et al. (2002) cDNA microarray analysis of gene expression during Fe-deficiency stress in barley suggests that polar transport of vesicles is implicated in phytosiderophore secretion in Fe-deficient barley roots. *Plant Journal*, 30(1), 83–94. Available from: <https://doi.org/10.1046/j.1365-313X.2002.01270.x>
- Nishiyama, R., Kato, M., Nagata, S., Yanagisawa, S. & Yoneyama, T. (2012) Identification of Zn-nicotianamine and Fe-2'-deoxymugineic acid in the phloem sap from rice plants (*Oryza sativa* L.). *Plant and Cell Physiology*, 53(2), 381–390. Available from: <https://doi.org/10.1093/pcp/pcr188>
- Nishizawa, N. & Mori, S. (1987) The particular vesicle appearing in barley root cells and its relation to mugineic acid secretion. *Journal of Plant Nutrition*, 10(9–16), 1013–1020. Available from: <https://doi.org/10.1080/01904168709363629>
- Nozoye, T., Nagasaka, S., Kobayashi, T., Sato, Y., Uozumi, N., Nakanishi, H. et al. (2015) The phytosiderophore efflux transporter TOM2 is involved in metal transport in rice. *Journal of Biological Chemistry*, 290(46), 27688–27699. Available from: <https://doi.org/10.1074/jbc.M114.635193>
- Nozoye, T., Nagasaka, S., Kobayashi, T., Takahashi, M., Sato, Y., Sato, Y., et al. (2011) Phytosiderophore efflux transporters are crucial for iron acquisition in graminaceous plants. *Journal of Biological Chemistry*, 286(7), 5446–5454. Available from: <https://doi.org/10.1074/jbc.M110.180026>
- Nozoye, T., Tsunoda, K., Nagasaka, S., Bashir, K., Takahashi, M., Kobayashi, T., et al. (2014) Rice nicotianamine synthase localizes to particular vesicles for proper function. *Plant Signaling and Behavior*, 9(5), e28660. Available from: <https://doi.org/10.4161/psb.28660>
- Nozoye, T., von Wirén, N., Sato, Y., Higashiyama, T., Nakanishi, H. & Nishizawa, N.K. (2019) Characterization of the nicotianamine exporter ENA1 in rice. *Frontiers in Plant Science*, 10, 10. Available from: <https://doi.org/10.3389/fpls.2019.00502>
- Ogo, Y., Itai, R.N., Nakanishi, H., Kobayashi, T., Takahashi, M., et al. (2007) The rice bHLH protein OsIRO2 is an essential regulator of the genes involved in Fe uptake under Fe-deficient conditions. *Plant Journal*, 51(3), 366–377. Available from: <https://doi.org/10.1111/j.1365-313X.2007.03149.x>
- Olsen, L.I., Hansen, T.H., Larue, C., Østerberg, J.T., Hoffmann, R.D., Liesche, J., et al. (2016) Mother-plant-mediated pumping of zinc into the developing seed. *Nature Plants*, 2(5), 1–6. Available from: <https://doi.org/10.1038/NPLANTS.2016.36>
- Olsen, L.I. & Palmgren, M.G. (2014) Many rivers to cross: the journey of zinc from soil to seed. *Frontiers in Plant Science*, 5, 30. Available from: <https://doi.org/10.3389/fpls.2014.00030>
- Østerberg, J.T. & Palmgren, M. (2018) Heavy metal pumps in plants: structure, function and origin. *Advances in Botanical Research*, 87, 57–89. Available from: <https://doi.org/10.1016/bs.abr.2018.09.004>
- Ova, E.A., Kutman, U.B., Ozturk, L. & Cakmak, I. (2015) High phosphorus supply reduced zinc concentration of wheat in native soil but not in autoclaved soil or nutrient solution. *Plant and Soil*, 393(1–2), 147–162. Available from: <https://doi.org/10.1007/s11104-015-2483-8>
- Palmgren, M.G., Clemens, S., Williams, L.E., Krämer, U., Borg, S., Schjørring, J.K. et al. (2008) Zinc biofortification of cereals: problems and solutions. *Trends in Plant Science*, 13(9), 464–473. Available from: <https://doi.org/10.1016/j.tplants.2008.06.005>
- Paul, S., Datta, S.K. & Datta, K. (2015) miRNA regulation of nutrient homeostasis in plants. *Frontiers in Plant Science*, 6, 232. Available from: <https://doi.org/10.3389/fpls.2015.00232>
- Pearce, S., Tabbita, F., Cantu, D., Buffalo, V., Avni, R., Vazquez-Gross, H., et al. (2014) Regulation of Zn and Fe transporters by the GPC1 gene during early wheat monocarpic senescence. *BMC Plant Biology*, 14(1), 1–23. Available from: <https://doi.org/10.1186/s12870-014-0368-2>
- Pedas, P., Schjoerring, J.K. & Husted, S. (2009) Identification and characterization of zinc-starvation-induced ZIP transporters from barley roots. *Plant Physiology and Biochemistry*, 47(5), 377–383. Available from: <https://doi.org/10.1016/j.plaphy.2009.01.006>
- Pedas, P., Ytting, C.K., Fuglsang, A.T., Jahn, T.P., Schjoerring, J.K. & Husted, S. (2008) Manganese efficiency in Barley: identification and characterization of the metal ion transporter HvIRT1. *Plant Physiology*, 148(1), 455–466. Available from: <https://doi.org/10.1104/pp.108.118851>
- Peiter, E., Montanini, B., Gobert, A., Pedas, P., Husted, S., Maathuis, F.J., et al. (2007) A secretory pathway-localized cation diffusion facilitator confers plant manganese tolerance. *Proceedings of the National Academy of Sciences of the United States of America*, 104(20), 8532–8537. Available from: <https://doi.org/10.1073/pnas.0609507104>
- Persson, D.P., de Bang, T.C., Pedas, P.R., Kutman, U.B., Cakmak, I., Andersen, B., et al. (2016) Molecular speciation and tissue compartmentation of zinc in durum wheat grains with contrasting nutritional status. *The New Phytologist*, 211(4), 1255–1265. Available from: <https://doi.org/10.1111/nph.13989>
- Pilon, M. (2017) The copper microRNAs. *New Phytologist*, 213(3), 1030–1035. Available from: <https://doi.org/10.1111/nph.14244>

- Pineau, C., Loubet, S., Lefoulon, C., Chalies, C., Fizames, C., Lacombe, B., et al. (2012) Natural variation at the FRD3 MATE transporter locus reveals cross-talk between Fe homeostasis and Zn tolerance in *Arabidopsis thaliana*. *PLoS Genetics*, 8(12), 1003120. Available from: <https://doi.org/10.1371/journal.pgen.1003120>
- Podar, D., Scherer, J., Noordally, Z., Herzyk, P., Nies, D. & Sanders, D. (2012) Metal selectivity determinants in a family of transition metal transporters. *Journal of Biological Chemistry*, 287(5), 3185–3196. Available from: <https://doi.org/10.1074/jbc.M111.305649>
- Ram, H., Kaur, A., Gandass, N., Singh, S., Deshmukh, R., et al. (2019) Molecular characterization and expression dynamics of MTP genes under various spatio-temporal stages and metal stress conditions in rice. *PLOS One*, 14, e0217360. Available from: <https://doi.org/10.1371/journal.pone.0217360>
- Ramesh, S.A., Shin, R., Eide, D.J. & Schachtman, D.P. (2003) Differential metal selectivity and gene expression of two zinc transporters from rice. *Plant Physiology*, 133(1), 126–134. Available from: <https://doi.org/10.1104/pp.103.026815>
- Rawat, N., Neelam, K., Tiwari, V.K. & Dhaliwal, H.S. (2013) Biofortification of cereals to overcome hidden hunger. *Plant Breeding*, 132(5), 437–445. Available from: <https://doi.org/10.1111/pbr.12040>
- Ricachenevsky, F.K., Punshon, T., Lee, S., Oliveira, B., Trenz, T.S., Maraschin, F., et al. (2018) Elemental profiling of rice FOX lines leads to characterization of a new Zn plasma membrane transporter, OsZIP7. *Frontiers in Plant Science*, 9, 865. Available from: <https://doi.org/10.3389/fpls.2018.00865>
- Ricachenevsky, F.K., Sperotto, R.A., Menguer, P.K., Sperb, E.R., Lopes, K.L. & Fett, J.P. (2011) Zinc-induced facilitator-like family in plants: lineage-specific expansion in monocotyledons and conserved genomic and expression features among rice (*Oryza sativa*) paralogs. *BMC Plant Biology*, 11(1), 1–22. Available from: <https://doi.org/10.1186/1471-2229-11-20>
- Rodriguez-Villalon, A. & Brady, S.M. (2019) Single cell RNA sequencing and its promise in reconstructing plant vascular cell lineages. *Current Opinion in Plant Biology*, 48, 47–56. Available from: <https://doi.org/10.1016/j.pbi.2019.04.002>
- Rogers, E.E., Eide, D.J. & Gueriot, M.L. (2000) Altered selectivity in an *Arabidopsis* metal transporter. *Proceedings of the National Academy of Sciences of the United States of America*, 97(22), 12356–12360. Available from: <https://doi.org/10.1073/pnas.210214197>
- Ryu, K.H., Huang, L., Kang, H.M. & Schiefelbein, J. (2019) Single-cell RNA sequencing resolves molecular relationships among individual plant cells. *Plant Physiology*, 179(4), 1444–1456. Available from: <https://doi.org/10.1104/pp.18.01482>
- Saenchai, C., Bouain, N., Kisko, M., Prom-U-Thai, C., Dumas, P. & Rouached, H. (2016) The involvement of OsPHO1;1 in the regulation of iron transport through integration of phosphate and Zinc deficiency signaling. *Frontiers in Plant Science*, 7, 396. Available from: <https://doi.org/10.3389/fpls.2016.00396>
- Salt, D.E., Prince, R.C., Baker, A.J.M., Raskin, I. & Pickering, I.J. (1999) Zinc ligands in the metal hyperaccumulator *Thlaspi caerulescens* as determined using x-ray absorption spectroscopy. *Environmental Science and Technology*, 33(5), 713–717. Available from: <https://doi.org/10.1021/es980825x>
- Sasaki, A., Yamaji, N. & Ma, J.F. (2014) Overexpression of OsHMA3 enhances Cd tolerance and expression of Zn transporter genes in rice. *Journal of Experimental Botany*, 65(20), 6013–6021. Available from: <https://doi.org/10.1093/jxb/eru340>
- Sasaki, A., Yamaji, N., Mitani-Ueno, N., Kashino, M. & Ma, J.F. (2015) A node-localized transporter OsZIP3 is responsible for the preferential distribution of Zn to developing tissues in rice. *Plant Journal*, 84(2), 374–384. Available from: <https://doi.org/10.1111/tpj.13005>
- Sasaki, A., Yamaji, N., Yokosho, K. & Ma, J.F. (2012) Nramp5 is a major transporter responsible for manganese and cadmium uptake in rice. *The Plant Cell*, 24(5), 2155–2167. Available from: <https://doi.org/10.1105/tpc.112.096925>
- Sasaki, T. & Burr, B. (2000) International rice genome sequencing project: the effort to completely sequence the rice genome. *Current Opinion in Plant Biology*, 3(2), 138–142. Available from: [https://doi.org/10.1016/S1369-5266\(99\)00047-3](https://doi.org/10.1016/S1369-5266(99)00047-3)
- Satoh-Nagasawa, N., Mori, M., Nakazawa, N., Kawamoto, T., Nagato, Y., Sakurai, K., et al. (2012) Mutations in rice (*oryza sativa*) heavy metal ATPase 2 (OsHMA2) restrict the translocation of zinc and cadmium. *Plant and Cell Physiology*, 53(1), 213–224. Available from: <https://doi.org/10.1093/pcp/pcr166>
- Schaaf, G., Ludewig, U., Erenoglu, B.E., Mori, S., Kitahara, T. & von Wirén, N. (2004) ZmYS1 functions as a proton-coupled symporter for phytosiderophore- and nicotianamine-chelated metals. *Journal of Biological Chemistry*, 279(10), 9091–9096. Available from: <https://doi.org/10.1074/jbc.M311799200>
- Scheepers, M., Spielmann, J., Boulanger, M., Carnol, M., Bosman, B., De Pauw, E., et al. (2020) Intertwined metal homeostasis, oxidative and biotic stress responses in the *Arabidopsis frd3* mutant. *The Plant Journal*, 102(1), 34–52. Available from: <https://doi.org/10.1111/tpj.14610>
- Seigneurin-Berny, D., Gravot, A., Auroy, P., Mazard, C., Kraut, A., Finazzi, G., et al. (2006) HMA1, a new Cu-ATPase of the chloroplast envelope, is essential for growth under adverse light conditions. *Journal of Biological Chemistry*, 281(5), 2882–2892. Available from: <https://doi.org/10.1074/jbc.M508333200>
- Shahzad, Z., Rouached, H. & Rakha, A. (2014) Combating mineral malnutrition through iron and zinc biofortification of cereals. *Comprehensive Reviews in Food Science and Food Safety*, 13(3), 329–346. Available from: <https://doi.org/10.1111/1541-4337.12063>
- Shanmugam, V., Lo, J.C., Wu, C.L., Wang, S.L., Lai, C.C., Connolly, E.L., et al. (2011) Differential expression and regulation of iron-regulated metal transporters in *Arabidopsis halleri* and *Arabidopsis thaliana*—the role in zinc tolerance. *New Phytologist*, 190(1), 125–137. Available from: <https://doi.org/10.1111/j.1469-8137.2010.03606.x>
- Shanmugam, V., Tsednee, M. & Yeh, K.C. (2012) Zinc tolerance induced by iron 1 reveals the importance of glutathione in the cross-homeostasis between zinc and iron in *Arabidopsis thaliana*. *Plant Journal*, 69(6), 1006–1017. Available from: <https://doi.org/10.1111/j.1365-3113X.2011.04850.x>
- Sharma, S., Kaur, G., Kumar, A., Meena, V., Kaur, J. & Pandey, A.K. (2019) Overlapping transcriptional expression response of wheat zinc-induced facilitator-like transporters emphasize important role during Fe and Zn stress. *BMC Molecular Biology*, 20(1), 1–17. Available from: <https://doi.org/10.1186/s12867-019-0139-6>
- Shojima, S., Nishizawa, N.K., Fushiya, S., Nozoe, S., Irifune, T. & Mori, S. (1990) Biosynthesis of phytosiderophores: In vitro biosynthesis of 2'-deoxymugineic acid from L-methionine and nicotianamine. *Plant Physiology*, 93(4), 1497–1503. Available from: <https://doi.org/10.1104/pp.93.4.1497>
- Sinclair, S.A. & Krämer, U. (2012) The zinc homeostasis network of land plants. *Biochimica et Biophysica Acta-Molecular Cell Research*, 1823(9), 1553–1567. Available from: <https://doi.org/10.1016/j.bbamcr.2012.05.016>
- Sinclair, S.A., Senger, T., Talke, I.N., Cobbett, C.S., Haydon, M.J. & Krämer, U. (2018) Systemic upregulation of MTP2- and HMA2-mediated Zn partitioning to the shoot supplements local Zn deficiency responses. *The Plant Cell*, 30(10), 2463–2479. Available from: <https://doi.org/10.1105/tpc.18.00207>
- Singh, S.P., Keller, B., Gruissem, W. & Bhullar, N.K. (2017) Rice nicotianamine synthase 2 expression improves dietary iron and zinc levels in wheat. *Theoretical and Applied Genetics*, 130(2), 283–292. Available from: <https://doi.org/10.1007/s00122-016-2808-x>

- Sommer, A.L. & Lipman, C.B. (1926) Evidence on the indispensable nature of zinc and boron for higher green plants. *Plant Physiology*, 1(3), 231–249. Available from: <https://doi.org/10.1104/pp.1.3.231>
- Sperotto, R.A. (2013) Zn/Fe remobilization from vegetative tissues to rice seeds: Should I stay or should I go? Ask Zn/Fe supply!. *Frontiers in Plant Science*, 4, 464. Available from: <https://doi.org/10.3389/fpls.2013.00464>
- Sperotto, R.A., Ricachenevsky, F.K., Duarte, G.L., Boff, T., Lopes, K.L., Sperb, E.R., et al. (2009) Identification of up-regulated genes in flag leaves during rice grain filling and characterization of OsNAC5, a new ABA-dependent transcription factor. *Planta*, 230(5), 985–1002. Available from: <https://doi.org/10.1007/s00425-009-1000-9>
- Spielmann, J., Ahmadi, H., Scheepers, M., Weber, M., Nitsche, S., Carnol, M., et al. (2020) The two copies of the zinc and cadmium ZIP6 transporter of *Arabidopsis halleri* have distinct effects on cadmium tolerance. *Plant, Cell & Environment*, 43(9), 2143–2157. Available from: <https://doi.org/10.1111/pce.13806>
- Stomph, T.Jan, Jiang, W. & Struik, P.C. (2009) Zinc biofortification of cereals: rice differs from wheat and barley. *Trends in Plant Science*, 14(3), 123–124. Available from: <https://doi.org/10.1016/j.tplants.2009.01.001>
- Sunkar, R., Kapoor, A. & Zhu, J.K. (2006) Posttranscriptional induction of two Cu/Zn superoxide dismutase genes in *Arabidopsis* is mediated by downregulation of miR398 and important for oxidative stress tolerance. *The Plant Cell*, 18(8), 2051–2065. Available from: <https://doi.org/10.1105/tpc.106.041673>
- Suzuki, M., Bashir, K., Inoue, H., Takahashi, M., Nakanishi, H. & Nishizawa, N.K. (2012) Accumulation of starch in Zn-deficient rice. *Rice*, 5(1), 1–8. Available from: <https://doi.org/10.1186/1939-8433-5-9>
- Suzuki, M., Takahashi, M., Tsukamoto, T., Watanabe, S., Matsuhashi, S., Yazaki, J., et al. (2006) Biosynthesis and secretion of mugineic acid family phytosiderophores in zinc-deficient barley. *Plant Journal*, 48(1), 85–97. Available from: <https://doi.org/10.1111/j.1365-313X.2006.02853.x>
- Suzuki, M., Tsukamoto, T., Inoue, H., Watanabe, S., Matsuhashi, S., Takahashi, M., et al. (2008) Deoxymugineic acid increases Zn translocation in Zn-deficient rice plants. *Plant Molecular Biology*, 66(6), 609–617. Available from: <https://doi.org/10.1007/s11103-008-9292-x>
- Swamy, B., Rahman, M.A., Inabangan-Asilo, M.A., Amparado, A., Manito, C., Chadha-Mohanty, P., et al. (2016) Advances in breeding for high grain zinc in rice. *Rice*, 9(1), 1–16. Available from: <https://doi.org/10.1186/s12284-016-0122-5>
- Takahashi, R., Ishimaru, Y., Shimo, H., Ogo, Y., Senoura, T., Nishizawa, N.K. et al. (2012) The OsHMA2 transporter is involved in root-to-shoot translocation of Zn and Cd in rice. *Plant, Cell and Environment*, 35(11), 1948–1957. Available from: <https://doi.org/10.1111/j.1365-3040.2012.02527.x>
- Talke, I.N., Hanikenne, M. & Krämer, U. (2006) Zinc-dependent global transcriptional control, transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of the hyperaccumulator *Arabidopsis halleri*. *Plant Physiology*, 142(1), 148–167. Available from: <https://doi.org/10.1104/pp.105.076232>
- Tan, J., Wang, J., Chai, T., Zhang, Y., Feng, S., et al. (2013) Functional analyses of TaHMA2, a P 1B -type ATPase in wheat. *Plant Biotechnology Journal*, 11(4), 420–431. Available from: <https://doi.org/10.1111/pbi.12027>
- Tan, L., Qu, M., Zhu, Y., Peng, C., Wang, J., et al. (2020) Zinc transporter5 and zinc transporter9 function synergistically in zinc/cadmium uptake1. *Plant Physiology*, 183(3), 1235–1249. Available from: <https://doi.org/10.1104/pp.19.01569>
- Tan, L., Zhu, Y., Fan, T., Peng, C., Wang, J., et al. (2019) OsZIP7 functions in xylem loading in roots and inter-vascular transfer in nodes to deliver Zn/Cd to grain in rice. *Biochemical and Biophysical Research Communications*, 512(1), 112–118. Available from: <https://doi.org/10.1016/j.bbrc.2019.03.024>
- Tang, H., Bowers, J.E., Wang, X. & Paterson, A.H. (2010) Angiosperm genome comparisons reveal early polyploidy in the monocot lineage. *Proceedings of the National Academy of Sciences of the United States of America*, 107(1), 472–477. Available from: <https://doi.org/10.1073/pnas.0908007107>
- Tauris, B., Borg, S., Gregersen, P.L. & Holm, P.B. (2009) A roadmap for zinc trafficking in the developing barley grain based on laser capture microdissection and gene expression profiling. *Journal of Experimental Botany*, 60(4), 1333–1347. Available from: <https://doi.org/10.1093/jxb/erp023>
- Tiong, J., McDonald, G., Genc, Y., Shirley, N., Langridge, P. & Huang, C.Y. (2015) Increased expression of six ZIP family genes by zinc (Zn) deficiency is associated with enhanced uptake and root-to-shoot translocation of Zn in barley (*Hordeum vulgare*). *New Phytologist*, 207(4), 1097–1109. Available from: <https://doi.org/10.1111/nph.13413>
- Tiong, J., McDonald, G.K., Genc, Y., Pedas, P., Hayes, J.E., Toubia, J., et al. (2014) HvZIP7 mediates zinc accumulation in barley (*Hordeum vulgare*) at moderately high zinc supply. *New Phytologist*, 201(1), 131–143. Available from: <https://doi.org/10.1111/nph.12468>
- Tong, J., Sun, M., Wang, Y., Zhang, Y., Rasheed, A., Li, M., et al. (2020) Dissection of molecular processes and genetic architecture underlying iron and zinc homeostasis for biofortification: From model plants to common wheat. *International Journal of Molecular Sciences*, 21(23), 9280. Available from: <https://doi.org/10.3390/ijms21239280>
- Turgeon, R. & Wolf, S. (2009) Phloem transport: cellular pathways and molecular trafficking. *Annual Review of Plant Biology*, 60, 207–221. Available from: <https://doi.org/10.1146/annurev.arplant.043008.092045>
- Uauy, C., Distelfeld, A., Fahima, T., Blechl, A. & Dubcovsky, J. (2006) A NAC gene regulating senescence improves grain protein, zinc, and iron content in wheat. *Science*, 314(5803), 1298–1301. Available from: <https://doi.org/10.1126/science.1133649>
- Ueno, D., Sasaki, A., Yamaji, N., Miyaji, T., Fujii, Y., Takemoto, Y., et al. (2015) A polarly localized transporter for efficient manganese uptake in rice. *Nature plants*, 1(12), 1–8. Available from: <https://doi.org/10.1038/nplants.2015.170>
- Ueno, D., Yamaji, N., Kono, I., Huang, C.F., Ando, T., Yano, M. et al. (2010) Gene limiting cadmium accumulation in rice. *Proceedings of the National Academy of Sciences of the United States of America*, 107(38), 16500–16505. Available from: <https://doi.org/10.1073/pnas.1005396107>
- Valdés-López, O., Yang, S.S., Aparicio-Fabre, R., Graham, P.H., Reyes, J.L., Vance, C.P. et al. (2010) MicroRNA expression profile in common bean (*Phaseolus vulgaris*) under nutrient deficiency stresses and manganese toxicity. *New Phytologist*, 187(3), 805–818. Available from: <https://doi.org/10.1111/j.1469-8137.2010.03320.x>
- Vallee, B.L. & Falchuk, K.H. (1993) The biochemical basis of zinc physiology. *Physiological Reviews*, 73(1), 79–118. Available from: <https://doi.org/10.1152/physrev.1993.73.1.79>
- Vert, G., Grotz, N., Dédaldéchamp, F., Gaymard, F., Gueriot, M.L., Briat, J.F. et al. (2002) 'IRT1, an Arabidopsis transporter essential for iron uptake from the soil and for plant growth'. *The Plant Cell*, 14(6), 1223–1233. Available from: <https://doi.org/10.1105/tpc.001388>
- Wang, H.L., Offler, C.E. & Patrick, J.W. (1994) Nucellar projection transfer cells in the developing wheat grain. *Protoplasma*, 182(1–2), 39–52. Available from: <https://doi.org/10.1007/BF01403687>
- Waters, B.M., McInturf, S.A. & Stein, R.J. (2012) Rosette iron deficiency transcript and microRNA profiling reveals links between copper and iron homeostasis in *Arabidopsis thaliana*. *Journal of Experimental Botany*, 63(16), 5903–5918. Available from: <https://doi.org/10.1093/jxb/ers239>

- Waters, B.M., Uauy, C., Dubcovsky, J. & Grusak, M.A. (2009) Wheat (*Triticum aestivum*) NAM proteins regulate the translocation of iron, zinc, and nitrogen compounds from vegetative tissues to grain. *Journal of Experimental Botany*, 60(15), 4263–4274. Available from: <https://doi.org/10.1093/jxb/erp257>
- White, P.J. & Broadley, M.R. (2009) Biofortification of crops with seven mineral elements often lacking in human diets—iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytologist*, 182(1), 49–84. Available from: <https://doi.org/10.1111/j.1469-8137.2008.02738.x>
- White, P.J. (2001) The pathways of calcium movement to the xylem. *Journal of Experimental Botany*, 52(358), 891–899. Available from: <https://doi.org/10.1093/jexbot/52.358.891>
- Williams, L.E. & Mills, R.F. (2005) P1B-ATPases—an ancient family of transition metal pumps with diverse functions in plants. *Trends in Plant Science*, 10(10), 491–502. Available from: <https://doi.org/10.1016/j.tplants.2005.08.008>
- Von Wirén, N., Marschner, H. & Römheld, V. (1996) Roots of iron-efficient maize also absorb phytosiderophore-chelated zinc. *Plant Physiology*, 111(4), 1119–1125. Available from: <https://doi.org/10.1104/pp.111.4.1119>
- Wirth, J., Poletti, S., Aeschlimann, B., Yakandawala, N., Drosse, B., Osorio, S., et al. (2009) Rice endosperm iron biofortification by targeted and synergistic action of nicotianamine synthase and ferritin. *Plant Biotechnology Journal*, 7(7), 631–644. Available from: <https://doi.org/10.1111/j.1467-7652.2009.00430.x>
- Wu, C.Y., Lu, L.L., Yang, X.E., Feng, Y., Wei, Y.Y., Hao, H.L., et al. (2010) Uptake, translocation, and remobilization of zinc absorbed at different growth stages by rice genotypes of different Zn densities. *Journal of Agricultural and Food Chemistry*, 58(11), 6767–6773. Available from: <https://doi.org/10.1021/jf100017e>
- Xia, H. et al. (2020) Elucidating the source–sink relationships of zinc biofortification in wheat grains: a review. *Food and Energy Security*, 9(4), e243. Available from: <https://doi.org/10.1002/fes3.243>
- Xue, Y.-F., Yue, S.C., Liu, D.Y., Zhang, W., Chen, X.P. et al. (2019) Dynamic zinc accumulation and contributions of Pre- and/or post-silking zinc uptake to grain zinc of maize as affected by nitrogen supply. *Frontiers in Plant Science*, 10, 1203. Available from: <https://doi.org/10.3389/fpls.2019.01203>
- Yamaguchi, N., Ishikawa, S., Abe, T., Baba, K., Arao, T. & Terada, Y. (2012) Role of the node in controlling traffic of cadmium, zinc, and manganese in rice. *Journal of Experimental Botany*, 63(7), 2729–2737. Available from: <https://doi.org/10.1093/jxb/err455>
- Yamaji, N., Sakurai, G., Mitani-Ueno, N. & Ma, J.F. (2015) Orchestration of three transporters and distinct vascular structures in node for intervascular transfer of silicon in rice. *Proceedings of the National Academy of Sciences of the United States of America*, 112(36), 11401–11406. Available from: <https://doi.org/10.1073/pnas.1508987112>
- Yamaji, N., Xia, J., Mitani-Ueno, N., Yokosho, K. & Feng Ma, J. (2013) Preferential delivery of zinc to developing tissues in rice is mediated by P-type heavy metal ATPase OsHMA2. *Plant Physiology*, 162(2), 927–939. Available from: <https://doi.org/10.1104/pp.113.216564>
- Yang, M., Li, Y., Liu, Z., Tian, J., Liang, L., Qiu, Y., et al. (2020) A high activity zinc transporter OsZIP9 mediates zinc uptake in rice. *Plant Journal*, 103(5), 1695–1709. Available from: <https://doi.org/10.1111/tpj.14855>
- Yang, X., Huang, J., Jiang, Y. & Zhang, H.S. (2009) Cloning and functional identification of two members of the ZIP (ZRT, IRT-like protein) gene family in rice (*Oryza sativa* L.). *Molecular Biology Reports*, 36(2), 281–287. Available from: <https://doi.org/10.1007/s11033-007-9177-0>
- Yang, Z., Wu, Y., Li, Y., Ling, H.Q. & Chu, C. (2009) OsMT1a, a type 1 metallothionein, plays the pivotal role in zinc homeostasis and drought tolerance in rice. *Plant Molecular Biology*, 70(1–2), 219–229. Available from: <https://doi.org/10.1007/s11103-009-9466-1>
- Yokosho, K., Yamaji, N. & Ma, J.F. (2016) OsFRDL1 expressed in nodes is required for distribution of iron to grains in rice. *Journal of Experimental Botany*, 67(18), 5485–5494. Available from: <https://doi.org/10.1093/jxb/erw314>
- Yokosho, K., Yamaji, N., Ueno, D., Mitani, N. & Ma, J.F. (2009) OsFRDL1 is a citrate transporter required for efficient translocation of iron in rice. *Plant Physiology*, 149(1), 297–305. Available from: <https://doi.org/10.1104/pp.108.128132>
- Yoneyama, T., Ishikawa, S. & Fujimaki, S. (2015) Route and regulation of zinc, cadmium, and iron transport in rice plants (*Oryza sativa* L.) during vegetative growth and grain filling: metal transporters, metal speciation, grain Cd reduction and Zn and Fe biofortification. *International Journal of Molecular Sciences*, 16(8), 19111–19129. Available from: <https://doi.org/10.3390/ijms160819111>
- Yordem, B.K., Conte, S.S., Ma, J.F., Yokosho, K., Vasques, K.A., Gopalsamy, S.N. et al. (2011) Brachypodium distachyon as a new model system for understanding iron homeostasis in grasses: phylogenetic and expression analysis of Yellow Stripe-Like (YSL) transporters. *Annals of Botany*, 108(5), 821–833. Available from: <https://doi.org/10.1093/aob/mcr200>
- Yuan, L., Yang, S., Liu, B., Zhang, M. & Wu, K. (2012) Molecular characterization of a rice metal tolerance protein, OsMTP1. *Plant Cell Reports*, 31(1), 67–79. Available from: <https://doi.org/10.1007/s00299-011-1140-9>
- Zargar, S.M., Fujiwara, M., Inaba, S., Kobayashi, M., Kurata, R., Ogata, Y. et al. (2015) Correlation analysis of proteins responsive to Zn, Mn, or Fe deficiency in Arabidopsis roots based on iTRAQ analysis. *Plant Cell Reports*, 34(1), 157–166. Available from: <https://doi.org/10.1007/s00299-014-1696-2>
- Zhang, J., Wang, S., Song, S., Xu, F., Pan, Y. & Wang, H. (2019) Transcriptomic and proteomic analyses reveal new insight into chlorophyll synthesis and chloroplast structure of maize leaves under zinc deficiency stress. *Journal of Proteomics*, 199, 123–134. Available from: <https://doi.org/10.1016/j.jpro.2019.03.001>
- Zhang, M. & Liu, B. (2017) Identification of a rice metal tolerance protein OsMTP11 as a manganese transporter. *PLOS One*, 12, e0174987. Available from: <https://doi.org/10.1371/journal.pone.0174987>
- Zhang, T.Q., Xu, Z.G., Shang, G.D. & Wang, J.W. (2019) A single-cell RNA sequencing profiles the developmental landscape of arabidopsis root. *Molecular Plant*, 12(5), 648–660. Available from: <https://doi.org/10.1016/j.molp.2019.04.004>
- Zhang, Y., Xu, Y.H., Yi, H.Y. & Gong, J.M. (2012) Vacuolar membrane transporters OsVIT1 and OsVIT2 modulate iron translocation between flag leaves and seeds in rice. *Plant Journal*, 72(3), 400–410. Available from: <https://doi.org/10.1111/j.1365-3113.2012.05088.x>

How to cite this article: Amini, S., Arsova, B. & Hanikenne, M. (2022) The molecular basis of zinc homeostasis in cereals. *Plant, Cell & Environment*, 1–23. <https://doi.org/10.1111/pce.14257>