

EDITORIAL COMMENT

Concomitant Cardiac Amyloidosis in Severe Aortic Stenosis



The Trojan Horse?*

Philippe Pibarot, DVM, PhD,^a Patrizio Lancellotti, MD, PhD,^{b,c} Jagat Narula, MD^d

Per the *Aeneid* of Virgil, the Trojans were defeated after the Greeks left behind a huge wooden horse at the gates of Troy and pretended to sail back home after a 10-year siege. The wooden horse was filled with Greek warriors who invaded Troy after the Trojans brought the horse inside the city walls. Quite like the hidden warriors, cardiac amyloidosis (CA) is an insidious and often undetected process. Aortic stenosis (AS) and transthyretin (ATTR) amyloidosis are both more prevalent in elderly people, and may affect functional and clinical outcomes (1). However, there are, hitherto, few data pertaining to the exact prevalence and severity of CA and its impact on outcomes in patients with AS both before and after aortic valve replacement (AVR).

SEE PAGE 128

In this issue of the *Journal*, Nitsche et al. (2) report results of a multicenter international registry of the prevalence, predictors, and outcomes in 408 consecutive severe AS patients with AS evaluated for transcatheter aortic valve replacement (TAVR) at 3 referral centers. Bone scintigraphy with ^{99m}Tc-DPD

was performed in all and assessed by a core laboratory using the Perugini grading system (grade 0, negative scan, and grades 1 to 3 for increasingly positive scans). ATTR-CA was diagnosed by a positive DPD result, and the absence of a clonal immunoglobulin and light-chain CA by a tissue biopsy.

PREVALENCE, TYPE, AND SEVERITY OF CA IN SEVERE AS

In the study by Nitsche et al. (2), CA was detected in 48 (11.8%) patients, which is consistent with previous reports of prevalence between 9% and 16% (1,3-6). Of these 48 patients, one-third (n = 16, 3.9% of the total population) had grade 1 and 7.9% had grade 2 or 3 DPD uptake. In this elderly population (age 83.4 ± 6.5 years) with severe AS, ATTR-CA was confirmed in all patients except 1 with light-chain CA. Compared to patients with lone AS, those with AS and CA were older and had worse functional status, worse cardiac remodeling, higher circulating N-terminal pro-brain natriuretic peptide and troponin levels, and more frequently exhibited a low-flow, low-gradient AS pattern.

RED FLAGS FOR CONCOMITANT CA IN SEVERE AS

The study by Nitsche et al. (2) confirms that CA is frequent (1 out of 8 patients) in patients with severe AS referred for TAVR evaluation. However, systematic screening for CA using bone scintigraphy and light-chain analysis does not appear to be an optimal approach from logistic and economic standpoints. The challenge, in this context, is to differentiate a wooden horse (lone AS) from a Trojan horse (AS with CA). Hence, it is of paramount importance to develop and validate red flags for identification of AS patients

*Editorials published in the *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of JACC or the American College of Cardiology.

From the ^aInstitut Universitaire de Cardiologie et de Pneumologie de Québec/Québec Heart and Lung Institute, Université Laval, Québec City, Québec, Canada; ^bUniversity of Liège Hospital, GIGA Cardiovascular Sciences, Departments of Cardiology, Heart Valve Clinic, CHU Sart Tilman, Liège, Belgium; ^cGruppo Villa Maria Care and Research, Anthea Hospital, Bari, Italy; and the ^dMount Sinai Hospital Morningside, New York, New York, USA.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

who are likely to carry concomitant CA and who should therefore undergo confirmatory diagnostic testing. However, CA shares several clinical, electrocardiographic, and echocardiographic features with AS phenotype, including old age, left ventricular concentric hypertrophy, diastolic dysfunction, and depressed global longitudinal strain (1). Hence, the challenge in the AS population is to identify red flags that are more specific to CA. In the study by Nitsche et al. (2), a clinical score (i.e., the RAISE [remodeling, age, injury, system, and electrical] score, including several clinical, electrocardiographic, and echocardiographic parameters, and blood biomarkers) was developed to predict the presence of CA in patients with severe AS. This multiparameter score assessed age ≥ 85 years (1 point), history of carpal tunnel syndrome (3 points), presence of right bundle branch block (2 points), Sokolow/Lyon index < 1.9 mV (1 point), high sensitivity troponin level > 20 ng/ml (1 point), and E/A ratio > 1.4 (1 point), and a score ≥ 2 showed a high sensitivity (84%) and a score ≥ 3 had a high specificity (94%) for the detection of CA. Among the aforementioned 6 parameters, carpal tunnel syndrome and low electrocardiogram voltage despite the presence of left ventricular hypertrophy were more specific for co-existing CA. Old age, advanced diastolic dysfunction, and reduced longitudinal strain, even if associated with apical sparing, are not necessarily specific for co-existent CA and are frequently encountered in severe AS. Furthermore, the E/A ratio is not an optimal determinant of diastolic dysfunction. Despite limitations, this simple score provides reasonable accuracy to raise suspicion for CA, and a score ≥ 2 should prompt confirmatory tests including bone scintigraphy and light-chain analysis. The RAISE score allows room for further improvement by inclusion of other potentially valuable red flags, such as the presence of biventricular hypertrophy, tricuspid annulus S' velocity < 9 cm/s, or TAPSE (tricuspid annular plane systolic excursion) < 14 mm (Figure 1) (1,4). This score will need to be validated in a larger and independent population.


RESPONSE TO TREATMENT AND OUTCOMES IN PATIENTS WITH AS AND CONCOMITANT CA

The risk of futility of AVR has been a widely held concern in the management of AS patients with concomitant CA. Indeed, initial reports had suggested that such patients carry higher risk of futility of AVR and have worse outcomes compared to those with lone AS (1,5,6). These previous studies, however,

suffered from selection bias. The study by Nitsche et al. (2), which included a consecutive series of patients with systematic detection algorithm for CA, is reassuring in that patients with AS and CA had outcomes similar to those with lone AS. However, there was a trend ($p = 0.05$) for higher mortality at 1 year in AS-CA versus lone AS, but the former had worse risk profile at baseline, and multivariable analysis demonstrated no independently adverse effect of coexisting CA on outcomes both in patients undergoing TAVR and those treated medically. Assessment of treatment utility versus futility not only includes patients' vital status but also improvement in symptomatic and functional status and quality of life. The study by Nitsche et al. (2) only reported all-cause mortality obtained from national registries. Further studies are thus needed to assess the impact of CA on other relevant clinical outcomes, including rehospitalization for heart failure, functional class, and quality of life. Also, this study is limited to 3-year follow-up, whereas CA may have an impact on longer-term outcomes.

VALVULAR AMYLOIDOSIS AND THE LONGEVITY OF BIOPROSTHESIS

The Trojan Horse has come to mean any stratagem that causes a target to invite a foe into a protected location, and this analogy is not only consequential for the myocardium but also the valve (Figure 1). The amyloid protein may also deposit within the aortic valve leaflets and could therefore contribute to the valve leaflet thickening and the progression of AS (7,8). Aortic valve calcium scoring measured by non-contrast computed tomography (CT) may be useful to confirm stenosis severity in the context of low-flow, low-gradient AS, which is more prevalent in patients with AS-CA (Figure 1) (1). However, noncontrast CT only captures mineralized tissues and would underestimate AS severity in patients whose valve stenosis is predominantly caused by noncalcific tissues, such as fibrous tissues or even amyloid substance. New imaging methods based on contrast CT are currently being developed to quantitate the volume of both calcific and noncalcific valvular tissues. These methods could eventually become helpful to confirm severity of AS in patients with low-flow, low-gradient AS and CA. It is plausible that amyloid proteins may also infiltrate and accumulate within the bioprosthetic valve leaflets and contribute to accelerated structural deterioration of the prosthetic valve. Further studies are needed to determine whether AS-CA patients will have shorter valve durability following AVR.

FIGURE 1 Screening, Diagnosis, and Management of Cardiac Amyloidosis in Patients With Aortic Stenosis

Adapted with permission from Ternacle et al. (1). Artwork by Mia Pibarot. AL = light chain; AS = aortic stenosis; ATTR = transthyretin; AVA = aortic valve area; AVR = aortic valve replacement; CA = cardiac amyloidosis; ECV = extracellular volume; GLS = global longitudinal strain; HF = heart failure; IVS = interventricular septum; LGE = late gadolinium enhancement; LVH = left ventricular hypertrophy; MG = mean transvalvular gradient; MI = myocardial infarction; RVW = right ventricular wall; SLI = Sokolow/Lyon index; SVi = stroke volume index; TAPSE = tricuspid plane systolic excursion; TAVR = transcatheter aortic valve replacement; TTE = transthoracic echocardiography.

ALGORITHM FOR SCREENING, DIAGNOSIS, AND MANAGEMENT OF CA IN SEVERE AS

Due to a substantial prevalence of CA in severe AS, it is prudent to follow a systematic approach for the care of such patients (Figure 1). As the first step, it is important to recognize red flags for CA. If present, the second step is to perform additional tests including DPD scintigraphy, monoclonal light-chain analysis in blood and urine, and/or myocardial biopsy to confirm the diagnosis and type of CA. The third step should confirm the severity of AS by documenting either the echocardiographic evidence of high gradient, or severe aortic valve calcification on noncontrast CT in the case of low-flow, low-gradient AS. The last step should allow the selection of the type of treatment; patients with confirmed severe AS should promptly undergo AVR, and TAVR should be preferred in patients with concomitant CA. Indeed, patients with AS and CA often have advanced cardiac damage (9) with frequent impairment of the right heart including right ventricular dysfunction, tricuspid regurgitation, and/or pulmonary hypertension. In this vulnerable subset of patients, transfemoral TAVR is usually associated with better outcomes than surgical AVR (10,11). Finally, treatment with tafamidis should be considered in patients with confirmed ATTR-CA, and chemotherapy in those with light-chain CA, regardless of the AS severity or the decision to perform AVR (Figure 1) (1).

CONCLUSIONS

CA occurs in 1 out of 8 patients with severe AS evaluated for TAVR and is associated with worse risk profile at baseline but similar prognosis following TAVR when compared to lone AS. The RAISE score ≥ 2 raises the suspicion for CA and should prompt confirmatory tests. The diagnosis of CA in symptomatic patients with severe AS should not preclude or delay the consideration for TAVR. Further, it should trigger the consideration for a pharmaco-therapeutic approach to target CA (e.g., tafamidis in case of ATTR-CA). Like the Trojan Horse, CA is more dangerous if it is not recognized and not dealt with.

AUTHOR DISCLOSURES

Dr. Pibarot has received funding from Edwards Lifesciences for echocardiography core laboratory analyses in the field of transcatheter aortic valve replacement and from Medtronic for in vitro analyses, with no personal compensation. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

ADDRESS FOR CORRESPONDENCE: Dr. Philippe Pibarot, Institut Universitaire de Cardiologie et de Pneumologie de Québec, 2725 Chemin Sainte-Foy, Québec G1V-4G5, Canada. E-mail: philippe.pibarot@med.ulaval.ca.

REFERENCES

1. Ternacle J, Krapf L, Mohty D, et al. Aortic stenosis and cardiac amyloidosis: JACC review topic of the week. *J Am Coll Cardiol* 2019;74:2638-51.
2. Nitsche C, Scully PR, Patel KP, et al. Prevalence and outcomes of concomitant aortic stenosis and cardiac amyloidosis. *J Am Coll Cardiol* 2021;77:128-39.
3. Scully PR, Patel KP, Treibel TA, et al. Prevalence and outcome of dual aortic stenosis and cardiac amyloid pathology in patients referred for transcatheter aortic valve implantation. *Eur Heart J* 2020;41:2759-67.
4. Castañó A, Narotsky DL, Hamid N, et al. Unveiling transthyretin cardiac amyloidosis and its predictors among elderly patients with severe aortic stenosis undergoing transcatheter aortic valve replacement. *Eur Heart J* 2017;38:2879-87.
5. Cavalcante JL, Rijal S, Abdelkarim I, et al. Cardiac amyloidosis is prevalent in older patients with aortic stenosis and carries worse prognosis. *J Cardiovasc Magn Reson* 2017;19:98.
6. Treibel TA, Fontana M, Gilbertson JA, et al. Occult transthyretin cardiac amyloid in severe calcific aortic stenosis: prevalence and prognosis in patients undergoing surgical aortic valve replacement. *Circ Cardiovasc Imaging* 2016;9:e005066.
7. Kristen AV, Schnabel PA, Winter B, et al. High prevalence of amyloid in 150 surgically removed heart valves—a comparison of histological and clinical data reveals a correlation to atheroinflammatory conditions. *Cardiovasc Pathol* 2010;19:228-35.
8. Audet A, Côté N, Couture C, et al. Amyloid substance within stenotic aortic valves promotes mineralization. *Histopathology* 2012;61:610-9.
9. Tastet L, Tribouilloy C, Maréchaux S, et al. Staging cardiac damage in patients with asymptomatic aortic valve stenosis. *J Am Coll Cardiol* 2019;74:550-63.
10. Pibarot P, Salaun E, Dahou A, et al. Echocardiographic results of transcatheter versus surgical aortic valve replacement in low-risk patients: the PARTNER 3 trial. *Circulation* 2020;141:1527-37.
11. Cremer PC, Zhang Y, Alu M, et al. The incidence and prognostic implications of worsening right ventricular function after surgical or transcatheter aortic valve replacement: insights from PARTNER IIA. *Eur Heart J* 2018;39:2659-67.

KEY WORDS aortic stenosis, cardiac amyloidosis, TAVR