



Contents lists available at ScienceDirect

Schizophrenia Research

journal homepage: www.elsevier.com/locate/schres

Anatomical integrity within the inferior fronto-occipital fasciculus and semantic processing deficits in schizophrenia spectrum disorders

Werner Surbeck^{a,b,g,*,1}, Jürgen Hänggi^{c,1}, Felix Scholtes^{b,d}, Petra V. Viher^e, André Schmidt^f, Katharina Stegmayer^e, Erich Studerus^f, Udine E. Lang^f, Anita Riecher-Rössler^j, Werner Strik^e, Erich Seifritz^{g,h}, Stefan Borgwardt^{f,i,2}, Boris B. Quednow^{a,h,2}, Sebastian Walther^{e,2}

^a Experimental and Clinical Pharmacopsychology, Department of Psychiatry, Psychotherapy and Psychosomatics, Psychiatric Hospital of the University of Zurich, Zurich, Switzerland

^b Department of Neuroanatomy, Faculty of Medicine, University of Liège, Liège, Belgium

^c Division Neuropsychology, Department of Psychology, University of Zurich, Zurich, Switzerland

^d Department of Neurosurgery, University Hospital of Liège, Liège, Belgium

^e Translational Research Center, University Hospital of Psychiatry, University of Bern, Bern, Switzerland

^f Department of Psychiatry (UPK), University of Basel, Basel, Switzerland

^g Department of Psychiatry, Psychotherapy and Psychosomatics, Psychiatric Hospital of the University of Zurich, Zurich, Switzerland

^h Neuroscience Center Zurich, University and ETH, Zurich, Switzerland

ⁱ Department of Psychiatry and Psychotherapy, University of Lübeck, Germany

^j Medical Faculty, University of Basel, Basel, Switzerland

ARTICLE INFO

Article history:

Received 10 July 2019

Received in revised form 25 November 2019

Accepted 20 December 2019

Available online xxx

Keywords:

Schizophrenia

First episode psychosis

Inferior frontooccipital fasciculus

Diffusion tensor imaging

Formal thought disorder

Semantic impairment

ABSTRACT

The core symptoms of schizophrenia spectrum disorders (SSD) include abnormal semantic processing which may rely on the ventral language stream of the human brain. Thus, structural disruption of the ventral language stream may play an important role in semantic deficits observed in SSD patients.

Therefore, we compared white matter tract integrity in SSD patients and healthy controls using diffusion tensor imaging combined with probabilistic fiber tractography. For the ventral language stream, we assessed the inferior fronto-occipital fasciculus [IFOF], inferior longitudinal fasciculus, and uncinate fasciculus. The arcuate fasciculus and corticospinal tract were used as control tracts. In SSD patients, the relationship between semantic processing impairments and tract integrity was analyzed separately. Three-dimensional tract reconstructions were performed in 45/44 SSD patients/controls ("Bern sample") and replicated in an independent sample of 24/24 SSD patients/controls ("Basel sample").

Multivariate analyses of fractional anisotropy, mean, axial, and radial diffusivity of the left IFOF showed significant differences between SSD patients and controls ($p_{(FDR-corr)} < 0.001$, $\eta_p^2 = 0.23$) in the Bern sample. Axial diffusivity (AD) of the left UF was inversely correlated with semantic impairments ($r = -0.454$, $p_{(FDR-corr)} = 0.035$). In the Basel sample, significant group differences for the left IFOF were replicated ($p < .01$, $\eta_p^2 = 0.29$), while the correlation between AD of the left IFOF and semantic processing decline ($r = -0.376$, $p = .09$) showed a statistical trend. No significant effects were found for the dorsal language stream.

This is direct evidence for the importance of the integrity of the ventral language stream, in particular the left IFOF, in semantic processing deficits in SSD.

© 2020 Elsevier B.V. All rights reserved.

1. Introduction

Since the introduction of the modern conceptualization of schizophrenia, formal thought disorder (TD) has been considered to be a core symptom of the disease (Bleuler, 1911; Kraepelin, 1919). Clinically, this corresponds to disturbances in oral communication. Particularly positive TD, such as semantic violations, seem to be highly disease-specific for schizophrenia spectrum disorders (SSD) (Kircher et al., 2018; Levy et al., 2010; Morgan et al., 2017; Salavera et al., 2013). Semantic anomalies have thus been incorporated into scales designed to

* Corresponding author at: Department of Psychiatry, Psychotherapy and Psychosomatics, Psychiatric Hospital of the University of Zurich, Lenggstrasse 31, CH-8032 Zurich, Switzerland.

E-mail address: werner.surbeck@puk.zh.ch (W. Surbeck).

¹ Shared first authorship.

² Shared last authorship in alphabetical order.

assess TD, such as the Scale for the Assessment of Thought, Language and Communication (TLC) (Andreasen, 1979; Levy et al., 2010), and into general psychopathological scales devoted to psychotic syndromes, for example the Bern Psychopathology scale (BPS) (Strik et al., 2010).

Positive TD has been reported to be associated with structural alterations of the perisylvian language network of the brain, including the classic language areas of Wernicke and Broca (Fusar-Poli et al., 2012; Shepherd et al., 2012), as well as the middle longitudinal fasciculus (MLF) (Asami et al., 2013). However, the course and termination of the MLF in humans is still a matter of debate. Its role in language remains unclear (Champfleuer et al., 2012; Maldonado et al., 2013). Indeed, increasing evidence indicates that language processing mainly relies on a dual-stream architecture, analogous to that of the visual system (Hickok and Poeppel, 2007; Hickok and Poeppel, 2004; Saur et al., 2008; Ungerleider and Haxby, 1994): a dorsal phonological stream – including the arcuate fasciculus (AF) (Hickok and Poeppel, 2004) – and a ventral stream. The latter, linking the posteroinferior, occipitotemporal, part of the brain to the frontal lobe, depends on a direct pathway, the inferior fronto-occipital fasciculus (IFOF), and an indirect pathway, formed by the inferior longitudinal fasciculus (ILF) linked to the uncinate fasciculus (UF) (Sarubbo et al., 2016). Functional disruption of the IFOF may well participate in verbal semantic processing disturbances: inhibitory electrical stimulation of the IFOF during awake surgery induces semantic disturbances (Duffau et al., 2014) and reduced integrity of the left IFOF was found to correlate with subjects' semantic performance in healthy elderly (de Zubicaray et al., 2011) as well as brain-injured individuals (Han et al., 2013). This may also apply to TD within SSD. We therefore investigated the relationship between verbal semantic processing impairments and the integrity of the white matter (WM) integrity underlying language function in patients with schizophrenia and first episode psychosis (FEP) in two independent studies using diffusion tensor imaging (DTI). We hypothesized IFOF integrity is altered in patients with SSD and that semantic anomalies are correlated with these alterations.

2. Patients and methods

We retrieved preexisting data sets from the University Hospital of Psychiatry, Bern, Switzerland, and the Department of Psychiatry of the University of Basel, Switzerland. Detailed clinical and demographic characteristics of both SSD samples have been reported elsewhere (Schmidt et al., 2017; Schmidt et al., 2015; Stegmayer et al., 2017; Walther et al., 2017; Walther et al., 2015). Behavioral, perfusion and resting-state functional connectivity findings of the Bern sample (Schmidt et al., 2017, Schmidt et al., 2015; Stegmayer et al., 2017; Walther et al., 2017, Walther et al., 2015) and structural connectivity findings other than reported in the present manuscript of the Basel sample (Schmidt et al., 2017, Schmidt et al., 2015; Stegmayer et al., 2017; Walther et al., 2017, Walther et al., 2015) can be found in the referenced publications.

2.1. Participants

2.1.1. Patients and control subjects of the Bern sample

Thirty male and 15 female (total 45) schizophrenia patients and 26 male and 18 female (total 44) healthy control subjects matched for age and education were included in this study. Subjects were recruited from the inpatient and outpatient departments of the University Hospital of Psychiatry, Bern, Switzerland. Healthy controls were recruited among staff and via advertisements. All subjects were right-handed as determined by the Edinburgh handedness inventory (Oldfield, 1971). Exclusion criteria for both groups included current substance abuse or dependence other than nicotine; past or current medical or neurological conditions impairing movements, such as dystonia, idiopathic parkinsonism, and stroke; history of head trauma with concurrent loss of consciousness, and a history of electroconvulsive treatment. Exclusion

criteria limited to control subjects were a history of any psychiatric disorder and any first-degree relatives with schizophrenia or schizoaffective disorder. All participants were interviewed with the Mini International Neuropsychiatric Interview (Sheehan et al., 1998) and the Comprehensive Assessment of Symptoms and History (Andreasen et al., 1992). Diagnoses were given according to Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition criteria. All but 4 patients were under antipsychotic pharmacotherapy, and dosages were computed as chlorpromazine (CPZ) equivalents (Woods, 2003). Further assessment of schizophrenia psychopathology included the Positive and Negative Syndrome Scale (PANSS) (Kay et al., 1987). Demographic, global brain and clinical characteristics are given in Table 1. The study was approved by the local ethics committee and all participants provided written informed consent.

2.1.2. Patients and control subjects of the Basel sample

Twenty-four patients with FEP and 24 healthy control subjects matched for age were included in this study. Subjects were recruited from the specialized outpatient clinic for early detection of psychosis at the Department of Psychiatry, University of Basel. As reported in Supplementary Table 3, most participants were right handers ($\chi^2_{(df=1)} = 0.76, p = .67$). Exclusion criteria for both groups included current substance abuse according to ICD-10 research criteria, history of previous psychotic disorder; psychotic symptomatology secondary to an organic disorder; psychotic symptomatology associated with an affective psychosis or a borderline personality disorder; age under 18 years; inadequate knowledge of the German language; and verbal IQ <70, measured with the German vocabulary test “Mehrfachwahl Wortschatz-Test Form B” (MWT-B) (Lehrl et al., 1995). Patients were further interviewed with the Basel Screening Instrument for Psychosis (BSIP) (Riecher-Rössler et al., 2008) and the Brief Psychiatric Rating Scale (BPRS) (Lukoff et al., 1986). All met the transition criteria of Young et al. (Yung et al., 1998). Only 9 patients received antipsychotics whose dosages were computed as CPZ equivalents (Woods, 2003). The study was also approved by the local ethics committee and all participants provided written informed consent.

2.2. Assessment of verbal semantic performance

2.2.1. Bern sample

Clinical data with respect to verbal semantic performance was retrieved from the following inventories (Table 1):

- BPS is a validated clinical rating scale developed to group psychotic symptoms in the domains of language, affectivity, and motor-behavior (Strik et al., 2010). The items *mistaking identity, coherence of speech, interruptions (including neologism), naming and apprehension of meaning* were selected for the assessment of semantic processing.
- The TLC is a widely-used instrument for the assessment of language dysfunction (Goldberg et al., 1998). The items *semantic paraphasia, neologisms, word approximations, derailment, and incoherence* were selected for the assessment of semantic processing.

For the correlation of semantic impairments with measures of tract integrity, ten scales considered most representative for semantic processing (Levy et al., 2010) were selected (five from the BPS and five from TLC) and the values of these scales added up.

2.2.2. Basel sample

In the Basel sample, the available clinical tests were different from those of the Bern sample. Data relevant for verbal semantic performance were thus retrieved from the California verbal learning test (CVLT) (Woods et al., 2006) (see Supplementary Table 3). We used the List-Based Semantic Clustering Index of the CVLT (Stricker et al., 2002).

Table 1

Demographic, global brain and clinical characteristics of the schizophrenia patients and healthy control subjects of the Berne sample.

Measures	Schizophrenia (n = 45)				Control subjects (n = 44)				t-Value	df	p
	Mean	SD	Min.	Max.	Mean	SD	Min.	Max.			
Age (years)	38.1	11.5	19.0	63.0	38.8	13.6	18.0	64.0	0.26	84.1	0.79
Education (years)	13.5	3.1	8.0	21.0	14.1	2.7	8.0	20.0	1.10	87.0	0.27
Duration of illness (years)	12.0	12.4	0.0	45.1	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Number of episodes	6.4	7.1	1.0	30.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Total intracranial volume (liter)	1.553	0.159	1.212	1.903	1.550	0.149	1.307	1.907	-0.11	87.0	0.91
Total gray matter volume (liter)	0.619	0.081	0.424	0.806	0.639	0.057	0.548	0.788	1.33	79.2	0.19
Total white matter volume (liter)	0.453	0.057	0.307	0.584	0.460	0.060	0.347	0.590	0.59	87.0	0.56
Average fractional anisotropy (0–1)	0.275	0.013	0.248	0.309	0.278	0.010	0.261	0.301	1.52	82.0	0.13
Average axial diffusivity (mm ² /s)	0.0012	0.00007	0.0011	0.0014	0.0012	0.00005	0.0011	0.0013	-1.95	78.4	0.054
Average radial diffusivity (mm ² /s)	0.0008	0.00006	0.0007	0.0010	0.0008	0.00004	0.0007	0.0009	-2.72	74.3	0.008
Average mean diffusivity (mm ² /s)	0.0010	0.00007	0.0009	0.0011	0.0009	0.00005	0.0008	0.0010	-2.07	78.7	0.042
Average motion translation (mm)	1.523	0.359	0.938	2.382	1.513	0.335	0.793	2.250	-0.13	87.0	0.90
Average motion rotation (degree)	0.007	0.0023	0.003	0.013	0.006	0.0023	0.002	0.012	-1.55	87.0	0.12
Percent bad slices	0.022	0.0565	0.0	0.29	0.0043	0.0222	0.00	0.14	-1.94	57.5	0.058
Average dropout scores	1.032	0.091	1.00	1.44	1.011	0.058	1.00	1.37	-1.30	75.2	0.20
Bern Psychopathology Scale ^a	4.09	3.43	0.0	13.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Thought, Language, and Communication	6.16	7.25	0.0	29.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A
PANSS (sum of positive symptoms)	18.3	6.41	7.0	33.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A
PANSS (sum of negative symptoms)	18.3	4.94	10.0	35.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A
PANSS (sum of total symptoms)	72.8	17.4	40.0	112.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A
	Frequency				Frequency				χ²	df	p
Sex (male/female)	30/15				26/18				0.55	1	0.52

All measures were compared between groups using student's t-tests for independent samples. Abbreviations: df, degrees of freedom; Max., maximum; Min., minimum; N/A, not available; p, p-value; PANSS, positive and negative symptom scale; SD, standard deviation.

^a Negative values (-1) were converted to positive values (1) to compute the sum of each patient.

2.3. Image acquisition

At both sites, imaging was performed on a 3 T MRI scanner (Bern: Siemens Magnetom Trio; Basel: Magnetom Verio, both Siemens Healthcare, Erlangen, Germany) with a 12-channel head coil for signal reception. 3D-T1-weighted images as well as DTI measurements were acquired. Technical details can be found in the Supplementary Methods.

2.4. Image processing

First, T1-weighted images were processed using the FreeSurfer software (<http://surfer.nmr.mgh.harvard.edu/fswiki>) (Dale et al., 1999; Fischl et al., 2004; Fischl et al., 2002). We used this tool to obtain the seed points needed for fiber tractography. The processing steps are described elsewhere (<http://surfer.nmr.mgh.harvard.edu/fswiki/FreeSurferMethodsCitation>). DTI data were then processed with the tracts constrained by underlying anatomy (TRACULA) toolbox (<https://surfer.nmr.mgh.harvard.edu/fswiki/Tracula>).

The TRACULA toolbox, which relies on a global probabilistic tractography approach with anatomical priors, was used for the automated reconstruction of a set of 18 major WM pathways from DTI images. Prior distributions on the neighboring anatomical structures of each pathway are derived from an atlas and combined with the FreeSurfer cortical parcellation scheme and subcortical segmentation to constrain the tractography solutions (Yendiki et al., 2014; Yendiki et al., 2011). In the present work, we focused on the specific set of the WM fiber tracts of interest named above, previously linked to semantic information processing in healthy subjects, i.e., the IFOF, UF and ILF (Friederici, 2009; Saur et al., 2008) (Fig. 1). For two control tracts, no effects were expected: we chose the arcuate fasciculus (AF) as a language-related tract which seems not to be involved in semantic processing (Friederici, 2009), as well as the corticospinal tract (CST) responsible for motor control of the extremities and no language function (Brodal and Walberg, 1982; Zarei et al., 2007). The probabilistic tractography is described in detail in the Supplementary Methods and the reconstructed tracts of interest are shown in Fig. 1. For all tracts, fractional anisotropy (FA), axial diffusivity (AD), radial diffusivity (RD), and mean

diffusivity (MD) values were extracted. Further details with respect to the TRACULA tractography toolbox and the output measures it provides can be found in the online documentation (<https://surfer.nmr.mgh.harvard.edu/fswiki/Tracula>) as well as in the relevant publications (Yendiki et al., 2016; Yendiki et al., 2014; Yendiki et al., 2011).

2.5. Statistical analysis

Comparisons of demographic and global brain measures as well as motion parameters across groups were performed using student's t-tests for independent samples and χ^2 -Tests for gender and handedness. DTI-related measures (FA, AD, RD and MD) of the IFOF, ILF, UF as well as of the AF and CST (control tracts) were compared between groups using parametric univariate and multivariate analysis of covariance (ANCOVA and MANCOVA) models controlling for global diffusivity values. Because of the interdependency of the various DTI-derived measures, multivariate measures are more appropriate than univariate ones. We also tested whether the residuals after regressing out the global diffusivity measures were normally distributed using the Kolmogorov-Smirnov normality test. Almost all residuals were approximately normally distributed (Supplementary Table 1 and 2).

The associations between the DTI-related measures and semantic impairments were conducted using Spearman's rank order correlations based on residuals after regressing out global DTI-derived measures, age due to the wide age range (19–63 years) as well as gender, education, PANSS scores, and CPZ equivalents of the schizophrenia patients in the discovery (Bern) sample. Due to the smaller age range (18–41 years) in the FEP patients of the replication (Basel) sample, age was not regressed out. We also decided to not correct for gender, education, PANSS scores, and CPZ equivalents in the replication (Basel) sample because the sample size is very small (only 14 patients) so that adding additional covariates of no interest would reduce the number of degrees of freedom considerably and hence increasing the risk of false negative results.

Effect sizes are reported based on the correlation coefficient r (correlations) and η_p^2 (ANCOVA and MANCOVA). According to Cohen, effect sizes are denoted small if $r = 0.10$ and $\eta_p^2 = 0.01$, moderate if $r =$




Fig. 1. White matter pathways investigated in the present study. DTI data processed with TRACULA reveals the three major white matter pathways of the ventral language processing stream, i.e., the inferior fronto-occipital fasciculus (IFOF, in red), inferior longitudinal fasciculus (ILF, in blue) and the uncinate fasciculus (UF, in green). The arcuate fasciculus and corticospinal tract (not shown) served as controls. A. right lateral view; B. left lateral view; C. inferior view; D. superior view; E. anterior view; F. posterior view. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

0.30 and $\eta_p^2 = 0.06$, and large if $r = 0.50$ and $\eta_p^2 = 0.14$. All tests were performed using IBM SPSS Statistics software version 25 (www.ibm.com/SPSS/Statistics). Error probability was set at $p < .05$ and, if not otherwise stated, two-tailed hypothesis testing has been performed. To account for the multiple comparisons performed, we applied the false discovery rate (FDR) correction. FDR correction was applied to the discovery (Bern) sample but not to the replication (Basel) sample. Given that results in the Basel data set are similar as those found with the Bern data set, the analyses of which are FDR corrected, the probability that the findings in the Basel data set represent false positives is very low, even if we do not correct the analyses of the Basel data set for multiple comparisons (Miller, 2009). For the group comparisons, we FDR corrected for the semantic-related tracts of interests (IFOF, ILF and UF) as well as for the language- and not semantic-related control tract (AF), but we did not FDR correct for the motor-related control tract (CST). Correlations have been performed only for the IFOF because the IFOF was the only tract for which we found a significant group difference with our MANCOVA models in both samples. In total, we FDR corrected for 52 statistical models, i.e., for the 32 ANCOVAs (4 measures \times 4 tracts \times 2 hemispheres), 12 MANCOVAs (4 tracts \times left/right/both) as well as for the 8 Spearman correlations with semantic

impairments (4 measures \times 1 tract \times 2 hemispheres). For the correlations in the replication sample, we used one-tailed hypothesis testing because of the anticipated direction of this association.

3. Results

3.1. Demographic and global brain measures

3.1.1. Bern sample

There were no significant differences between schizophrenia patients and healthy controls with respect to sex, age, education, total intracranial volume, total gray matter volume, and total WM volume (t -tests for independent samples, all $p > .19$, Table 1). The BPS and TLC scores are reported in Table 1.

DTI-derived global measures were significantly different between schizophrenia patients and healthy controls (Table 1 and Supplementary Results). Consequently, we used these global DTI measures as covariates-of-no-interest in our statistical models performed to investigate the integrity of the WM fiber pathways (see below).

With respect to head motion, no significant differences were found, neither for the translations and rotations nor for the dropout scores (all

$p > .12$) and the trend ($p = .058$) in percent slices with poor resolution/image quality (slightly increased in patients) is neglectable due to the very small number of inadequate slices (<1% in both groups; Table 1). Note that most of the participants (79 out of 89, 88%) have 0% bad slices rendering this variable not suitable for regression analysis and therefore we refrained from using percent bad slices as a covariate of no interest in our statistical models.

3.1.2. Basel sample

Patients with FEP and healthy controls did not differ in sex, handedness, age, total intracranial volume, total gray matter volume, and total white matter volume (all $p > .15$, Supplementary Table 3). However, there was one statistically significant group difference regarding years of education that was higher in control subjects (Supplementary Table 3).

DTI-derived global measures were significantly different between groups. We therefore used these parameters as covariates-of-no-interest in our statistical models performed to investigate the integrity of the white matter fiber pathways.

With respect to head motion, no significant differences were found, neither for the translations, nor rotations, nor dropout scores, nor the percentage of inadequate slices (all $p > .14$).

3.2. Group comparisons of WM fiber pathways

3.2.1. Bern sample

The reconstructed WM fiber pathways of interest (IFOF, ILF and UF) are shown in Fig. 1 and the IFOF diffusivity values of the Bern sample (and that of the Basel sample as well) are listed in Supplementary Table 4.

3.2.1.1. Inferior fronto-occipital fasciculus. If considering the interdependency between the four diffusivity measures of the IFOF and applying the more appropriate MANCOVA model while correcting for the four global diffusivity measures, a robust and statistically significant difference between schizophrenia patients and healthy controls was found for the left and right IFOF together ($F_{(df=8,76)} = 5.18$, $p_{(FDR-corrected)} = 0.002$, $\eta_p^2 = 0.35$), as well as for the left IFOF ($F_{(df=4,80)} = 5.87$, $p_{(FDR-corrected)} = 0.009$, $\eta_p^2 = 0.23$), and for the right IFOF separately ($F_{(df=4,80)} = 2.78$, $p = .032$, $\eta_p^2 = 0.12$). The effect of the right IFOF lost statistical significance after FDR correction for multiple comparisons ($p_{(FDR-corrected)} = 0.19$). These effects were driven by decreased FA and increased AD, RD, and MD in patients compared to controls.

There were no significant group differences in any of the diffusivity measures (FA, AD, RD, and MD) when each measure was investigated in isolation while correcting for the corresponding global measure (ANCOVA).

3.2.1.2. Inferior longitudinal fasciculus, uncinata fasciculus, arcuate fasciculus and corticospinal tract. If all measures were analyzed simultaneously (MANCOVA), no significant group differences were found for almost all fiber bundles, except for the right AF ($F_{(df=6,78)} = 3.11$, $p_{(FDR-corrected)} = 0.041$, $\eta_p^2 = 0.19$).

When focusing on the four diffusivity measures of the tract center of the ILF, UF, AF, and CST, no significant group differences were found when analyzing each measure in isolation (ANCOVA), except for the right AF where patients showed increased AD compared to controls ($F_{(df=1,86)} = 6.09$, $p < .05$, $\eta_p^2 = 0.066$). Note that this effect is relatively weak compared with the ones reported for the IFOF, does not survive FDR correction ($p_{(FDR-corrected)} = 0.14$), and it is restricted to the right non-language dominant hemisphere.

3.2.2. Basel sample

3.2.2.1. Inferior fronto-occipital fasciculus. If considering the interdependency between the four diffusivity measures of the IFOF and applying

a MANCOVA model while correcting for the four global diffusivity measures, the same pattern of diffusivity differences as reported for the Bern sample emerged, except for the right IFOF: a strong and statistically significant difference between patients and healthy controls was found for the left and right IFOF together ($F_{(df=8,35)} = 2.68$, $p = .021$, $\eta_p^2 = 0.38$), for the left IFOF ($F_{(df=4,39)} = 3.89$, $p = .009$, $\eta_p^2 = 0.29$), but there was no significant group difference for the right IFOF ($F_{(df=4,39)} = 1.00$, $p = .42$, $\eta_p^2 = 0.093$).

In contrast to the Bern sample where univariate analyses did not reveal any significant effect of the four diffusivity measures, the left IFOF of the FEP patients of the Basel sample showed reduced FA ($F_{(df=1,45)} = 6.12$, $p = .017$, $\eta_p^2 = 0.120$), increased RD ($F_{(df=1,45)} = 6.30$, $p = .016$, $\eta_p^2 = 0.123$), and increased MD ($F_{(df=1,45)} = 4.77$, $p = .034$, $\eta_p^2 = 0.096$), whereas no effects were found for the right IFOF (FA: $F_{(df=1,45)} = 0.55$, $p = .46$, $\eta_p^2 = 0.012$; RD: $F_{(df=1,45)} = 0.35$, $p = .56$, $\eta_p^2 = 0.008$; MD: $F_{(df=1,45)} = 0.20$, $p = .66$, $\eta_p^2 = 0.004$). AD did not significantly differ for both IFOFs when each measure was investigated in isolation while correcting for the corresponding global measure (ANCOVA).

3.2.2.2. Inferior longitudinal fasciculus, uncinata fasciculus, arcuate fasciculus, and corticospinal tract. If all measures were analyzed simultaneously (MANCOVA), statistically significant differences between FEP patients and healthy controls were found for the left and right UF together ($F_{(df=6,37)} = 2.79$, $p = .024$, $\eta_p^2 = 0.31$), for the left UF ($F_{(df=3,40)} = 4.76$, $p = .006$, $\eta_p^2 = 0.26$), as well as the right ILF ($F_{(df=3,40)} = 3.75$, $p = .018$, $\eta_p^2 = 0.22$). For all other tracts, there were no statistically significant multivariate differences between the groups.

If focusing on the four diffusivity measures FA, AD, RD and MD of the tract center of the ILF, UF, AF, and CST no significant group differences were found when each measure was analyzed in isolation (ANCOVA).

3.3. Association of tract integrity with impairments in semantic processing

3.3.1. Bern sample

Semantic data were available for patients only. There were significant inverse correlations between semantic impairments and the diffusivity values (after regressing out the corresponding global diffusivity measure as well as age, sex, education, PANSS scores, and CPZ equivalents) of the left as well as right IFOF (see Table 2).

The strongest inverse correlation was found between semantic impairments and AD of the left IFOF ($r = -0.454$, $p_{(FDR-corrected)} = 0.002$), whereas semantic impairments and AD of the right IFOF were less strongly associated ($r = -0.331$, $p = .026$) and did not survive correction for multiple comparisons ($p_{(FDR-corrected)} = 0.188$) (scatterplots are shown in Fig. 2). Beyond AD, MD of the left IFOF correlated also inversely with semantic impairments ($r = -0.326$, $p = .029$). Note that the correlations of semantic impairments with AD of the right IFOF and that of MD of the left IFOF would not survive the FDR correction for multiple comparisons ($p_{(FDR-corrected)} = 0.188$ and $p_{(FDR-corrected)} = 0.187$, respectively). Moreover, the explained variance of the left IFOF ($r^2 = 0.206$) is about twice in magnitude than that of the right IFOF ($r^2 = 0.110$).

3.3.2. Basel sample

In order to correlate semantic impairments with measures of tract integrity, we selected the LBC of the CVLT. We report rank-ordered correlations (Spearman's rho) to safeguard against potential outliers in the semantic scores. These correlations are based on the residuals after regressing out the corresponding global diffusivity measure only. There was a significant correlation between the LBC and axial diffusivity in the left IFOF in the total sample of patients and controls (14 FEP patients and 18 controls, $r = -0.317$, $p < .05$, see Supplementary Table 5), which was reduced to a statistical trend in the patient group only (14 FEP patients, $r = -0.376$, $p = .0925$). Of note, the latter correlation is based on 14 patients only due to missing CVLT data in 10

Table 2
Correlations between semantic processing impairments and measures of tract integrity within the schizophrenia patients of the Berne sample.

		Fractional anisotropy	Axial diffusivity	Radial diffusivity	Mean diffusivity
IFOF left	Spearman's rho	-0.048	-0.454	-0.214	-0.326
	p-Value (FDR-corr.)	0.935	0.035	0.519	0.187
IFOF right	Spearman's rho	-0.054	-0.331	-0.196	-0.281
	p-Value (FDR-corr.)	0.935	0.188	0.569	0.288

Partial Spearman rank-ordered correlations corrected for the effects of global diffusivity measures, age, sex, education, PANSS scores, and CPZ equivalents. Significant correlations are printed in bold. Abbreviations: FDR-corr., false discovery rate corrected; CPZ, chlorpromazine; IFOF, inferior fronto-occipital fasciculus; PANSS, positive and negative symptom scale.

patients. However, again, these correlations indicate greater semantic impairments associated with reduced integrity of the left IFOF (scatterplots see in Supplementary Fig. 1).

4. Discussion

Compared to control subjects, schizophrenia and FEP patients in this investigation appear to have significantly structural alterations of the white matter tracts underlying semantic processing in the ventral language stream of the brain. While these structural, anatomical alterations

have been previously described in FEP (Cheung et al., 2008; Epstein and Kumra, 2015; Lee et al., 2013; Lu et al., 2011; Melicher et al., 2015; Szeszko et al., 2008) and chronic schizophrenia (Boos et al., 2013; Schneiderman et al., 2009; Skelly et al., 2008), little was known about a potential functional correlate. Furthermore, a current study on tract specific age effects reported an abnormal maturation and accelerated aging for the IFOF (Cetin-Karayumak et al., 2019) as compared to controls. Again, whether these age-related group differences were associated with specific functional decline remains unanswered. Only recently, Viher et al. (2018) reported a linear brain-behavior




Fig. 2. Associations between semantic processing impairments and axial diffusivity of the inferior fronto-occipital fasciculus (IFOF). Scatterplots of the correlations between semantic processing impairments and the axial diffusivity standardized residuals (after regressing out the effects of global axial diffusivity, age, sex, education, PANSS score, and CPZ equivalents) of the left IFOF ($r = -0.454$, $p_{\text{FDR-corrected}} = 0.035$, upper panel) and the right IFOF ($r = -0.331$, $p_{\text{FDR-corrected}} = 0.188$, lower panel). Abbreviations: CPZ, chlorpromazine; IFOF, inferior fronto-occipital fasciculus; PANSS, positive and negative symptom scale.

association between WM abnormalities in a predominantly left fronto-temporal language network and TD in schizophrenia. In that investigation, however, all fiber tracts of the ventral language stream (UF, ILF and IFOF) were combined. That investigation indicated that lower total FA was associated with increasing vulnerability to positive TD. Furthermore, Kubicki et al. (2011) reported a correlation between semantic processing, as retrieved from several tests of language and verbal memory, and the ventral connectivity between the inferior frontal gyrus and the superior temporal gyrus in schizophrenia patients, also suggesting an implication of the ventral language stream in semantic processing problems in schizophrenia.

In the present investigation, AD changes within both the IFOFs' were negatively correlated with semantic processing impairment in schizophrenia patients ("Bern sample"). Although bilaterally significant, the effect sizes of the left IFOF are about twice as strong as the effect sizes in the right IFOF, both for group comparison and the correlation with semantic impairments. This correlation is in line with previous reports on similar anatomofunctional correlations within language domains (Viher et al., 2018). This suggests that structural disintegration of white matter pathways can indeed be interpreted as representative of the corresponding functional decline. Severity of schizophrenia psychopathology, however, as represented by the PANSS scores, does not correlate statistically significantly with AD, neither for the left nor for the right IFOF. With a trend towards statistical significance, the observed negative correlation between anatomical integrity and semantic performance was reproduced in our replication sample ("Basel sample"). In addition, in the latter, when anatomofunctional correlations are computed across FEP patients and healthy controls, the correlation between semantic processing impairment and AD changes becomes significant for the left (but not the right) IFOF. A left side's predominance of the IFOF in verbal semantic processing is generally supported by data from intraoperative direct electrostimulation and semantic performance in healthy elderly individuals (Duffau et al., 2014). Compared to healthy controls FEP patients' in this investigation exhibit significant structural alterations in the left, but not the right IFOF. With regard to schizophrenia patients, the same difference emerged for both the left and the right IFOF. This observation might represent a compensatory mechanism seen in FEP and lost in chronic SSD. On the other hand, progressive bilateral structural alteration related to verbal semantic processing, exhibiting the same left-sided predominance, could also explain these findings.

The present study indicates that the involvement of the IFOF with respect to semantic processing is selective in patients suffering from confirmed psychotic illness: There are no significant correlations for the AF in neither the Bern nor the Basel sample. Concerning the ILF we found a negative correlation for the FA on the right in the Bern sample, which lose significance after FDR correction. With regard to the Basel sample there is a correlation between semantic processing decline and AD of the left and right UF. However, when anatomofunctional correlations are computed across FEP patients and healthy controls, the correlation remains significant only for the right non-language dominant UF.

The isolated correlation of semantic decline and anomalies in AD, not reflected by other DTI measures, might appear surprising. While the majority of DTI studies in schizophrenia have shown decreased FA in long-range association tracts, alterations in the radial or axial diffusivity have attracted less attention until now (Koch et al., 2013). Isolated AD abnormalities have been interpreted as a preliminary stage before more severe changes (Qiu et al., 2008; Zivadinov et al., 2013) and were attributed among others to possible changes in axonal number, size or compactness (Harsan et al., 2006; Schonberger et al., 2006; Song et al., 2003). Such changes in AD, without FA anomalies have been observed within the ventral and dorsal language systems of non-psychotic high risk patients—with high risk subjects demonstrating a deviation from the normal maturation trajectory (Kubicki et al., 2013). These changes might indeed precede further pathway pathologies (including myelin changes) (Kubicki et al., 2013). Nevertheless,

neuroanatomical correlates of AD changes are still under investigation and conflicting results are reported for AD changes (Zikou et al., 2016).

The retrospective nature and the relatively small size of the control (Basel) sample limit the interpretation of the present investigation. In order to gain more precise insight into the anatomical correlates of semantic processing alterations in SSD, broader prospective data are needed. Finally, in the scatterplots of Fig. 2, the amount of "zeros" is explained by the absence of clinically apparent semantic impairments in more than half of the patients (23 out of 45) in the Bern sample. Nevertheless, (1) in the Basel sample, where no such "zeros" occurred, a similar association has been found; (2) most schizophrenia patients with no semantic impairments showed positive residual AD values in the IFOF, whereas the patients with semantic impairments showed negative residual AD values in the IFOF, (3) even if statistical significance is lost due to lack of power, the direction of the correlation remains the same when excluding the 23 patients with no semantic impairments, and (4) when using the mean of all BPS and TCS scales instead of the mean of only the semantic related scales, there is still an inverse relationship although almost all "zeros" disappeared.

5. Conclusion

Structural alterations of the IFOF correlate with patients' semantic processing impairments in schizophrenia spectrum disorders. This is the first direct evidence for a contribution of disrupted IFOF integrity to semantic processing deficits in these patients.

Acknowledgments

This study received funding from the Swiss National Science Foundation (grant #152619 to Sebastian Walther) and the Bangerter-Rhyner-Foundation (to Sebastian Walther).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.schres.2019.12.025>.

References

- Andreasen, N.C., 1979. Thought, language, and communication disorders. I. Clinical assessment, definition of terms, and evaluation of their reliability. *Arch. Gen. Psychiatry* 36, 1315–1321.
- Andreasen, N.C., Flaum, M., Arndt, S., 1992. The Comprehensive Assessment of Symptoms and History (CASH). An instrument for assessing diagnosis and psychopathology. *Arch. Gen. Psychiatry* 49, 615–623.
- Asami, T., Saito, Y., Whitford, T.J., Makris, N., Niznikiewicz, M., McCarley, R.W., Shenton, M.E., Kubicki, M., 2013. Abnormalities of middle longitudinal fascicle and disorganization in patients with schizophrenia. *Schizophr. Res.* 143, 253–259.
- Bleuler, E., 1911. *Dementia Praecox or the Group of Schizophrenias*. International Universities Press, New York.
- Boos, H.B.M., Mandl, R.C.W., van Haren, N.E.M., Cahn, W., van Baal, G.C.M., Kahn, R.S., Hulshoff Pol, H.E., 2013. Tract-based diffusion tensor imaging in patients with schizophrenia and their non-psychotic siblings. *Eur. Neuropsychopharmacol.* 23, 295–304.
- Brodal, A., Walberg, F., 1982. A re-evaluation of the question of ascending fibers in the pyramidal tract. *Brain Res.* 232, 271–281.
- Cetin-Karayumak, S., Di Biase, M.A., Chunga, N., Reid, B., Somes, N., Lyall, A.E., Kelly, S., Solgun, B., Pasternak, O., Vangel, M., Pearson, G., Tamminga, C., Sweeney, J.A., Clementz, B., Schretlen, D., Viher, P.V., Stegmayer, K., Walther, S., Lee, J., Crow, T., James, A., Voineskos, A., Buchanan, R.W., Szeszko, P.R., Malhotra, A.K., Hegde, R., McCarley, R., Keshavan, M., Shenton, M., Rathi, Y., Kubicki, M., 2019. White matter abnormalities across the lifespan of schizophrenia: a harmonized multi-site diffusion MRI study. *Mol. Psychiatry* <https://doi.org/10.1038/s41380-019-0509-y> [Epub ahead of print].
- Champfleury, N.M.D., Maldonado, I.L., Moritz-gasser, S., Machi, P., Bars, E.L., Bonafé, A., Duffau, H., 2012. Middle longitudinal fasciculus delineation within language pathways: a diffusion tensor imaging study in human. *Eur. J. Radiol.* 82 (1), 151–157.
- Cheung, V., Cheung, C., McAlonan, G.M., Deng, Y., Wong, J.G., Yip, L., Tai, K.S., Khong, P.L., Sham, P., Chua, S.E., 2008. A diffusion tensor imaging study of structural dysconnectivity in never-medicated, first-episode schizophrenia. *Psychol. Med.* 38, 877–885.
- Dale, A.M., Fischl, B., Sereno, M.I., 1999. Cortical Surface-Based Analysis I. Segmentation and surface reconstruction. *Neuroimage* 9, 179–194.
- Duffau, H., Moritz-Gasser, S., Mandonnet, E., 2014. A re-examination of neural basis of language processing: proposal of a dynamic hodotopical model from data provided by brain stimulation mapping during picture naming. *Brain Lang.* 131, 1–10.

- Epstein, K.A., Kumra, S., 2015. White matter fractional anisotropy over two time points in early onset schizophrenia and adolescent cannabis use disorder: a naturalistic diffusion tensor imaging study. *Psychiatry Res.* 232, 34–41.
- Fischl, B., Salat, D.H., Busa, E., Albert, M., Dieterich, M., Haseilgrove, C., van der Kouwe, A., Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., Dale, A.M., 2002. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* 33, 341–355.
- Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D.H., Busa, E., Seidman, L.J., Goldstein, J., Kennedy, D., Caviness, V., Makris, N., Rosen, B., Dale, A.M., 2004. Automatically parcellating the human cerebral cortex. *Cereb. Cortex* 14, 11–22.
- Friederici, A.D., 2009. Pathways to language: fiber tracts in the human brain. *Trends Cogn. Sci.* 13, 175–181.
- Fusar-Poli, P., Radua, J., McGuire, P., Borgwardt, S., 2012. Neuroanatomical maps of psychosis onset: voxel-wise meta-analysis of antipsychotic-naïve VBM studies. *Schizophr. Bull.* 38, 1297–1307.
- Goldberg, T.E., Aloia, M.S., Gourovitch, M.L., Missar, D., Pickar, D., Weinberger, D.R., 1998. Cognitive substrates of thought disorder, I: the semantic system. *Am. J. Psychiatry* 155, 1671–1676.
- Han, Z., Ma, Y., Gong, G., He, Y., Caramazza, A., Bi, Y., 2013. White matter structural connectivity underlying semantic processing: evidence from brain damaged patients. *Brain* 136, 2952–2965.
- Harsan, L.A., Poulet, P., Guignard, B., Steibel, J., Parizel, N., de Sousa, P.L., Boehm, N., Grucker, D., Ghandour, M.S., 2006. Brain dysmyelination and recovery assessment by noninvasive in vivo diffusion tensor magnetic resonance imaging. *J. Neurosci. Res.* 83, 392–402.
- Hickok, G., Poeppel, D., 2004. Dorsal and ventral streams: a framework for understanding aspects of the functional anatomy of language. *Cognition* 92, 67–99.
- Hickok, G., Poeppel, D., 2007. The cortical organization of speech processing. *Nat. Rev. Neurosci.* 8, 393–402.
- Kay, S.R., Fiszbein, A., Opler, L.A., 1987. The Positive and Negative Syndrome Scale (PANSS) for schizophrenia. *Schizophr. Bull.* 13, 261–276.
- Kircher, T., Bröhl, H., Meier, F., Engelen, J., 2018. Formal thought disorders: from phenomenology to neurobiology. *Lancet Psychiatry* 5, 515–526.
- Koch, K., Wagner, G., Schachtzabel, C., Schultz, C.C., Gü, D., Rgen, J., Reichenbach, R., Sauer, H., Schlö, R.G.M., 2013. Neural activation and radial diffusivity in schizophrenia: combined fMRI and diffusion tensor imaging study. *Schizophr. Res.* 150, 93–100.
- Kraepelin, E., 1919. *Dementia Praecox and Paraphrenia*. E & S Livingstone, Edinburgh.
- Kubicki, M., Alvarado, J.L., Westin, C.-F., Tate, D.F., Markant, D., Terry, D.P., Whitford, T.J., De Siebenthal, J., Bouix, S., McCarley, R.W., Kikinis, R., Shenton, M.E., 2011. Stochastic tractography study of inferior frontal gyrus anatomical connectivity in schizophrenia. *Neuroimage* 55, 1657–1664.
- Kubicki, M., Shenton, M.E., Maciejewski, P.K., Pelavin, P.E., Hawley, K.J., Ballinger, T., Swisher, T., Jabbar, G.A., Thermenos, H.W., Keshavan, M.S., Seidman, L.J., Delisi, L.E., 2013. Decreased axial diffusivity within language connections: a possible biomarker of schizophrenia risk. *Schizophr. Res.* 148, 67–73.
- Lee, S.-H., Kubicki, M., Asami, T., Seidman, L.J., Goldstein, J.M., Mesholam-Gately, R.I., McCarley, R.W., Shenton, M.E., 2013. Extensive white matter abnormalities in patients with first-episode schizophrenia: a diffusion tensor imaging (DTI) study. *Schizophr. Res.* 143, 231–238.
- Lehrl, S., Triebig, G., Fischer, B., 1995. Multiple choice vocabulary test MWT as a valid and short test to estimate premorbid intelligence. *Acta Neurol. Scand.* 91, 335–345.
- Levy, D.L., Coleman, M.J., Sung, H., Ji, F., Matthyse, S., Mendell, N.R., Titone, D., 2010. The genetic basis of thought disorder and language and communication disturbances in schizophrenia. *J. Neurolinguistics* 23, 176.
- Lu, L.H., Zhou, X.J., Keedy, S.K., Reilly, J.L., Sweeney, J.A., 2011. White matter microstructure in untreated first episode bipolar disorder with psychosis: comparison with schizophrenia. *Bipolar Disord.* 13, 604–613.
- Lukoff, D., Liberman, R.P., Nuechterlein, K.H., 1986. Symptom monitoring in the rehabilitation of schizophrenic patients. *Schizophr. Bull.* 12, 578–602.
- Maldonado, L.L., de Champfleury, N.M., Velut, S., Destrieux, C., Zemmoura, I., Duffau, H., 2013. Evidence of a middle longitudinal fasciculus in the human brain from fiber dissection. *J. Anat.* 223, 38–45.
- Melicher, T., Horacek, J., Hlinka, J., Spaniel, F., Tintera, J., Ibrahim, I., Mikolas, P., Novak, T., Mohr, P., Hoschl, C., 2015. White matter changes in first episode psychosis and their relation to the size of sample studied: a DTI study. *Schizophr. Res.* 162, 22–28.
- Miller, J., 2009. What is the probability of replicating a statistically significant effect? *Psychon. Bull. Rev.* 16, 617–640. <https://doi.org/10.3758/PBR.16.4.617>.
- Morgan, C.J., Coleman, M.J., Ulgen, A., Boling, L., Cole, J.O., Johnson, F.V., Lerbinger, J., Bodkin, J.A., Holzman, P.S., Levy, D.L., 2017. Thought disorder in schizophrenia and bipolar disorder probands, their relatives, and nonpsychiatric controls. *Schizophr. Bull.* 43, 523–535.
- Oldfield, R.C., 1971. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 9, 97–113.
- Qiu, D., Tan, L.-H., Zhou, K., Khong, P.-L., 2008. Diffusion tensor imaging of normal white matter maturation from late childhood to young adulthood: Voxel-wise evaluation of mean diffusivity, fractional anisotropy, radial and axial diffusivities, and correlation with reading development. *Neuroimage* 41, 223–232. <https://doi.org/10.1016/j.neuroimage.2008.02.023>.
- Riecher-Rössler, A., Aston, J., Ventura, J., Merlo, M., Borgwardt, S., Gschwandtner, U., Stieglitz, R.-D., 2008. The Basel Screening Instrument for Psychosis (BSIP): development, structure, reliability and validity. *Fortschr. Neurol. Psychiatr.* 76, 207–216.
- Salavera, C., Puyuelo, M., Antoñanzas, J.L., Teruel, P., 2013. Semantics, pragmatics, and formal thought disorders in people with schizophrenia. *Neuropsychiatr. Dis. Treat.* 9, 177–183.
- Sarubbo, S., De Benedictis, A., Merler, S., Mandonnet, E., Barbareschi, M., Dallabona, M., Chioffi, F., Duffau, H., 2016. Structural and functional integration between dorsal and ventral language streams as revealed by blunt dissection and direct electrical stimulation. *Hum. Brain Mapp.* 37, 3858–3872.
- Saur, D., Kreher, B.W., Schnell, S., Kummerer, D., Kellmeyer, P., Vry, M.-S., Umarova, R., Musso, M., Glauche, V., Abel, S., Huber, W., Rijntjes, M., Hennig, J., Weiller, C., 2008. Ventral and dorsal pathways for language. *Proc. Natl. Acad. Sci.* 105, 18035–18040.
- Schmidt, A., Lenz, C., Smieskova, R., Harrisberger, F., Walter, A., Riecher-Rössler, A., Simon, A., Lang, U.E., McGuire, P., Fusar-Poli, P., Borgwardt, S., 2015. Brain diffusion changes in emerging psychosis and the impact of state-dependent psychopathology. *Neurosignals* 23, 71–83.
- Schmidt, A., Crossley, N.A., Harrisberger, F., Smieskova, R., Lenz, C., Riecher-Rössler, A., Lang, U.E., McGuire, P., Fusar-Poli, P., Borgwardt, S., 2017. Structural network disorganization in subjects at clinical high risk for psychosis. *Schizophr. Bull.* 43, 583–591.
- Schneiderman, J.S., Buchsbaum, M.S., Haznedar, M.M., Hazlett, E.A., Brickman, A.M., Shihabuddin, L., Brand, J.G., Torosjan, Y., Newmark, R.E., Canfield, E.L., Tang, C., Aronowitz, J., Paul-Oudouard, R., Hof, P.R., 2009. Age and diffusion tensor anisotropy in adolescent and adult patients with schizophrenia. *Neuroimage* 45, 662–671.
- Schonberger, R.B., Worden, W.S., Shahmohammadi, K., Menn, K., Silverman, T.J., Stout, R.G., Shelley, K.H., Silverman, D.G., 2006. Topical non-iontophoretic application of acetylcholine and nitroglycerin via a transcutaneous patch: a new means for assessing microvascular reactivity. *Yale J. Biol. Med.* 79, 1–7.
- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., Dunbar, G.C., 1998. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J. Clin. Psychiatry* 59 (Suppl. 2) (22–33–57).
- Shepherd, A.M., Laurens, K.R., Matheson, S.L., Carr, V.J., Green, M.J., 2012. Systematic meta-review and quality assessment of the structural brain alterations in schizophrenia. *Neurosci. Biobehav. Rev.* 36, 1342–1356.
- Skelly, L.R., Calhoun, V., Meda, S.A., Kim, J., Mathalon, D.H., Pearlson, G.D., 2008. Diffusion tensor imaging in schizophrenia: relationship to symptoms. *Schizophr. Res.* 98, 157–162.
- Song, S.-K., Sun, S.-W., Ju, W.-K., Lin, S.-J., Cross, A.H., Neufeld, A.H., 2003. Diffusion tensor imaging detects and differentiates axon and myelin degeneration in mouse optic nerve after retinal ischemia. *Neuroimage* 20, 1714–1722.
- Stegmayer, K., Stettler, M., Strik, W., Federspiel, A., Wiest, R., Bohlhalter, S., Walther, S., 2017. Resting state perfusion in the language network is linked to formal thought disorder and poor functional outcome in schizophrenia. *Acta Psychiatr. Scand.* 136, 506–516.
- Stricker, J.L., Brown, G.G., Wixted, J., Baldo, J.V., Delis, D.C., 2002. New semantic and serial clustering indices for the California Verbal Learning Test-Second Edition: background, rationale, and formulae. *J. Int. Neuropsychol. Soc.* 8, 425–435.
- Strik, W., Wopfler, A., Horn, H., Koschorke, P., Razavi, N., Walther, S., Wirtz, G., 2010. The Bern psychopathology scale for the assessment of system-specific psychotic symptoms. *Neuropsychobiology* 61, 197–209.
- Szeszko, P.R., Robinson, D.G., Ashtari, M., Vogel, J., Betensky, J., Sevy, S., Ardekani, B. a, Lencz, T., Malhotra, A.K., McCormack, J., Miller, R., Lim, K.O., Gunduz-Bruce, H., Kane, J.M., Bilder, R.M., 2008. Clinical and neuropsychological correlates of white matter abnormalities in recent onset schizophrenia. *Neuropsychopharmacology* 33, 976–984.
- Ungerleider, L.G., Haxby, J.V., 1994. “What” and “where” in the human brain. *Curr. Opin. Neurobiol.* 4, 157–165.
- Viher, P.V., Stegmayer, K., Giezendanner, S., Federspiel, A., Bohlhalter, S., Wiest, R., Strik, W., Walther, S., 2018. White matter correlates of the disorganized speech dimension in schizophrenia. *Eur. Arch. Psychiatry Clin. Neurosci.* 268, 99–104.
- Walther, S., Stegmayer, K., Sulzbacher, J., Vanbellingen, T., Müri, R., Strik, W., Bohlhalter, S., 2015. Nonverbal social communication and gesture control in schizophrenia. *Schizophr. Bull.* 41, 338–345.
- Walther, S., Stegmayer, K., Federspiel, A., Bohlhalter, S., Wiest, R., Viher, P.V., 2017. Aberrant Hyperconnectivity in the motor system at rest is linked to motor abnormalities in schizophrenia spectrum disorders. *Schizophr. Bull.* 43, 982–992.
- Woods, S.W., 2003. Chlorpromazine equivalent doses for the newer atypical antipsychotics. *J. Clin. Psychiatry* 64, 663–667. <https://doi.org/10.4088/jcp.v64n0607>.
- Woods, S.P., Delis, D.C., Scott, J.C., Kramer, J.H., Holdnack, J.A., 2006. The California verbal learning test – second edition: test-retest reliability, practice effects, and reliable change indices for the standard and alternate forms. *Arch. Clin. Neuropsychol.* 21, 413–420.
- Yendiki, A., Panneck, P., Srinivasan, P., Stevens, A., Zöllei, L., Augustinack, J., Wang, R., Salat, D., Ehrlich, S., Behrens, T., Jbabdi, S., Gollub, R., Fischl, B., 2011. Automated probabilistic reconstruction of white-matter pathways in health and disease using an atlas of the underlying anatomy. *Front. Neuroinform.* 5, 23.
- Yendiki, A., Koldewyn, K., Kakunoori, S., Kanwisher, N., Fischl, B., 2014. Spurious group differences due to head motion in a diffusion MRI study. *Neuroimage* 88, 79–90.
- Yendiki, A., Reuter, M., Wilkens, P., Rosas, H.D., Fischl, B., 2016. Joint reconstruction of white-matter pathways from longitudinal diffusion MRI data with anatomical priors. *Neuroimage* 127, 277–286.
- Yung, A.R., Phillips, L.J., McGorry, P.D., McFarlane, C.A., Francey, S., Harrigan, S., Patton, G.C., Jackson, H.J., 1998. Prediction of psychosis. A step towards indicated prevention of schizophrenia. *Br. J. Psychiatry. Suppl.* 172, 14–20.
- Zarei, M., Johansen-Berg, H., Jenkinson, M., Ciccarelli, O., Thompson, A.J., Matthews, P.M., 2007. Two-dimensional population map of cortical connections in the human internal capsule. *J. Magn. Reson. Imaging* 25, 48–54.

- Zikou, A.K., Xydis, V.G., Astrakas, L.G., Nakou, I., Tzarouchi, L.C., Tzoufi, M., Argyropoulou, M.I., 2016. Diffusion tensor imaging in children with tuberous sclerosis complex: tract-based spatial statistics assessment of brain microstructural changes. *Pediatr. Radiol.* 46, 1158–1164. <https://doi.org/10.1007/s00247-016-3582-2>.
- Zivadinov, R., Shucard, J.L., Hussein, S., Durfee, J., Cox, J.L., Bergsland, N., Dwyer, M.G., Benedict, R.H.B., Ambrus, J., Shucard, D.W., 2013. Multimodal imaging in systemic lupus erythematosus patients with diffuse neuropsychiatric involvement. *Lupus* 22, 675–683. <https://doi.org/10.1177/0961203313486193>.
- de Zubicaray, G.I., Rose, S.E., McMahon, K.L., 2011. The structure and connectivity of semantic memory in the healthy older adult brain. *Neuroimage* 54, 1488–1494.