



Abstract #15506

Abstract Information

Abstract ID:**15506****Abstract Title:****Experimental study of plagioclase growth, nucleation rates, and shape evolution during cooling of an anhydrous basaltic andesite.****Authors (presenting author in bold):****Melvyn M. R. Billon**¹, Bernard Charlier², Olivier Namur³ and Jacqueline E. Vander Auwera¹, (1)University of Liege, (2)University of Liège, (3)KU Leuven**Abstract Text:**

Crystal textures are commonly used to assess the physical and chemical state of pre and syn-eruptive magmatic processes. Plagioclase nucleation and growth rates have been investigated in a range of composition [1,2], but basaltic andesite is underrepresented in the experimental dataset. In this study, a series of experiments were performed at 1 atm on an anhydrous natural basaltic andesite from Osorno volcano (Central Southern Volcanic Zone, Chile) at NNO to determine its nucleation and growth rates and to constrain the role of the cooling rate, isothermal dwells, and initial superheating on the plagioclase habit. Three series of cooling-driven experiments were conducted: (A) initial equilibration at 1190°C, above the liquidus (1180 ± 5 °C), during 24h followed by different cooling rates (1°C/h, 3°C/h, 9°C/h); (B) same as in (A) followed by several isothermal dwells; (C) initial heating at 1450°C followed by a 1°C/h cooling rate. All runs were quenched at different temperatures (1165°C, 1140°C, 1120°C, 1100°C). We used different methods to calculate the growth and nucleation rates: batch method (image analyses), maximum size (l_{max}) and Crystal Size Distribution (CSD). They range from 1.05×10^{-7} to 3.6×10^{-6} mm.s⁻¹ and from 3.17×10^{-2} to 2.8 mm⁻³.s⁻¹, respectively. These results and especially the growth rates are comparable with those of [2] (10^{-7} to 10^{-6} mm.s⁻¹) who performed hydrous experiments on a basaltic andesite. With decreasing temperature, there is a general trend in plagioclase shape to evolve from elongate to prismatic/bladed as observed by [2]. However, we observe a large variability of shape for each undercooling value (3D modeling with ShapeCalc [3]). The proportion of the largest and small crystals respectively increases and decreases at constant crystallinity and with increasing dwell time suggesting Ostwald ripening. Finally, the superheating step removes potential

nuclei/germs (more anorthitic residual cores) and induces the crystallization of larger crystals at temperatures significantly lower than the liquidus.

[1] Pupier, E. et al. (2008). *Contributions to Mineralogy and Petrology* 155, 555. [2] Shea T., Hammer J. E. (2013). *Journal of Volcanology and Geothermal Research* 260, 127. [3] Mangler, M.F. et al. (2022). *Contributions to Mineralogy and Petrology* 177, 64.

Final Paper##:

19

Keywords:

Experiments, Growth and Nucleation

Submitter's E-mail Address:

melvyn.billon@uliege.be

Submitter Full Name:

Melvyn BILLON

Preferred Presentation Style:

Oral

Conference Attendance:

In person

Registration:

I understand that the presenting author named in this submission must register for the conference by 31 May 2023.

Submission Permission and Copyright:

I grant non-exclusive copyright for my abstract and any related uploaded content to the Conference Exchange and the conference owners. I give permission for my details to be stored in the conference database. I, and the presenting author (if different than the submitter), agree to be contacted by the conference organizers about this submission. I agree to the recording and online streaming of any presentations scheduled as a result of this submission. I confirm all listed authors have agreed to these terms.

First Author**Presenting Author**

Melvyn M. R. Billon

Email: melvyn.billon@uliege.be

University of Liege
Petrology and geochemistry of igneous rocks
PhD Student
Allée du six Août 12, Department of Geology (B20)
Liege Belgium 4000
Belgium

Second Author

Bernard Charlier
Email: b.charlier@uliege.be

University of Liège
Geology
Liège 4000
Belgium

Third Author

Olivier Namur
Email: olivier.namur@kuleuven.be

KU Leuven
Earth and Environmental Sciences
Celestijnenlaan 200E
Heverlee 3001
Belgium

Fourth Author

Jacqueline E. Vander Auwera
Email: jvdauwera@uliege.be

University of Liege
Geology
Prof.
Sart Tilman (B20)
Liege Liege 4000
Belgium