

Reconsideration of marine bivalves as mercury biomonitoring tool: a national isotopic survey along the coast of South Korea.

ORAL PRESENTATION

Presented by Lucien Besnard



Monday, 10 July 2023



15:00 - 15:15



R17 (Building B, Level 2, Lyon Congress Center)

Session: 12aO2 - Mercury cycling in the context of global change

Symposium Session: 12a - Mercury cycling in the context of global change

Theme: Theme 12: Environmental geochemistry and human health


Abstract


Coastal sediments act as a reservoir for inorganic and organic contaminants. Among them, mercury (Hg), a toxic and bioaccumulative heavy metal, is efficiently transferred from sediment to marine biota and ultimately to humans via fisheries consumption. Marine bivalves have been recognized as an effective bioindicator for sediment pollution as their suspension or deposit feeding behaviors are physically and chemically tied to the sediment. Due to their broad geographic distribution, bivalve-based monitoring programs, known as the 'Mussel Watch', had been developed around the world including South Korea. However, due to multiple natural and anthropogenic Hg sources, absence of significant Hg relationships between bivalves and sediment have been reported in coastal regions. To investigate bivalve Hg sources and efficiency as coastal sediment Hg bioindicators, paired sediment and bivalves (blue mussel, *Mytilus edulis*, and Pacific oyster, *Magallana gigas*) were sampled in 50 coastal sites in South Korea and measured for total Hg (THg) concentration and Hg isotope ratios. Sediment THg varied from 0.7 to 195.7 ng·g⁻¹, except for two highly contaminated sites at Pohang (600.1 ng·g⁻¹) and Onsan (1417.9 ng·g⁻¹). Bivalve THg ranged between 21.1 and 225.2 ng·g⁻¹ and was not correlated with sediment THg (linear regression, $R^2=0.01$, $p>0.05$). Only a handful of

studies have compared Hg isotope ratios between bivalves and sediment and observed isotopic differences (mainly $\delta^{202}\text{Hg}$ values) were either attributed to biogeochemical processes occurring in the different fractions of the sediment prior to exposure or to the preferential accumulation of dissolved Hg from the water column by bivalves [1,2]. Based on these identified shifts, Hg isotope ratios are used to identify biogeochemical and/or ecological factors dictating the extent of sediment Hg bioaccumulation to bivalves. Finally, we further plan to quantify carbon, nitrogen, and sulfur isotopes to assess the importance of species feeding behavior and dietary sources on Hg bioaccumulation.

[1] Kwon et al., Mercury isotope study of sources and exposure pathways of methylmercury in estuarine food webs in the Northeastern US. *Environ.Sci.Technol.* **2014**, 48, 10089-10097.

[2] Li et al., Environmental origins of methylmercury accumulated in subarctic estuarine fish indicated by mercury stable isotopes. *Environ.Sci.Technol.* **2016**, 50, 11559-11568.





Presenting Author



Lucien Besnard
Pohang University of Science and Technology

Authors



Kongtae Ra
Korea Institute of Ocean Science and Technology



Gilles Lepoint
University of Liège (ULg), Laboratoire d'Ecologie trophique et isotopique (LETIS)



Saebom Jung
Pohang University of Science and Technology



Sae Yun Kwon
Pohang University of Science and Technology

