

GFR in Current Clinical Guidelines

Pierre Delanaye, MD, PhD
University of Liège
CHU Sart Tilman
BELGIUM






Category	Disclosure Information
Employer	Nothing to disclose.
Ownership Interest	Nothing to disclose.
Consultancy	IDS; Nephrolyx; Alentis Therapeutics; ARK Bioscience; Astellas
Research Funding	Nothing to disclose.
Honoraria	IDS; Fresenius Kabi; Fresenius Medical Care; Nephrolyx; Alentis Therapeutics; ARK Bioscience; AstraZeneca; Bayer
Patents or Royalties	Nothing to disclose.
Advisory or Leadership Role	Nothing to disclose.
Speakers Bureau	Nothing to disclose.
Other Interests or Relationships	Nothing to disclose.
Disclosure Updated Date	07/18/2023

The Glomerular Filtration Rate is usually the best parameter to assess the global kidney function.



Homer William Smith (January 2, 1895 – March 25, 1962)

- GFR is estimated with biomarkers
- Serum creatinine is one the most prescribed analysis
- The most important is probably to know the limitations...



NephroTest Cohort (France)
 Which GFR for patients with
 serum creatinine measured
 at $80 \mu\text{mol/L}$ (0.9 mg/dL)?

IC 95% for subjects <65 years old
 IC 95% for subjects >65 years old

S. Creatinine lab
 normality range

Other Limitations

Analytical

- Jaffe methods
- Enzymatic methods
- Jaffe and enzymatic methods gives slightly different results
- Pseudochromogen: glucose, fructose, ascorbate, proteins, urate, acetoacetate, acetone, pyruvate => false positive
- Bilirubins: false negative

Physiological: Tubular secretion

- 10 to 40%
- Increase with decreased GFR
- Unpredictable at the individual level !

Physiological: Muscular mass

- Production (relatively) constant but muscular production => serum creatinine is dependent of muscular mass, not only GFR (age? sex/gender? race/population?)
- Extra-renal production

Perrone RD, Clin Chem, 1992, 38, p1933

Delanaye P, Nephron, 2017, 136, p302

Creatinine: to the trash?

- Very cheap (0.04€ /Jaffe)
- Good specificity
- Good analytical CV, IDMS traceability

Creatinine clearance

- Not recommended
- Creatinine tubular secretion
- Lack of precision:

errors in urine collection


22 to 27% for « trained » patients

50 to 70 % for others

large intra-individual variability for
creatinine excretion

Statistics

- Good correlation: a “*sine qua non*” condition but insufficient
- Bias: mean difference between two values = the systematic error
- Precision: SD around the bias = the random error
- Accuracy 30% = % of eGFR between $\pm 30\%$ of measured GFR



Which one?

- Cockcroft
- CKD-EPI
- EKFC

Which one?

- ~~• Cockcroft~~
- CKD-EPI
- EKFC

The CKD-EPI equation

A New Equation to Estimate Glomerular Filtration Rate

Andrew S. Levey, MD; Lesley A. Stevens, MD, MS; Christopher H. Schmid, PhD; Yaping (Lucy) Zhang, MS; Alejandro F. Castro III, MPH; Harold I. Feldman, MD, MSCE; John W. Kusek, PhD; Paul Eggers, PhD; Frederick Van Lente, PhD; Tom Greene, PhD; and Josef Coresh, MD, PhD, MHS, for the CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration)*


Ann Intern Med. 2009;150:604-612.

Table 2. The CKD-EPI Equation for Estimating GFR on the Natural Scale*

Race and Sex	Serum Creatinine Level, $\mu\text{mol/L}$ (mg/dL)	Equation
Black		
Female	≤ 62 (≤ 0.7)	$\text{GFR} = 166 \times (\text{Scr}/0.7)^{-0.329} \times (0.993)^{\text{Age}}$
	> 62 (> 0.7)	$\text{GFR} = 166 \times (\text{Scr}/0.7)^{-1.209} \times (0.993)^{\text{Age}}$
Male	≤ 80 (≤ 0.9)	$\text{GFR} = 163 \times (\text{Scr}/0.9)^{-0.411} \times (0.993)^{\text{Age}}$
	> 80 (> 0.9)	$\text{GFR} = 163 \times (\text{Scr}/0.9)^{-1.209} \times (0.993)^{\text{Age}}$
White or other		
Female	≤ 62 (≤ 0.7)	$\text{GFR} = 144 \times (\text{Scr}/0.7)^{-0.329} \times (0.993)^{\text{Age}}$
	> 62 (> 0.7)	$\text{GFR} = 144 \times (\text{Scr}/0.7)^{-1.209} \times (0.993)^{\text{Age}}$
Male	≤ 80 (≤ 0.9)	$\text{GFR} = 141 \times (\text{Scr}/0.9)^{-0.411} \times (0.993)^{\text{Age}}$
	> 80 (> 0.9)	$\text{GFR} = 141 \times (\text{Scr}/0.9)^{-1.209} \times (0.993)^{\text{Age}}$

- CKD-EPI
- Development dataset: n=5504
- Internal validation: n=2750
- External validation: n=3896
- Creatinine calibrated
- Median GFR in the development = 68 mL/min/1.73 m²

Figure. Performance of the CKD-EPI and MDRD Study equations in estimating measured GFR in the external validation data set.



KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease

VOLUME 3 | ISSUE 1 | JANUARY 2013

<http://www.kidney-international.org>



Revised in 2023, Public Review ended

CKD-EPI: What else?

Development and Validation of a Modified Full Age Spectrum Creatinine-Based Equation to Estimate Glomerular Filtration Rate

A Cross-sectional Analysis of Pooled Data

Hans Pottel, PhD*; Jonas Björk, PhD*; Marie Courbebaisse, MD, PhD; Lionel Couzi, MD, PhD; Natalie Ebert, MD, MPH; Björn O. Eriksen, MD, PhD; R. Neil Dalton, PhD; Laurence Dubourg, MD, PhD; François Gaillard, MD, PhD; Cyril Garrouste, MD; Anders Grubb, MD, PhD; Lola Jacquemont, MD, PhD; Magnus Hansson, MD, PhD; Nassim Kamar, MD, PhD; Edmund J. Lamb, PhD; Christophe Legendre, MD; Karin Littmann, MD; Christophe Mariat, MD, PhD; Toralf Melsom, MD, PhD; Lionel Rostaing, MD, PhD; Andrew D. Rule, MD; Elke Schaeffner, MD, PhD, MSc; Per-Ola Sundin, MD, PhD; Stephen Turner, MD, PhD; Arend Bökenkamp, MD; Ulla Berg, MD, PhD; Kajsa Åsling-Monemi, MD, PhD; Luciano Selistre, MD, PhD; Anna Åkesson, BSc; Anders Larsson, MD, PhD; Ulf Nyman, MD, PhD†; and Pierre Delanaye, MD, PhD†

Measured GFR and IDMS traceable creatinine

N=11,251 in the development and internal validation dataset

N=8,378 in the external validation dataset

N=1,254 between 2 and 18 years

7+6 cohorts

White people

Figure 1. The new EKFC equation.

Age	SCr/Q	Equation
2–40 y	<1	$107.3 \times (SCr/Q)^{-0.322}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132}$
>40 y	<1	$107.3 \times (SCr/Q)^{-0.322} \times 0.990^{(Age - 40)}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132} \times 0.990^{(Age - 40)}$

Q Values

For ages 2–25 y:

Males:

$$\ln(Q) = 3.200 + 0.259 \times Age - 0.543 \times \ln(Age) - 0.00763 \times Age^2 + 0.0000790 \times Age^3$$

Females:

$$\ln(Q) = 3.080 + 0.177 \times Age - 0.223 \times \ln(Age) - 0.00596 \times Age^2 + 0.0000686 \times Age^3$$

For ages >25 y:

Males:

$$Q = 80 \mu\text{mol/L (0.90 mg/dL)}$$

Females:

$$Q = 62 \mu\text{mol/L (0.70 mg/dL)}$$

SCr and Q in $\mu\text{mol/L}$ (to convert to mg/dL, divide by 88.4)

Q values (in $\mu\text{mol/L}$ or mg/dL) correspond to the median SCr values for the age- and sex-specific populations. EKFC = European Kidney Function Consortium; SCr = serum creatinine.

Figure 1. The new EKFC equation.

Age	SCr/Q	Equation
2–40 y	<1	$107.3 \times (SCr/Q)^{-1.322}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132}$
>40 y	<1	$107.3 \times (SCr/Q)^{-1.322} \times 0.990^{(Age - 40)}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132} \times 0.990^{(Age - 40)}$

Q Values

For ages 2–25 y:

Males:

$$\ln(Q) = 3.200 + 0.259 \times Age - 0.543 \times \ln(Age) - 0.00763 \times Age^2 + 0.0000790 \times Age^3$$

Females:

$$\ln(Q) = 3.080 + 0.177 \times Age - 0.223 \times \ln(Age) - 0.00596 \times Age^2 + 0.0000686 \times Age^3$$

For ages >25 y:

Males:

$$Q = 80 \mu\text{mol/L (0.90 mg/dL)}$$

Females:

$$Q = 62 \mu\text{mol/L (0.70 mg/dL)}$$

SCr and Q in $\mu\text{mol/L}$ (to convert to mg/dL, divide by 88.4)

Q values (in $\mu\text{mol/L}$ or mg/dL) correspond to the median SCr values for the age- and sex-specific populations. EKFC = European Kidney Function Consortium; SCr = serum creatinine.

Figure 1. The new EKFC equation.

Age	SCr/Q	Equation
2–40 y	<1	$107.3 \times (SCr/Q)^{-0.322}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132}$
>40 y	<1	$107.3 \times (SCr/Q)^{-0.322} \times 0.990^{(Age - 40)}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132} \times 0.990^{(Age - 40)}$

Q Values

For ages 2–25 y:

Males:

$$\ln(Q) = 3.200 + 0.259 \times Age - 0.543 \times \ln(Age) - 0.00763 \times Age^2 + 0.0000790 \times Age^3$$

Females:

$$\ln(Q) = 3.080 + 0.177 \times Age - 0.223 \times \ln(Age) - 0.00596 \times Age^2 + 0.0000686 \times Age^3$$

For ages >25 y:

Males:

$$Q = 80 \mu\text{mol/L (0.90 mg/dL)}$$

Females:

$$Q = 62 \mu\text{mol/L (0.70 mg/dL)}$$

SCr and Q in $\mu\text{mol/L}$ (to convert to mg/dL, divide by 88.4)

Q values (in $\mu\text{mol/L}$ or mg/dL) correspond to the median SCr values for the age- and sex-specific populations. EKFC = European Kidney Function Consortium; SCr = serum creatinine.

Pierre Delanaye*, François Gaillard, Jessica van der Weijden, Geir Mjøen, Ingela Ferhman-Ekholm, Laurence Dubourg, Natalie Ebert, Elke Schaeffner, Torbjörn Åkerfeldt, Karolien Goffin, Lionel Couzi, Cyril Garrouste, Lionel Rostaing, Marie Courbebaisse, Christophe Legendre, Maryvonne Hourmant, Nassim Kamar, Etienne Cavalier, Laurent Weekers, Antoine Bouqueneau, Martin H. de Borst, Christophe Mariat, Hans Pottel and Marco van Londen

Age-adapted percentiles of measured glomerular filtration in healthy individuals: extrapolation to living kidney donors over 65 years

Before 40 y: mGFR = 107 mL/min/1.73m²
...and it seems universal...




Figure 3: mGFR according to age in the development (dark dots) and external validation cohort (n=329) (gray dots). Red lines are percentiles 5, 10, 50, 90 and 95, calculated from kidney donors younger than 65 years and extrapolated for ages >65 years.

Pierre Delanaye*, François Gaillard, Jessica van der Weijden, Geir Mjøen, Ingela Ferhman-Ekholm, Laurence Dubourg, Natalie Ebert, Elke Schaeffner, Torbjörn Åkerfeldt, Karolien Goffin, Lionel Couzi, Cyril Garrouste, Lionel Rostaing, Marie Courbebaisse, Christophe Legendre, Maryvonne Hourmant, Nassim Kamar, Etienne Cavalier, Laurent Weekers, Antoine Bouqueneau, Martin H. de Borst, Christophe Mariat, Hans Pottel and Marco van Londen

Age-adapted percentiles of measured glomerular filtration in healthy individuals: extrapolation to living kidney donors over 65 years

Before 40 y: mGFR = 107 mL/min/1.73m²
...and it seems universal...






Figure 3: mGFR according to age in the development (dark dots) and external validation cohort (n=329) (gray dots). Red lines are percentiles 5, 10, 50, 90 and 95, calculated from kidney donors younger than 65 years and extrapolated for ages >65 years.



N=83,257 from three labs
(Sweden, Belgium)

62 $\mu\text{mol/L}$ = 0,70 mg/dL



80 $\mu\text{mol/L}$ = 0,90 mg/dL

Figure 1. The new EKFC equation.

Age	SCr/Q	Equation
2–40 y	<1	$107.3 \times (SCr/Q)^{-0.322}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132}$
>40 y	<1	$107.3 \times (SCr/Q)^{-0.322} \times 0.990^{(Age - 40)}$
	≥ 1	$107.3 \times (SCr/Q)^{-1.132} \times 0.990^{(Age - 40)}$

Q Values

For ages 2–25 y:

Males:

$$\ln(Q) = 3.200 + 0.259 \times Age - 0.543 \times \ln(Age) - 0.00763 \times Age^2 + 0.0000790 \times Age^3$$

Females:

$$\ln(Q) = 3.080 + 0.177 \times Age - 0.223 \times \ln(Age) - 0.00596 \times Age^2 + 0.0000686 \times Age^3$$

For ages >25 y:

Males:

$$Q = 80 \mu\text{mol/L (0.90 mg/dL)}$$

Females:

$$Q = 62 \mu\text{mol/L (0.70 mg/dL)}$$

SCr and Q in $\mu\text{mol/L}$ (to convert to mg/dL, divide by 88.4)

Q values (in $\mu\text{mol/L}$ or mg/dL) correspond to the median SCr values for the age- and sex-specific populations. EKFC = European Kidney Function Consortium; SCr = serum creatinine.

Figure 1. The new EKFC equation.

Age	SCr/Q	Equation
2–40 y	<1	$107.3 \times (\text{SCr}/Q)^{-0.322}$
	≥ 1	$107.3 \times (\text{SCr}/Q)^{-1.132}$
>40 y	<1	$107.3 \times (\text{SCr}/Q)^{-0.322} \times 0.990^{(\text{Age} - 40)}$
	≥ 1	$107.3 \times (\text{SCr}/Q)^{-1.132} \times 0.990^{(\text{Age} - 40)}$

Q Values

For ages 2–25 y:

Males:

$$\ln(Q) = 3.200 + 0.259 \times \text{Age} - 0.543 \times \ln(\text{Age}) - 0.00763 \times \text{Age}^2 + 0.0000790 \times \text{Age}^3$$

Females:

$$\ln(Q) = 3.080 + 0.177 \times \text{Age} - 0.223 \times \ln(\text{Age}) - 0.00596 \times \text{Age}^2 + 0.0000686 \times \text{Age}^3$$

For ages >25 y:

Males:


$$Q = 80 \mu\text{mol/L} (0.90 \text{ mg/dL})$$

Females:

$$Q = 62 \mu\text{mol/L} (0.70 \text{ mg/dL})$$

SCr and Q in $\mu\text{mol/L}$ (to convert to mg/dL, divide by 88.4)

Q values (in $\mu\text{mol/L}$ or mg/dL) correspond to the median SCr values for the age- and sex-specific populations. EKFC = European Kidney Function Consortium; SCr = serum creatinine.






Figure S8. P30-accuracy against age for the EKFC, FAS, CKiD and CKD-EPI equation in the external validation dataset. P30 (%) was graphically presented across the age spectrum using cubic splines with two free knots and using 3rd degree polynomials.

EKFC: added value

- Better performance (not more expensive)
- More « physiological»: correction at the serum creatinine level (sex, race), age better conceptualized, « Q » specific to specific populations
- Valid from 2y to old ages
- Children: no need for height
- No implausible jump at transition adolescence/young adults

Debate on the race factor in USA

Serum creatinine is different between Black and non-Black people in USA (and we don't know why!)

(normal) mGFR is not different

The race coefficient in the CKD-EPI₂₀₀₉ was considered as a source of discrimination



ORIGINAL ARTICLE

New Creatinine- and Cystatin C–Based Equations to Estimate GFR without Race

L.A. Inker, N.D. Eneanya, J. Coresh, H. Tighiouart, D. Wang, Y. Sang, D.C. Crews, A. Doria, M.M. Estrella, M. Froissart, M.E. Grams, T. Greene, A. Grubb, V. Gudnason, O.M. Gutiérrez, R. Kalil, A.B. Karger, M. Mauer, G. Navis, R.G. Nelson, E.D. Poggio, R. Rodby, P. Rossing, A.D. Rule, E. Selvin, J.C. Seegmiller, M.G. Shlipak, V.E. Torres, W. Yang, S.H. Ballew, S.J. Couture, N.R. Powe, and A.S. Levey, for the Chronic Kidney Disease Epidemiology Collaboration*

➤ [N Engl J Med. 2021 Nov 4;385\(19\):1737-1749.](#)

Table 3. Accuracy of Current and New Approaches for GFR Estimation as Compared with Measured GFR in the Validation Data Set.

Filtration Marker and Equation*	Black Participants	Non-Black Participants	Difference between Black Participants and Non-Black Participants (95% CI)†
Bias: Median Difference between Measured GFR and eGFR (95% CI)‡			
<i>milliliters per minute per 1.73 square meters</i>			
Creatinine			
eGFRcr(ASR), current	-3.7 (-5.4 to -1.8)	-0.5 (-0.9 to 0.0)	-3.2 (-5.0 to -1.3)
eGFRcr(ASR-NB), new	7.1 (5.9 to 8.8)	-0.5 (-0.9 to 0.0)	7.6 (6.1 to 9.0)
eGFRcr(AS), new	3.6 (1.8 to 5.5)	-3.9 (-4.4 to -3.4)	7.6 (5.6 to 9.5)
Creatinine			
eGFRcr(ASR), current	85.1 (82.2 to 87.9)	89.5 (88.5 to 90.4)	-4.4 (-7.6 to -1.2)
eGFRcr(ASR-NB), new	86.4 (83.4 to 89.1)	89.5 (88.5 to 90.4)	-3.1 (-6.2 to 0)
eGFRcr(AS), new	87.2 (84.5 to 90.0)	86.5 (85.4 to 87.6)	0.7 (-2.4 to 3.8)

NKF and ASN Release New Way to Diagnose Kidney Diseases



Both Organizations Recommend Race-Free Approach to Estimate GFR

Sept. 23, 2021, New York, NY - Today, the National Kidney Foundation (NKF) and the American Society of Nephrology (ASN) Task Force on Reassessing the Inclusion of Race in Diagnosing Kidney Diseases has released its final report, which outlines a new race-free approach to diagnose kidney disease. In the report, the NKF-ASN Task Force recommends the adoption of the **new eGFR 2021 CKD EPI creatinine** equation that estimates kidney function without a race variable. The task force also recommended increased use of **cystatin C** combined with serum (blood) creatinine, as a confirmatory assessment of GFR or kidney function.

Ethnic/race factor in Europe/Africa?

RESEARCH LETTER

Performance of GFR Estimating Equations in African Europeans: Basis for a Lower Race-Ethnicity Factor Than in African Americans

Flamant M et al Am J Kidney Dis, 2013, 62, p179

NO !

Hindawi
International Journal of Nephrology
Volume 2020, Article ID 2141035, 9 pages
<https://doi.org/10.1155/2020/2141035>



Research Article

No Race-Ethnicity Adjustment in CKD-EPI Equations Is Required for Estimating Glomerular Filtration Rate in the Brazilian Population

ARTICLE IN PRESS

www.kidney-international.org

[clinical investigation](http://clinicalinvestigation)

RESEARCH ARTICLE

Performance of glomerular filtration rate estimation equations in Congolese healthy adults: The inopportunity of the ethnic correction

Justine B. Bukabau^{1*}, Ernest K. Sumaili¹, Etienne Cavalier², Hans Pottel³, Bejos Kifakiou⁴, Aliocha Nkodila¹, Jean Robert R. Makulo¹, Vieux M. Mokoli¹, Chantal V. Zinga¹, Augustin L. Longo¹, Yannick M. Engole¹, Yannick M. Nlandu¹, François B. Lepira¹, Nazaire M. Nseka¹, Jean Marie Krzesinski⁴, Pierre Delanaye⁴

¹ Renal Unit, Department of Internal medicine, Kinshasa University Hospital, University of Kinshasa, Kinshasa, Democratic Republic of the Congo, ² Division of Clinical Chemistry, CHU Sart Tilman (ULg CHU), University of Liège, Liège, Belgium, ³ Division of Public Health and Primary Care, KU Leuven Campus Kulak Kortrijk, Kortrijk, Belgium, ⁴ Division of Nephrology-Dialysis-Transplantation, CHU Sart Tilman (ULg CHU), University of Liège, Liège, Belgium


* justinebuk@yahoo.fr

Performance of creatinine- or cystatin C-based equations to estimate glomerular filtration rate in sub-Saharan African populations

Justine B. Bukabau^{1,7}, Eric Yayo^{2,7}, Appolinaire Gnionsahé³, Dagui Monnet², Hans Pottel⁴, Etienne Cavalier⁵, Aliocha Nkodila¹, Jean Robert R. Makulo¹, Vieux M. Mokoli¹, François B. Lepira¹, Nazaire M. Nseka¹, Jean-Marie Krzesinski⁶, Ernest K. Sumaili^{1,7} and Pierre Delanaye^{6,7}

¹Renal Unit, Department of Internal Medicine, Kinshasa University Hospital, University of Kinshasa, Kinshasa, Democratic Republic of Congo; ²Département de Biochimie, UFR Sciences Pharmaceutiques et Biologiques, Université Felix Houphouët Boigny, Abidjan, Ivory Coast; ³Département de Néphrologie, UFR Sciences Médicales, Université Felix Houphouët Boigny, Abidjan, Ivory Coast; ⁴Department of Public Health and Primary Care, KU Leuven Campus Kulak Kortrijk, Kortrijk, Belgium; ⁵Division of Clinical Chemistry, CHU Sart Tilman (ULg CHU), University of Liège, Liège, Belgium; and ⁶Division of Nephrology-Dialysis-Transplantation, CHU Sart Tilman (ULg CHU), University of Liège, Liège, Belgium

Yayo ES, Nephrol Ther, 2016, 12, p454
Flamant M, Am J Kidney Dis, 2013, 62, p179
Bukabau JB, Plos One, 2018, 13, e0193384
Bukabau JB, Kidney Int, 2019, 95, p1181














Nephrology Dialysis Transplantation (2023) 38: 106–118

<https://doi.org/10.1093/ndt/gfac241>

Advance Access publication date 24 August 2022




Performance of creatinine-based equations to estimate glomerular filtration rate in White and Black populations in Europe, Brazil and Africa

Pierre Delanaye ^{1,2,*}, Emmanuelle Vidal-Petiot ^{3,*}, Jonas Björk ^{4,5}, Natalie Ebert ⁶, Björn O. Eriksen⁷, Laurence Dubourg⁸, Anders Grubb⁹, Magnus Hansson¹⁰, Karin Littmann¹¹, Christophe Mariat¹², Toralf Melsom⁷, Elke Schaeffner⁶, Per-Ola Sundin ¹³, Arend Bökenkamp¹⁴, Ulla B. Berg¹⁵, Kajsa Åsling-Monemi¹⁵, Anna Åkesson^{4,5}, Anders Larsson¹⁶, Etienne Cavalier ¹⁷, R. Neil Dalton¹⁸, Marie Courbebaisse¹⁹, Lionel Couzi ²⁰, Francois Gaillard ²¹, Cyril Garrouste²², Lola Jacquemont²³, Nassim Kamar²⁴, Christophe Legendre²⁵, Lionel Rostaing ²⁶, Thomas Stehlé ^{27,28}, Jean-Philippe Haymann²⁹, Luciano da Silva Selistre³⁰, Jorge P. Strogoff-de-Matos ³¹, Justine B. Bukabau³², Ernest K. Sumaili³², Eric Yayo³³, Dagui Monnet³³, Ulf Nyman³⁴, Hans Pottel^{35,†} and Martin Flamant^{36,†}


Methods

- Adults, measured GFR, IDMS creatinine
- EKFC consortium: 11 cohorts from Europe (n=17,321)
- Data from Paris (n=4,429, among them 964 Black Europeans)
- Data from Africa (RDC and Côte d'Ivoire, n=508)


White Europeans (n=17,321)



Black Europeans (n=964)



Black Africans (n=508)



EFLM Paper

Pierre Delanaye, Elke Schaeffner, Mario Cozzolino, Michel Langlois, Mario Plebani, Tomris Ozben and Etienne Cavalier*, on behalf of the Board members of the EFLM Task Group Chronic Kidney Diseases

The new, race-free, Chronic Kidney Disease Epidemiology Consortium (CKD-EPI) equation to estimate glomerular filtration rate: is it applicable in Europe? A position statement by the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM)


Nephrol Dial Transplant (2023) 38: 1–6

<https://doi.org/10.1093/ndt/gfac254>

Advance Access publication date 7 September 2022



What should European nephrology do with the new CKD-EPI equation?

Ron T. Gansevoort ¹, Hans-Joachim Anders², Mario Cozzolino³, Danilo Fliser⁴, Denis Fouque⁵, Alberto Ortiz^{6,7}, Maria José Soler⁸ and Christoph Wanner⁹

¹Department of Nephrology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, ²Renal Division, Hospital of the Ludwig Maximilians University, Munich, Germany, ³Department of Health Sciences, University of Milan, Renal Division, ASST Santi Paolo e Carlo, Milan, Italy, ⁴Department of Internal Medicine IV, Renal and Hypertensive Disease, University Medical Center, Homburg, Saar, Germany, ⁵Department of Nephrology, Hospices Civils de Lyon, Centre Hospitalier Lyon-Sud, Pierre-Benite, University of Lyon, France, ⁶Department of Nephrology, IIS-Fundacion Jimenez Diaz- UAM, Madrid, Spain, ⁷Department of Medicine, Universidad Autonoma de Madrid, Madrid, Spain, ⁸Department of Nephrology, Hospital Vall d'Hebron, Barcelona, Vall d'Hebron Research Institute (VHIR), Barcelona, Spain and ⁹Department of Internal Medicine I and Comprehensive Heart Failure Center, University Hospital Würzburg, Würzburg, Germany

Correspondence to: Ron T. Gansevoort; E-mail: r.t.gansevoort@umcg.nl

Validation of EKFC in US populations

Cohorts	Sample Size	Age (years)	Measured GFR (mL/min/1.73m ²)	% of women	% of Black subjects	Proportion of individuals with urinary clearance data available
<i>All</i>	12,854	56.0 [22.1]	57 [46]	44.3	21.7	93.2
AASK	1,844	54.5 [16.0]	57 [35]	35.9	100	100
ALTOLD	381	43.3 [19.0]	97 [18]	65.1	1.8	0
CRIC	1,194	59.0 [17.7]	48 [28]	44.4	44.7	100
CRISP	217	34.0 [13.0]	93 [34]	59.0	11.1	100
DCCT/EDIC	809	31.0 [9.0]	119 [25]	47.8	1.4	100
GENOA/ECAC	1,093	66.1 [12.1]	80 [27]	56.6	0	100
Mayo Clinic	5,069	59.0 [21.0]	50 [40]	44.6	2.0	100
MDRD	1,756	51.0 [21.0]	36 [29]	39.5	12.4	100
PERL	491	52.0 [15.0]	70 [25]	33.6	10.8	0

Results are expressed in % or Median [interquartile range].

GFR: glomerular filtration rate


Q-values could be population specific

Q-values determined in different populations


	Q value in women	Q value in men	Origine
White European	0.70	0.90	Large data from laboratories in Sweden and Belgium
Black European	0.74	1.02	Living kidney donors in Paris
Black Africans (Central Africa)	0.72	0.96	Healthy people in Congo
White US population-specific	0.73	0.93	Large data from laboratories from University of Washington Medicine System
Black US population-specific	0.73	1.00	Large data from laboratories from University of Washington Medicine System
White US population-specific	0.70	0.94	National Health and Nutrition Examination Survey
Black US population-specific	0.72	1.03	National Health and Nutrition Examination Survey
US race-free	0.73	0.97	Large data from laboratories from University of Washington Medicine System
China	0.62	0.88	27,830 healthy people

All results are expressed in mg/dL


A



B




Bias (A) and accuracy within 30% (P30) (B) for the CKD-EPI₂₀₂₁ and the EKFC in Black women (n=1,087) according to age.



Legend: CKD-EPI₂₀₂₁: race-free Chronic Kidney Disease Epidemiology, EKFC_{RF}: European Kidney Function Consortium with race-free Q-values. EKFC_{PS}: European Kidney Function with population specific Q-values

Bias (A) and accuracy within 30% (P30) (B) for the CKD-EPI₂₀₂₁, the EKFC_{RF} and EKFC_{PS} in Black men (n=1,703) according to age.



Legend: CKD-EPI₂₀₂₁: race-free Chronic Kidney Disease Epidemiology, EKFC_{RF}: European Kidney Function Consortium with race-free Q-values. EKFC_{PS}: European Kidney Function with population specific Q-values

Cystatin C

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Estimating Glomerular Filtration Rate from Serum Creatinine and Cystatin C

Lesley A. Inker, M.D., Christopher H. Schmid, Ph.D., Hocine Tighiouart, M.S.,
John H. Eckfeldt, M.D., Ph.D., Harold I. Feldman, M.D., Tom Greene, Ph.D.,
John W. Kusek, Ph.D., Jane Manzi, Ph.D., Frederick Van Lente, Ph.D.,
Yaping Lucy Zhang, M.S., Josef Coresh, M.D., Ph.D., and Andrew S. Levey, M.D.,
for the CKD-EPI Investigators*

Table 1. Characteristics of Study Participants, According to Data Set.*

Characteristic	Development and Internal Validation (N = 5352)	External Validation (N = 1119)	P Value
Age — yr	47±15	50±17	<0.001
Age group — no. (%)			
<40 yr	2008 (38)	357 (32)	<0.001
40–65 yr	2625 (49)	530 (47)	
>65 yr	719 (13)	232 (21)	
Male sex — no. (%)	3107 (58)	663 (59)	0.46
Black race — no. (%)†	2123 (40)	30 (3)	<0.001
Diabetes — no. (%)	1726 (32)	594 (53)	<0.001
Body-mass index‡			
Mean	28±6	25±4	<0.001
<20 — no. (%)	214 (4)	81 (7)	<0.001
20–24 — no. (%)	1585 (30)	503 (45)	
25–30 — no. (%)	1881 (35)	386 (35)	
>30 — no. (%)	1671 (31)	149 (13)	
Mean weight — kg	83±20	74±15	<0.001
Mean height — cm	171±10	170±9	0.017
Mean body-surface area — m ²	1.94±0.24	1.85±0.21	<0.001
Mean serum cystatin C — ml/liter	1.4±0.7	1.5±0.8	0.01
Mean serum creatinine — mg/dl§	1.6±0.9	1.6±1.1	0.15
Mean measured GFR — ml/min/1.73 m ² of body-surface area	68±39	70±41	0.13
Measured GFR — no. (%)			
<15 ml/min/1.73 m ²	160 (3)	51 (5)	<0.001
15–29 ml/min/1.73 m ²	785 (15)	166 (15)	
30–59 ml/min/1.73 m ²	1765 (33)	316 (28)	
60–89 ml/min/1.73 m ²	1105 (21)	215 (19)	
90–119 ml/min/1.73 m ²	862 (16)	199 (18)	
>120 ml/min/1.73 m ²	675 (13)	172 (15)	

Table 2. Creatinine Equation (CKD-EPI 2009), Cystatin C Equation (CKD-EPI 2012), and Creatinine–Cystatin C Equation (CKD-EPI 2012) for Estimating GFR, Expressed for Specified Sex, Serum Creatinine Level, and Serum Cystatin C Level.*

Basis of Equation and Sex	Serum Creatinine†	Serum Cystatin C	Equation for Estimating GFR
	mg/dl	mg/liter	
CKD-EPI creatinine equation‡			
Female	≤0.7		$144 \times (\text{Scr}/0.7)^{-0.329} \times 0.993^{\text{Age}} [\times 1.159 \text{ if black}]$
Female	>0.7		$144 \times (\text{Scr}/0.7)^{-1.209} \times 0.993^{\text{Age}} [\times 1.159 \text{ if black}]$
Male	≤0.9		$141 \times (\text{Scr}/0.9)^{-0.411} \times 0.993^{\text{Age}} [\times 1.159 \text{ if black}]$
Male	>0.9		$141 \times (\text{Scr}/0.9)^{-1.209} \times 0.993^{\text{Age}} [\times 1.159 \text{ if black}]$
CKD-EPI cystatin C equation§			
Female or male		≤0.8	$133 \times (\text{Scys}/0.8)^{-0.499} \times 0.996^{\text{Age}} [\times 0.932 \text{ if female}]$
Female or male		>0.8	$133 \times (\text{Scys}/0.8)^{-1.328} \times 0.996^{\text{Age}} [\times 0.932 \text{ if female}]$
CKD-EPI creatinine–cystatin C equation¶			
Female	≤0.7	≤0.8	$130 \times (\text{Scr}/0.7)^{-0.248} \times (\text{Scys}/0.8)^{-0.375} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$
		>0.8	$130 \times (\text{Scr}/0.7)^{-0.248} \times (\text{Scys}/0.8)^{-0.711} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$
Female	>0.7	≤0.8	$130 \times (\text{Scr}/0.7)^{-0.601} \times (\text{Scys}/0.8)^{-0.375} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$
		>0.8	$130 \times (\text{Scr}/0.7)^{-0.601} \times (\text{Scys}/0.8)^{-0.711} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$
Male	≤0.9	≤0.8	$135 \times (\text{Scr}/0.9)^{-0.207} \times (\text{Scys}/0.8)^{-0.375} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$
		>0.8	$135 \times (\text{Scr}/0.9)^{-0.207} \times (\text{Scys}/0.8)^{-0.711} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$
Male	>0.9	≤0.8	$135 \times (\text{Scr}/0.9)^{-0.601} \times (\text{Scys}/0.8)^{-0.375} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$
		>0.8	$135 \times (\text{Scr}/0.9)^{-0.601} \times (\text{Scys}/0.8)^{-0.711} \times 0.995^{\text{Age}} [\times 1.08 \text{ if black}]$

Table 3. Use of the CKD-EPI Creatinine Equation (2009), CKD-EPI Cystatin C Equation (2012), and CKD-EPI Creatinine–Cystatin C Equations (2012) in the External-Validation Data Set Comprising 1119 Participants.*

Variable	Estimated GFR			
	Overall	<60	60–89	≥90
	<i>ml/min/1.73 m² of body-surface area</i>			
Bias — median difference (95% CI)				
Creatinine equation	3.7 (2.8 to 4.6)	1.8 (1.1 to 2.5)	6.6 (3.5 to 9.2)	11.1 (8.0 to 12.5)
Cystatin C equation	3.4 (2.3 to 4.4)	0.4 (–0.5 to 1.4)	6.0 (4.6 to 8.5)	8.5 (6.5 to 11.2)
Creatinine–cystatin C equation	3.9 (3.2 to 4.5)	1.3 (0.5 to 1.8)	6.9 (5.0 to 8.9)	10.6 (9.5 to 12.7)
Average of creatinine and cystatin C†	3.5 (2.8 to 4.1)	0.4 (–0.3 to 0.8)	6.5 (4.6 to 8.4)	11.9 (9.9 to 13.9)
Precision — IQR of the difference (95% CI)				
Creatinine equation	15.4 (14.3 to 16.5)	10.0 (8.9 to 11.0)	19.6 (17.3 to 23.2)	25.0 (21.6 to 28.1)
Cystatin C equation	16.4 (14.8 to 17.8)	11.0 (10.0 to 12.4)	19.6 (16.1 to 23.1)	22.6 (18.8 to 26.3)
Creatinine–cystatin C equation	13.4 (12.3 to 14.5)	8.1 (7.3 to 9.1)	15.9 (13.9 to 18.1)	18.8 (16.8 to 22.5)
Average of creatinine and cystatin C equations†	13.9 (12.9 to 14.7)	7.9 (7.1 to 9.0)	15.8 (13.9 to 17.7)	18.6 (16.1 to 22.2)
Accuracy — % (95% CI)‡				
1–P ₃₀				
Creatinine equation	12.8 (10.9 to 14.7)	16.6 (13.6 to 19.7)	10.2 (6.4 to 14.2)	7.8 (5.1 to 11.0)
Cystatin C equation	14.1 (12.2 to 16.2)	21.4 (18.2 to 24.9)	12.7 (8.5 to 17.4)	2.2 (0.6 to 3.9)
Creatinine–cystatin C equation	8.5 (7.0 to 10.2)	13.3 (10.7 to 16.1)	5.3 (2.7 to 8.2)	2.3 (0.9 to 4.2)
Average of creatinine and cystatin C equations†	8.2 (6.7 to 9.9)	12.1 (9.5 to 14.8)	6.4 (3.6 to 9.7)	2.9 (1.3 to 4.9)
1–P ₂₀				
Creatinine equation	32.9 (30.1 to 35.7)	37.2 (33.1 to 41.2)	31.1 (25.1 to 37.4)	26.5 (21.7 to 31.4)
Cystatin C equation	33.0 (30.3 to 35.7)	42.1 (38.2 to 46.1)	29.3 (23.6 to 35.4)	19.4 (15.4 to 23.7)
Creatinine–cystatin C equation	22.8 (20.4 to 25.2)	28.6 (25.1 to 32.4)	17.8 (13.3 to 22.9)	16.2 (12.4 to 20.5)
Average of creatinine and cystatin C equations†	23.7 (21.3 to 26.1)	29.1 (25.7 to 32.8)	17.6 (13.2 to 22.4)	18.8 (14.6 to 23.2)

ORIGINAL ARTICLE

Cystatin C–Based Equation to Estimate GFR without the Inclusion of Race and Sex

H. Pottel, J. Björk, A.D. Rule, N. Ebert, B.O. Eriksen, L. Dubourg, E. Vidal-Petiot, A. Grubb, M. Hansson, E.J. Lamb, K. Littmann, C. Mariat, T. Melsom, E. Schaeffner, P.-O. Sundin, A. Åkesson, A. Larsson, E. Cavalier, J.B. Bukabau, E.K. Sumaili, E. Yayo, D. Monnet, M. Flamant, U. Nyman, and P. Delanaye


ABSTRACT

N Engl J Med 2023;388:333-43.

First step: cystatin C and age

Laboratory data from
Sweden
N=227,643
♀ 95,469
♂ 132,174

Figure S3. Cystatin C versus age and the median quantile line for the 227,643 included subjects.




$$\begin{aligned} \text{♀ } Q' &= 0.79 \text{ mg/L until 50 y,} \\ & Q' = 0.79 + 0.005 \times (\text{Age} - 50) \\ \text{♂ } Q' &= 0.86 \text{ mg/L until 50 y} \\ & Q' = 0.86 + 0.005 \times (\text{Age} - 50) \end{aligned}$$

First step: cystatin C and **sex**

Laboratory data from
Sweden
N=227,643
♀ 95,469
♂ 132,174


Figure S3. Cystatin C versus age and the median quantile line for the 227,643 included subjects.



$$\begin{aligned} \text{♀ } Q' &= 0.79 \text{ mg/L until 50 y,} \\ & Q' = 0.79 + 0.005 \times (\text{Age} - 50) \\ \text{♂ } Q' &= 0.86 \text{ mg/L until 50 y} \\ & Q' = 0.86 + 0.005 \times (\text{Age} - 50) \end{aligned}$$

Second step: cystatin C and sex

Figure S4. Median plasma cystatin C in one-year intervals against age for men and women. A mathematical model to define Q'-values is proposed (red solid line): for adults Q' = 0.79 mg/L (women, dashed line) and 0.86 mg/L (men, solid line) until 50 years and a linear increasing model thereafter.



$$Q' = 0.83 \text{ mg/L until 50 years}$$
$$Q' = 0.83 + 0.005 \times (\text{Age} - 50)$$

Third step: Cystatin C and race

- Data from the same center in France
- Same method for GFR (Cr-EDTA), creatinine and cystatin C measurements
- Black and White people

Table S3. Patient characteristics of the entire cohorts used for the matching analysis (mean \pm SD)

Ethnicity/Sex	N	Age (years)	BMI (kg/m ²)	mGFR (mL/min/1.73m ²)	SCr (mg/dL)	CysC (mg/L)
White Men	1296 (57%)	53.0 \pm 14.6	26.2 \pm 4.9	61.8 \pm 26.0	1.52 \pm 0.73	1.52 \pm 0.68
Black Men	436 (63%)	50.7 \pm 13.1	26.3 \pm 4.5	62.0 \pm 22.1	1.73 \pm 0.81	1.41 \pm 0.61
White Women	966 (43%)	52.5 \pm 15.2	25.8 \pm 6.2	62.8 \pm 26.8	1.16 \pm 0.61	1.38 \pm 0.73
Black Women	261 (37%)	51.9 \pm 15.2	27.4 \pm 5.8	59.1 \pm 25.6	1.40 \pm 0.79	1.46 \pm 0.76

Third step: Cystatin C and race

Matched analysis 1:1 for

- sex
- BMI ($\pm 2,5$ kg/m²)
- Measured GFR (± 3 mL/min/1.73m²)
- age (± 3 y)

Table S4. Demographic and renal characteristics of the matched White and Black subjects (mean \pm SD)

Sex	N	Age (years)	BMI (kg/m ²)	mGFR (mL/min/1.73m ²)	SCr (mg/dL)	CysC (mg/L)
White Men	377	51.1 \pm 12.2	25.7 \pm 3.4	63.8 \pm 21.0	1.43 \pm 0.62	1.41 \pm 0.56
Black Men	377	50.8 \pm 12.3	25.8 \pm 3.5	63.6 \pm 21.0	1.65 \pm 0.64	1.37 \pm 0.59
White Women	200	53.4 \pm 11.9	26.1 \pm 4.6	59.7 \pm 23.2	1.16 \pm 0.53	1.40 \pm 0.69
Black Women	200	53.3 \pm 11.9	26.2 \pm 4.6	59.8 \pm 23.1	1.33 \pm 0.61	1.41 \pm 0.64

Fourth Step: Validation of the new equation

$$\text{EKFC} - \text{eGFR} = 107.3 / [\text{Biomarker}/\text{Q}]^\alpha \times \\ [0.990^{(\text{Age}-40)} \text{ if age } > 40 \text{ years}],$$

with $\alpha = 0.322$ when biomarker/Q is less than 1
and $\alpha = 1.132$ when biomarker/Q is 1 or more.

Adults

Measured GFR, IDMS traceable creatinine, calibrated cystatin C
N=12,832

11 cohorts

White Europeans: n=7,727

White Europeans from Paris: n=2,646

White US: n=1,093

Black Europeans from Paris: n=858

Black Africans: n=508

Table 1. Performance of Single Biomarker (Serum Creatinine or Cystatin C)–Based Equations to Estimate the Glomerular Filtration Rate.*

Variable	Serum Creatinine–Based Equations		
	CKD-EPI eGFR _{cr} (ASR)	CKD-EPI eGFR _{cr} (AS)	EKFC eGFR _{cr}
EKFC cohort, 7727 White patients			
Median bias (95% CI) — ml/min/1.73 m ² †	3.96 (3.67 to 4.32)	7.40 (7.02 to 7.76)	0.58 (0.32 to 0.86)
IQR of estimated GFR– measured GFR— ml/min/1.73 m ² ‡	15.5 (–3.0 to 12.5)	16.3 (0.0 to 16.3)	14.5 (–6.5 to 8.0)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	14.8 (14.4 to 15.2)	16.3 (15.9 to 16.6)	13.1 (12.8 to 13.4)
P ₃₀ — % (95% CI)¶	40.3 (39.2 to 41.4)	34.7 (33.6 to 35.8)	43.3 (42.2 to 44.4)
P ₉₀ — % (95% CI)‖	81.6 (80.8 to 82.5)	75.7 (74.8 to 76.7)	85.8 (85.0 to 86.5)

7.40 (7.02 to 7.76)	0.58 (0.32 to 0.86)
16.3 (0.0 to 16.3)	14.5 (–6.5 to 8.0)
16.3 (15.9 to 16.6)	13.1 (12.8 to 13.4)
34.7 (33.6 to 35.8)	43.3 (42.2 to 44.4)
75.7 (74.8 to 76.7)	85.8 (85.0 to 86.5)

Table 1. Performance of Single Biomarker (Serum Creatinine or Cystatin C)-Based Equations to Estimate the Glomerular Filtration Rate.*

Variable	Cystatin C-Based Equations	
	CKD-EPI eGFR _{cys}	EKFC eGFR _{cys} without Sex
EKFC cohort, 7727 White patients		
Median bias (95% CI) — ml/min/1.73 m ² †	0.28 (-0.02 to 0.64)	0.00 (-0.37 to 0.27)
IQR of estimated GFR- measured GFR— ml/min/1.73 m ² ‡	19.1 (-7.9 to 11.2)	14.4 (-7.9 to 6.5)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	15.8 (15.5 to 16.1)	13.5 (12.9 to 14.1)
P ₃₀ — % (95% CI)¶	32.0 (31.0 to 33.0)	41.7 (40.6 to 42.8)
P ₉₀ — % (95% CI)‖	80.8 (79.9 to 81.7)	86.2 (85.4 to 87.0)

Cystatin C-Based Equations

CKD-EPI eGFR _{cys}	EKFC eGFR _{cys} without Sex
0.28 (-0.02 to 0.64)	0.00 (-0.37 to 0.27)
19.1 (-7.9 to 11.2)	14.4 (-7.9 to 6.5)
15.8 (15.5 to 16.1)	13.5 (12.9 to 14.1)
32.0 (31.0 to 33.0)	41.7 (40.6 to 42.8)
80.8 (79.9 to 81.7)	86.2 (85.4 to 87.0)

Table 1. Performance of Single Biomarker (Serum Creatinine or Cystatin C)–Based Equations to Estimate the Glomerular Filtration Rate.*

Variable	Serum Creatinine–Based Equations			Cystatin C–Based Equations	
	CKD-EPI eGFR _{cr} (ASR)	CKD-EPI eGFR _{cr} (AS)	EKFC eGFR _{cr}	CKD-EPI eGFR _{cys}	EKFC eGFR _{cys} without Sex
EKFC cohort, 7727 White patients					
Median bias (95% CI) — ml/min/1.73 m ² †	3.96 (3.67 to 4.32)	7.40 (7.02 to 7.76)	0.58 (0.32 to 0.86)	0.28 (–0.02 to 0.64)	0.00 (–0.37 to 0.27)
IQR of estimated GFR– measured GFR— ml/min/1.73 m ² ‡	15.5 (–3.0 to 12.5)	16.3 (0.0 to 16.3)	14.5 (–6.5 to 8.0)	19.1 (–7.9 to 11.2)	14.4 (–7.9 to 6.5)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	14.8 (14.4 to 15.2)	16.3 (15.9 to 16.6)	13.1 (12.8 to 13.4)	15.8 (15.5 to 16.1)	13.5 (12.9 to 14.1)
P ₃₀ — % (95% CI)¶	40.3 (39.2 to 41.4)	34.7 (33.6 to 35.8)	43.3 (42.2 to 44.4)	32.0 (31.0 to 33.0)	41.7 (40.6 to 42.8)
P ₉₀ — % (95% CI)‖	81.6 (80.8 to 82.5)	75.7 (74.8 to 76.7)	85.8 (85.0 to 86.5)	80.8 (79.9 to 81.7)	86.2 (85.4 to 87.0)

Table 1. Performance of Single Biomarker (Serum Creatinine or Cystatin C)-Based Equations to Estimate the Glomerular Filtration Rate.*

Variable	Serum Creatinine-Based Equations			Cystatin C-Based Equations	
	CKD-EPI eGFR _{cr} (AS)	CKD-EPI eGFR _{cr} (AS)	EKFC eGFR _{cr}	CKD-EPI eGFR _{cys}	EKFC eGFR _{cys} without Sex
U.S. cohort, 1093 White patients					
Median bias (95% CI) — ml/min/1.73 m ² †	2.83 (1.65 to 3.64)	7.11 (6.15 to 7.97)	-2.69 (-3.68 to -1.79)	12.1 (11.1 to 13.3)	4.26 (3.33 to 5.08)
IQR of estimated GFR- measured GFR— ml/min/1.73 m ² ‡	18.8 (-6.6 to 12.2)	18.7 (-2.2 to 16.5)	18.5 (-11.9 to 6.6)	21.5 (1.5 to 23.0)	18.3 (-5.3 to 13.0)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	16.2 (15.0 to 17.4)	17.5 (16.4 to 18.6)	16.2 (14.9 to 17.4)	21.3 (20.3 to 22.2)	16.8 (15.7 to 17.9)
P ₃₀ — % (95% CI)¶	41.9 (39.0 to 44.8)	39.4 (36.5 to 42.3)	41.4 (38.5 to 44.4)	29.4 (26.7 to 32.1)	41.5 (38.6 to 44.5)
P ₃₀ — % (95% CI)	86.0 (83.9 to 88.1)	81.0 (78.6 to 83.3)	89.3 (87.5 to 91.1)	72.9 (70.3 to 75.6)	83.9 (81.7 to 86.1)

Table 2. Performance of Combined Serum Creatinine– and Cystatin C–Based Equations to Estimate GFR.*

Variable	CKD-EPI eGFRcr-cys(ASR)	CKD-EPI eGFRcr-cys(AS)	EKFC eGFRcr-cys without Sex
EKFC cohort, 7727 White patients			
Median bias (95% CI) — ml/min/1.73 m ² †	2.50 (2.17 to 2.76)	5.04 (4.69 to 5.36)	0.37 (0.14 to 0.66)
IQR of estimated GFR – measured GFR — ml/min/1.73 m ² ‡	14.8 (–3.6 to 11.2)	16.7 (–1.8 to 14.9)	12.0 (–5.9 to 6.1)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	13.1 (12.8 to 13.4)	14.7 (14.4 to 15.0)	11.3 (11.0 to 11.6)
P ₁₀ — % (95% CI) ¶	41.5 (40.4 to 42.6)	37.2 (36.2 to 38.3)	48.9 (47.8 to 50.0)
P ₃₀ — % (95% CI)	88.3 (87.6 to 89.0)	84.2 (83.4 to 85.0)	90.4 (89.8 to 91.1)
Paris cohort, 2646 White patients			
Median bias (95% CI) — ml/min/1.73 m ² †	–1.35 (–1.82 to –0.97)	0.64 (0.16 to 1.15)	–0.65 (–1.06 to –0.23)
IQR of estimated GFR – measured GFR — ml/min/1.73 m ² ‡	13.4 (–7.5 to 5.8)	14.1 (–5.8 to 8.3)	12.4 (–6.8 to 5.6)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	12.1 (11.6 to 12.7)	12.6 (12.0 to 13.1)	11.8 (11.2 to 12.4)
P ₁₀ — % (95% CI) ¶	43.9 (42.0 to 45.8)	42.3 (40.4 to 44.1)	45.8 (43.9 to 47.7)
P ₃₀ — % (95% CI)	89.7 (88.5 to 90.8)	89.2 (88.0 to 90.4)	92.1 (91.1 to 93.1)
U.S. cohort, 1093 White patients			
Median bias (95% CI) — ml/min/1.73 m ² †	9.23 (8.45 to 10.10)	13.9 (13.1 to 14.9)	0.97 (0.01 to 2.12)
IQR of estimated GFR – measured GFR — ml/min/1.73 m ² ‡	18.4 (0.5 to 18.8)	18.1 (5.1 to 23.3)	17.4 (–8.2 to 9.2)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	18.1 (17.1 to 19.1)	21.0 (20.1 to 22.0)	15.5 (14.3 to 16.7)
P ₁₀ — % (95% CI) ¶	37.1 (34.3 to 40.0)	28.1 (25.4 to 30.8)	45.7 (42.7 to 48.6)
P ₃₀ — % (95% CI)	79.5 (77.1 to 81.9)	72.1 (69.4 to 74.8)	88.7 (86.9 to 90.6)
Paris cohort, 858 Black patients			
Median bias (95% CI) — ml/min/1.73 m ² †	–0.37 (–1.06 to 0.57)	–2.08 (–2.71 to –1.32)	–0.65 (–1.23 to 0.11)
IQR of estimated GFR – measured GFR — ml/min/1.73 m ² ‡	15.2 (–6.4 to 8.8)	14.0 (–7.9 to 6.1)	12.4 (–6.2 to 6.2)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	13.3 (11.9 to 14.6)	12.6 (11.2 to 13.9)	11.6 (10.0 to 13.0)
P ₁₀ — % (95% CI) ¶	38.7 (35.4 to 42.0)	38.9 (35.7 to 42.2)	48.3 (44.9 to 51.6)
P ₃₀ — % (95% CI)	87.9 (85.7 to 90.1)	89.0 (87.0 to 91.1)	92.0 (90.1 to 93.8)
African cohort, 508 Black patients			
Median bias (95% CI) — ml/min/1.73 m ² †	8.55 (6.87 to 10.30)	4.08 (2.37 to 5.78)	0.42 (–1.03 to 1.51)
IQR of estimated GFR – measured GFR — ml/min/1.73 m ² ‡	24.7 (–4.5 to 20.1)	22.0 (–7.4 to 14.7)	17.1 (–7.2 to 10.0)
Root-mean-square error (95% CI) — ml/min/1.73 m ² §	19.7 (18.2 to 21.1)	17.2 (15.8 to 18.5)	14.7 (13.3 to 16.0)
P ₁₀ — % (95% CI) ¶	28.7 (24.8 to 32.7)	34.3 (30.1 to 38.4)	43.5 (39.2 to 47.8)
P ₃₀ — % (95% CI)	75.0 (71.2 to 78.8)	77.6 (73.9 to 81.2)	84.3 (81.1 to 87.4)

Cystatin C/EKFC

- Cystatin C allows an eGFR without race nor sex
- EKFC is mathematically the same as EKFC creatinine, only Q is changing
- Continuum between children and adults for $EKFC_{crea}$
- Equations based on cystatin C are not better than equations based on creatinine
- EKFC equations are slightly better than corresponding CKD-EPI equations => **good alternative to CKD-Epi in Europe and Africa**
- Combined equations are better (P30 +5-10%)
- Standardisation
- More costly
- How to manage discrepant results?
- Place of EKFC and/or cystatin C in the next KDIGO?

<https://ekfccalculator.pages.dev/>

Limitations of equations = creatinine

Specific populations:
Equations are not magic!
Keep our clinical feeling!!

Anorexia Nervosa (Delanaye P, Clin Nephrol, 2009, 71, 482)

Cirrhosis (Skruzacek PA, Am J Kidney Dis, 2003, 42, 1169)

ICU (Delanaye P, BMC Nephrology, 2014, 15, 9)

Hospitalized (Poggio ED, Am J Kidney Dis, 2005, 46, 242)

Heart Transplanted (Delanaye P, Clin Transplant, 2006, 20, 596)

Kidney Transplanted (Masson I, Transplantation, 2013, 95, 1211)

Obesity (Bouquegneau A, NDT, 2013, 28, iv122)

Do not over-interpret an eGFR result...

All equations remain estimation...

Good at the population level

Lack of precision at the individual level




Fig. 1. Uncertainty of eGFR calculated using the CKD-EPI equations for African-Americans and non-African-Americans at various creatinine concentrations for a 50-year-old male. Circles (red, larger values) indicate African-American and diamonds (green, lower values) indicate non-African-American equations. Plot symbols are the eGFR values and error bars represent the 95% CI for each eGFR value.

$$eGFR = 60,25 \text{ ml/min/1.73m}^2$$




Fig. 1. Uncertainty of eGFR calculated using the CKD-EPI equations for African-Americans and non-African-Americans at various creatinine concentrations for a 50-year-old male. Circles (red, larger values) indicate African-American and diamonds (green, lower values) indicate non-African-American equations. Plot symbols are the eGFR values and error bars represent the 95% CI for each eGFR value.

$$\begin{aligned}
 \text{eGFR} &= \cancel{60,25} \text{ ml/min/1.73m}^2 \\
 &= 60 \text{ ml/min/1.73m}^2 \quad (\text{CI } 95\%: 33-87)
 \end{aligned}$$

The applicability of eGFR equations to different populations

Pierre Delanaye and Christophe Mariat



GO BACK to MEASURED GFR

Delanaye P, Nature Rev Nephrol, 2013, 9, p513

Ebert N, Clin Kidney J, 2021, 14, p1861

Agarwal R, Nephrol Dial Transplant, 2019, 34, p2001

Shafi T, Ann Intern Med, 2022, 175, p1073

Thanks for your attention



« The Cat » a Belgian Comic star...