
Cinquième partie :

Références générales

-
- ADAMS L.M., SIMMONS C.P., REZMANN L., STRUGNELL R.A., ROBINS-BROWNE R.M. Identification and characterization of a K88- and CS31A-like operon of a rabbit enteropathogenic *Escherichia coli* strain which encodes fimbriae involved in the colonization of rabbit intestine. *Infect. Immun.*, 1997, **65**, 5222-5230.
- ADU-BOBIE J., FRANKEL G., BAIN C., GONCALVES A.G., TRABULSI L.R., DOUCE G., KNUTTON S., DOUGAN G. Detection of intimins alpha, beta, gamma, and delta, four intimin derivatives expressed by attaching and effacing microbial pathogens. *J. Clin. Microbiol.*, 1998, **36**, 662-668.
- AN H., FAIRBROTHER J.M., DESAUTELS C., HAREL J. Distribution of a novel locus called Paa (porcine attaching and effacing associated) among enteric *Escherichia coli*. *Adv. Exp. Med. Biol.*, 1999, **473**, 179-184.
- ANONYMOUS Verotoxigenic *Escherichia coli*: Verotoxigenic *E. coli*; In Trends and Sources of Zoonotic Agents in Animals, Feeding Stuffs, Food and Man in the European Union and Norway in 2002. [en ligne] (25/10/2006) Adresse URL : http://ec.europa.eu/food/food/biosafety/salmonella/zoonoses_reps_2002_en.htm.
- BALDINI M.M., KAPER J.B., LEVINE M.M., CANDY D.C., MOON H.W. Plasmid-mediated adhesion in enteropathogenic *Escherichia coli*. *J. Pediatr. Gastroenterol. Nutr.*, 1983, **2**, 534-538.
- BATISSON I., GUIMOND M.P., GIRARD F., AN H., ZHU C., OSWALD E., FAIRBROTHER J.M., JACQUES M., HAREL J. Characterization of the novel factor paa involved in the early steps of the adhesion mechanism of attaching and effacing *Escherichia coli*. *Infect. Immun.*, 2003, **71**, 4516-4525.
- BAUER M.E., WELCH R.A. Characterization of an RTX toxin from enterohemorrhagic *Escherichia coli* O157:H7. *Infect. Immun.*, 1996, **64**, 167-175.
- BAUMLER A.J., HEFFRON F. Identification and sequence analysis of *lpfABCDE*, a putative fimbrial operon of *Salmonella typhimurium*. *J. Bacteriol.*, 1995, **177**, 2087-2097.
- BERENDSON R., CHENEY C.P., SCHAD P.A., BOEDEKER E.C. Species-specific binding of purified pili (AF/R1) from the *Escherichia coli* RDEC-1 to rabbit intestinal mucosa. *Gastroenterology*, 1983, **85**, 837-845.
- BERGAN T. Classification of *Escherichia coli*. *Methods Microbiol.*, 1984, **14**, 1-41.
- BERTSCHINGER H.U., NIEF V., TSCHAPE H. Active oral immunization of suckling piglets to prevent colonization after weaning by enterotoxigenic *Escherichia coli* with fimbriae F18. *Vet. Microbiol.*, 2000, **71**, 255-267.
- BETTELHEIM, K. A. Biochemical characteristics of *Escherichia coli*. In : Gyles, C. L. (Ed.), *Escherichia coli* in domestic animals and humans. Cab International : Wallingford, 1994, 3-30.
- BEUTIN L. *Escherichia coli* as a pathogen in dogs and cats. *Vet. Res.*, 1999, **30**, 285-298.
-

- BEUTIN, L. Plasmids in VTEC: their role in virulence and their use in typing. In : Duffy, G., Garvey, P., Coia, J., Wasteson, Y., and Mc Dowell, D. A. Eds (Ed.), Verocytotoxigenic *Escherichia coli* in Europe. 3. Pathogenicity and virulence (Proceedings). Concerted Action CT98-3935. The National Food Center : Dublin, 2000, 126-139.
- BEUTIN L., ALEKSIC S., ZIMMERMANN S., GLEIER K. Virulence factors and phenotypical traits of verotoxigenic strains of *Escherichia coli* isolated from human patients in Germany. *Med. Microbiol. Immunol. (Berl)*, 1994, **183**, 13-21.
- BIELASZEWSKA M., ZHANG W., TARR P.I., SONNTAG A.K., KARCH H. Molecular profiling and phenotype analysis of *Escherichia coli* O26:H11 and O26:NM: secular and geographic consistency of enterohemorrhagic and enteropathogenic isolates. *J. Clin. Microbiol.*, 2005, **43**, 4225-4228.
- BLANCO J.E., BLANCO M., ALONSO M.P., MORA A., DAHBI G., COIRA M.A., BLANCO J. Serotypes, virulence genes, and intimin types of Shiga toxin (verotoxin)-producing *Escherichia coli* isolates from human patients: prevalence in Lugo, Spain, from 1992 through 1999. *J. Clin. Microbiol.*, 2004, **42**, 311-319.
- BOERLIN P., MCEWEN S.A., BOERLIN-PETZOLD F., WILSON J.B., JOHNSON R.P., GYLES C.L. Associations between virulence factors of Shiga toxin-producing *Escherichia coli* and disease in humans. *J. Clin. Microbiol.*, 1999, **37**, 497-503.
- BOSWORTH B.T., DEAN-NYSTROM E.A., CASEY T.A., NEIBERGS H.L. Differentiation of F18ab+ from F18ac+ *Escherichia coli* by single-strand conformational polymorphism analysis of the major fimbrial subunit gene (*fedA*). *Clin. Diagn. Lab Immunol.*, 1998, **5**, 299-302.
- BOUVET J., MONTET M.P., ROSSEL R., LE ROUX A., BAVAI C., RAY-GUENIOT S., MAZUY C., ATRACHE V., VERNOSY-ROZAND C. Prevalence of verotoxin-producing *Escherichia coli* (VTEC) and *E. coli* O157:H7 in French pork. *J. Appl. Microbiol.*, 2002, **93**, 7-14.
- BURLAND V., SHAO Y., PERNA N.T., PLUNKETT G., SOFIA H.J., BLATTNER F.R. The complete DNA sequence and analysis of the large virulence plasmid of *Escherichia coli* O157:H7. *Nucleic Acids Res.*, 1998, **26**, 4196-4204.
- CAMPOS L.C., FRANZOLIN M.R., TRABULSI L.R. Diarrheagenic *Escherichia coli* categories among the traditional enteropathogenic *E. coli* O serogroups--a review. *Mem. Inst. Oswaldo Cruz*, 2004, **99**, 545-552.
- CANTEY J.R., BLAKE R.K., WILLIFORD J.R., MOSELEY S.L. Characterization of the *Escherichia coli* AF/R1 pilus operon: novel genes necessary for transcriptional regulation and for pilus-mediated adherence. *Infect. Immun.*, 1999, **67**, 2292-2298.
- CAPRIOLI A., MORABITO S., BRUGEREB H., OSWALD E. Enterohaemorrhagic *Escherichia coli*: emerging issues on virulence and modes of transmission. *Vet. Res.*, 2005, **36**, 289-311.

- CHALMERS R.M., SALMON R.L., WILLSHAW G.A., CHEASTY T., LOOKER N., DAVIES I., WRAY C. Vero-cytotoxin-producing *Escherichia coli* O157 in a farmer handling horses. *Lancet*, 1997, **349**, 1816-1816.
- CHAPMAN T.A., WU X.Y., BARCHIA I., BETTELHEIM K.A., DRIESEN S., TROTT D., WILSON M., CHIN J.J. Comparison of virulence gene profiles of *Escherichia coli* strains isolated from healthy and diarrheic swine. *Appl. Environ. Microbiol.*, 2006, **72**, 4782-4795.
- CHEN H.D., FRANKEL G. Enteropathogenic *Escherichia coli*: unravelling pathogenesis. *FEMS Microbiol. Rev.*, 2005, **29**, 83-98.
- CHENG D., SUN H., XU J., GAO S. Prevalence of fimbrial colonization factors F18ab and F18ac in *Escherichia coli* isolates from weaned piglets with edema and/or diarrhea in China. *Vet. Microbiol.*, 2005, **110**, 35-39.
- CHINA B. Attaching and effacing *Escherichia coli*. *Recent Res. Devel. Microbiology*, 2000, **4**, 1-30.
- CHINA B., GOFFAUX F. Secretion of virulence factors by *Escherichia coli*. *Vet. Res.*, 1999, **30**, 181-202
- CHINA B., PIRSON V., MAINIL J. Prevalence and molecular typing of attaching and effacing *Escherichia coli* among calf populations in Belgium. *Vet. Microbiol.*, 1998, **63**, 249-259.
- CLARKE S.C. Diarrhoeagenic *Escherichia coli*--an emerging problem? *Diagn. Microbiol. Infect. Dis.*, 2001, **41**, 93-98.
- CLARKE S.C., HAIGH R.D., FREESTONE P.P., WILLIAMS P.H. Virulence of enteropathogenic *Escherichia coli*, a global pathogen. *Clin. Microbiol. Rev.*, 2003, **16**, 365-378.
- CLEARY J., LAI L.C., SHAW R.K., STRAATMAN-IWANOWSKA A., DONNENBERG M.S., FRANKEL G., KNUTTON S. Enteropathogenic *Escherichia coli* (EPEC) adhesion to intestinal epithelial cells: role of bundle-forming pili (BFP), EspA filaments and intimin. *Microbiology*, 2004, **150**, 527-538.
- CORNICK N.A., HELGERSON A.F., MAI V., RITCHIE J.M., ACHESON D.W. In vivo transduction of an Stx-encoding phage in ruminants. *Appl. Environ. Microbiol.*, 2006, **72**, 5086-5088.
- CRAVIOTO A., GROSS R.J., SCOTLAND S.M., ROWE B. An adhesive factor found in strains of *Escherichia coli* belonging to the traditional infantile enteropathogenic serogroups. *Current. Microbiol.*, 1979, **3**, 95-99.
- CRAVIOTO A., TELLO A., NAVARRO A., RUIZ J., VILLAFAN H., URIBE F., ESLAVA C. Association of *Escherichia coli* HEp-2 adherence patterns with type and duration of diarrhoea. *Lancet*, 1991, **337**, 262-264.

- DEGRANDIS S., LAW H., BRUNTON J., GYLES C., LINGWOOD C.A. Globotetraosylceramide is recognized by the pig edema disease toxin. *J. Biol. Chem.*, 1989, **264**, 12520-12525.
- D'SOUZA J.M., WANG L., REEVES P. Sequence of the Escherichia coli O26 O antigen gene cluster and identification of O26 specific genes. *Gene*, 2002, **297**, 123-127.
- DEVINNEY R., GAUTHIER A., ABE A., FINLAY B.B. Enteropathogenic *Escherichia coli*: a pathogen that inserts its own receptor into host cells. *Cell Mol. Life Sci.*, 1999, **55**, 961-976.
- DONNENBERG M.S. (2006) [en ligne] (09/07/2006) Adresse URL : <http://medschool.umaryland.edu/infeMSD/Images.htm>, consulté le 30/04/2007.
- DONNENBERG M.S., GIRON J.A., NATARO J.P., KAPER J.B. A plasmid-encoded type IV fimbrial gene of enteropathogenic *Escherichia coli* associated with localized adherence. *Mol. Microbiol.*, 1992, **6**, 3427-3437.
- DOUGHTY S., SLOAN J., BENNETT-WOOD V., ROBERTSON M., ROBINS-BROWNE R.M., HARTLAND E.L. Identification of a novel fimbrial gene cluster related to long polar fimbriae in locus of enterocyte effacement-negative strains of enterohemorrhagic *Escherichia coli*. *Infect. Immun.*, 2002, **70**, 6761-6769.
- DOWNES F.P., BARRETT T.J., GREEN J.H., ALOISIO C.H., SPIKA J.S., STROCKBINE N.A., WACHSMUTH I.K. Affinity purification and characterization of Shiga-like toxin II and production of toxin-specific monoclonal antibodies. *Infect. Immun.*, 1988, **56**, 1926-1933.
- ELLIOTT S.J., KAPER J.B. Role of type 1 fimbriae in EPEC infections. *Microb. Pathog.*, 1997, **23**, 113-118.
- ELLIOTT S.J., WAINWRIGHT L.A., MCDANIEL T.K., JARVIS K.G., DENG Y.K., LAI L.C., MCNAMARA B.P., DONNENBERG M.S., KAPER J.B. The complete sequence of the locus of enterocyte effacement (LEE) from enteropathogenic *Escherichia coli* E2348/69. *Mol. Microbiol.*, 1998, **28**, 1-4.
- ESCHERICH T. Die Darmbakterien des Neugeborenen un Saglins. *Fortschr. Med.*, 1885, **3**, 515-522.
- FARMER J.J., III, DAVIS B.R., HICKMAN-BRENNER F.W., MCWHORTER A., HUNTLEY-CARTER G.P., ASBURY M.A., RIDDLE C., WATHEN-GRADY H.G., ELIAS C., FANNING G.R., . Biochemical identification of new species and biogroups of *Enterobacteriaceae* isolated from clinical specimens. *J. Clin. Microbiol.*, 1985, **21**, 46-76.
- FEKETE P.Z., GERARDIN J., JACQUEMIN E., MAINIL J.G., NAGY B. Replicon typing of F18 fimbriae encoding plasmids of enterotoxigenic and verotoxigenic *Escherichia coli* strains from porcine postweaning diarrhoea and oedema disease. *Vet. Microbiol.*, 2002, **85**, 275-284.

-
- FIEDERLING F., BOURY M., PETIT C., MILON A. Adhesive factor/rabbit 2, a new fimbrial adhesin and a virulence factor from *Escherichia coli* O103, a serogroup enteropathogenic for rabbits. *Infect. Immun.*, 1997, **65**, 847-851.
- FRANKEL G., PHILLIPS A.D., ROSENSHINE I., DOUGAN G., KAPER J.B., KNUTTON S. Enteropathogenic and enterohaemorrhagic *Escherichia coli*: more subversive elements. *Mol. Microbiol.*, 1998, **30**, 911-921.
- FRIEDRICH A.W., BIELASZEWSKA M., ZHANG W.L., PULZ M., KUCZIUS T., AMMON A., KARCH H. *Escherichia coli* harboring Shiga toxin 2 gene variants: frequency and association with clinical symptoms. *J. Infect. Dis.*, 2002, **185**, 74-84.
- GIRON J.A., HO A.S., SCHOOLNIK G.K. An inducible bundle-forming pilus of enteropathogenic *Escherichia coli*. *Science*, 1991, **254**, 710-713.
- GIRON J.A., HO A.S., SCHOOLNIK G.K. Characterization of fimbriae produced by enteropathogenic *Escherichia coli*. *J. Bacteriol.*, 1993, **175**, 7391-7403.
- GIRON J.A., TORRES A.G., FREER E., KAPER J.B. The flagella of enteropathogenic *Escherichia coli* mediate adherence to epithelial cells. *Mol. Microbiol.*, 2002, **44**, 361-379.
- GOFFAUX F., CHINA B., JANSSEN L., MAINIL J. Genotypic characterization of enteropathogenic *Escherichia coli* (EPEC) isolated in Belgium from dogs and cats. *Res. Microbiol.*, 2000, **151**, 865-871.
- GOFFAUX F., CHINA B., MAINIL J. Organisation and in vitro expression of *esp* genes of the LEE (locus of enterocyte effacement) of bovine enteropathogenic and enterohemorrhagic *Escherichia coli*. *Vet. Microbiol.*, 2001, **83**, 275-286.
- GOFFAUX F., JANSSEN L., PIRSON V., CHINA B., MAINIL J. Comparaison des *Escherichia coli* entéropathogènes (ECEP) et vérotoxino-gènes (ECVT) humaines et animales par typage de l'îlot de pathogénicité de type III (locus LEE). *Le Médecin Vétérinaire du Québec*, 2001, **31**, 16-18.
- GOMES T.A., BLAKE P.A., TRABULSI L.R. Prevalence of *Escherichia coli* strains with localized, diffuse, and aggregative adherence to HeLa cells in infants with diarrhea and matched controls. *J. Clin. Microbiol.*, 1989, **27**, 266-269.
- GOMEZ-DUARTE O.G., KAPER J.B. A plasmid-encoded regulatory region activates chromosomal *aeiA* expression in enteropathogenic *Escherichia coli*. *Infect. Immun.*, 1995, **63**, 1767-1776.
- GRIFFIN P.M., TAUXE R.V. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli*, and the associated hemolytic uremic syndrome. *Epidemiol. Rev.*, 1991, **13**, 60-98.
- HABIB N.F., JACKSON M.P. Identification of a B subunit gene promoter in the Shiga toxin operon of *Shigella dysenteriae* 1. *J. Bacteriol.*, 1992, **174**, 6498-6507.
- HAREL J., MARTIN C. Virulence gene regulation in pathogenic *Escherichia coli*. *Vet. Res.*, 1999, **30**, 131-155.
-

-
- HAYASHI T., MAKINO K., OHNISHI M., KUROKAWA K., ISHII K., YOKOYAMA K., HAN C.G., OHTSUBO E., NAKAYAMA K., MURATA T., TANAKA M., TOBE T., IIDA T., TAKAMI H., HONDA T., SASAKAWA C., OGASAWARA N., YASUNAGA T., KUHARA S., SHIBA T., HATTORI M., SHINAGAWA H. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.*, 2001, **8**, 11-22.
- HEAD S.C., KARMALI M.A., LINGWOOD C.A. Preparation of VT1 and VT2 hybrid toxins from their purified dissociated subunits. Evidence for B subunit modulation of a subunit function. *J. Biol. Chem.*, 1991, **266**, 3617-3621.
- HEAD S.C., KARMALI M.A., ROSCOE M.E., PETRIC M., STROCKBINE N.A., WACHSMUTH I.K. Serological differences between verocytotoxin 2 and shiga-like toxin II. *Lancet*, 1988, **2**, 751-751.
- HERRERO M., DE LORENZO V., TIMMIS K.N. Transposon vectors containing non-antibiotic resistance selection markers for cloning and stable chromosomal insertion of foreign genes in gram-negative bacteria. *J. Bacteriol.*, 1990, **172**, 6557-6567.
- HICKS S., FRANKEL G., KAPER J.B., DOUGAN G., PHILLIPS A.D. Role of intimin and bundle-forming pili in enteropathogenic *Escherichia coli* adhesion to pediatric intestinal tissue in vitro. *Infect. Immun.*, 1998, **66**, 1570-1578.
- HOLLAND R.E. Some infectious causes of diarrhea in young farm animals. *Clin. Microbiol. Rev.*, 1990, **3**, 345-375.
- IMBERECHTS H., DE GREVE H., SCHLICKER C., BOUCHET H., POHL P., CHARLIER G., BERTSCHINGER H., WILD P., VANDEKERCKHOVE J., VAN DAMME J., . Characterization of F107 fimbriae of *Escherichia coli* 107/86, which causes edema disease in pigs, and nucleotide sequence of the F107 major fimbrial subunit gene, *fedA*. *Infect. Immun.*, 1992, **60**, 1963-1971.
- IMBERECHTS H., BERTSCHINGER H.U., NAGY B., DEPREZ P., POHL P. Fimbrial colonisation factors F18ab and F18ac of *Escherichia coli* isolated from pigs with postweaning diarrhea and edema disease. *Adv. Exp. Med. Biol.*, 1997, **412**, 175-183.
- JALLAT C., DARFEUILLE-MICHAUD A., RICH C., JOLY B. Survey of clinical isolates of diarrhoeogenic *Escherichia coli*: diffusely adhering *E. coli* strains with multiple adhesive factors. *Res. Microbiol.*, 1994, **145**, 621-632.
- JANKE B.H., FRANCIS D.H., COLLINS J.E., LIBAL M.C., ZEMAN D.H., JOHNSON D.D., NEIGER R.D. Attaching and effacing *Escherichia coli* infection as a cause of diarrhea in young calves. *J. Am. Vet. Med. Assoc.*, 1990, **196**, 897-901.
- JENKINS C., PERRY N.T., CHEASTY T., SHAW D.J., FRANKEL G., DOUGAN G., GUNN G.J., SMITH H.R., PATON A.W., PATON J.C. Distribution of the *saa* gene in strains of Shiga toxin-producing *Escherichia coli* of human and bovine origins. *J. Clin. Microbiol.*, 2003, **41**, 1775-1778.
- JENSEN C.O. Ueber die Kälberrhur und deren Aetiologie. *Mh. Tierheilk.*, 1893, **4**, 97-124.
-

- JERSE A.E., YU J., TALL B.D., KAPER J.B. A genetic locus of enteropathogenic *Escherichia coli* necessary for the production of attaching and effacing lesions on tissue culture cells. *Proc. Natl. Acad. Sci. U. S. A.*, 1990, **87**, 7839-7843.
- JIANG S., WANG L., REEVES P. Molecular characterization of *S. pneumoniae* types 4, 6B, 8, and 18C capsular polysaccharide gene clusters. *Infect. Immun.*, 2001, **69**, 1244-1255.
- JOHNSON J.R., BERGGREN T., MANIVEL J.C. Histopathologic-microbiologic correlates of invasiveness in a mouse model of ascending unobstructed urinary tract infection. *J. Infect. Dis.*, 1992, **165**, 299-305.
- KARCH, H., SCHMIDT, H, et BRUNDER, W. Plasmid encoded determinants of *Escherichia coli* O157:H7. In : Kaper, J. B. and O'Brien, A. D. (Ed.), *Escherichia coli* O157:H7 and Other Shiga Toxin-producing *E. coli* strains. ASM Press : Washington DC, USA, 1998, 183-194.
- KARMALI M.A. Infection by verocytotoxin-producing *Escherichia coli*. *Clin. Microbiol. Rev.*, 1989, **2**, 15-38.
- KARMALI M.A., PETRIC M., LOUIE S., CHEUNG R. Antigenic heterogeneity of *Escherichia coli* verotoxins. *Lancet*, 1986, **1**, 164-165.
- KAUFFMANN F. The serology of the coli group. *J. Immunol.*, 1947, **57**, 71-100.
- KENNY B., DEVINNEY R., STEIN M., REINSCHIED D.J., FREY E.A., FINLAY B.B. Enteropathogenic *E. coli* (EPEC) transfers its receptor for intimate adherence into mammalian cells. *Cell*, 1997, **91**, 511-520.
- KERR P., BALL H., CHINA B., MAINIL J., FINLAY D., POLLOCK D., WILSON I., MACKIE D. Use of a monoclonal antibody against an *Escherichia coli* O26 surface protein for detection of enteropathogenic and enterohemorrhagic strains. *Clin. Diagn. Lab Immunol.*, 1999, **6**, 610-614.
- KLAPPROTH J.M., SCALETSKY I.C., MCNAMARA B.P., LAI L.C., MALSTROM C., JAMES S.P., DONNENBERG M.S. A large toxin from pathogenic *Escherichia coli* strains that inhibits lymphocyte activation. *Infect. Immun.*, 2000, **68**, 2148-2155.
- KNUTTON S., ROSENSHINE I., PALLAN M.J., NISAN I., NEVES B.C., BAIN C., WOLFF C., DOUGAN G., FRANKEL G. A novel EspA-associated surface organelle of enteropathogenic *Escherichia coli* involved in protein translocation into epithelial cells. *EMBO J.*, 1998, **17**, 2166-2176.
- KONOWALCHUK J., SPEIRS J.I., STAVRIC S. Vero response to a cytotoxin of *Escherichia coli*. *Infect. Immun.*, 1977, **18**, 775-779.
- KRAUSE G., ZIMMERMANN S., BEUTIN L. Investigation of domestic animals and pets as a reservoir for intimin- (*eae*) gene positive *Escherichia coli* types. *Vet. Microbiol.*, 2005, **106**, 87-95.
- KREJANY E.O., GRANT T.H., BENNETT-WOOD V., ADAMS L.M., ROBINS-BROWNE R.M. Contribution of plasmid-encoded fimbriae and intimin to capacity of rabbit-

- specific enteropathogenic *Escherichia coli* to attach to and colonize rabbit intestine. *Infect. Immun.*, 2000, **68**, 6472-6477.
- LALIOUI L., JOUVE M., GOUNON P., LE BOUGUENEC C. Molecular cloning and characterization of the afa-7 and afa-8 gene clusters encoding afimbrial adhesins in *Escherichia coli* strains associated with diarrhea or septicemia in calves. *Infect. Immun.*, 1999, **67**, 5048-5059.
- LEVINE M.M., NATARO J.P., KARCH H., BALDINI M.M., KAPER J.B., BLACK R.E., CLEMENTS M.L., O'BRIEN A.D. The diarrheal response of humans to some classic serotypes of enteropathogenic *Escherichia coli* is dependent on a plasmid encoding an enteroadhesiveness factor. *J. Infect. Dis.*, 1985, **152**, 550-559.
- LALIOUI L., JOUVE M., GOUNON P., LE BOUGUENEC C. Molecular cloning and characterization of the afa-7 and afa-8 gene clusters encoding afimbrial adhesins in *Escherichia coli* strains associated with diarrhea or septicemia in calves. *Infect. Immun.*, 1999, **67**, 5048-5059.
- LAW D. Adhesion and its role in the virulence of enteropathogenic *Escherichia coli*. *Clin. Microbiol. Rev.*, 1994, **7**, 152-173.
- LINGWOOD C.A. Verotoxins and their glycolipid receptors. *Adv. Lipid Res.*, 1993, **25**, 189-211.
- LIU D., REEVES P.R. *Escherichia coli* K12 regains its O antigen. *Microbiology*, 1994, **140** (Pt 1), 49-57.
- LOW A.S., DZIVA F., TORRES A.G., MARTINEZ J.L., ROSSER T., NAYLOR S., SPEARS K., HOLDEN N., MAHAJAN A., FINDLAY J., SALES J., SMITH D.G., LOW J.C., STEVENS M.P., GALLY D.L. Cloning, expression, and characterization of fimbrial operon F9 from enterohemorrhagic *Escherichia coli* O157:H7. *Infect. Immun.*, 2006a, **74**, 2233-2244.
- LOW A.S., HOLDEN N., ROSSER T., ROE A.J., CONSTANTINIDOU C., HOBMAN J.L., SMITH D.G., LOW J.C., GALLY D.L. Analysis of fimbrial gene clusters and their expression in enterohaemorrhagic *Escherichia coli* O157:H7. *Environ. Microbiol.*, 2006b, **8**, 1033-1047.
- LUDWIG, A., GOEBEL, W. Haemolysins of *Escherichia coli*. In : SUSSMAN M.(Ed) (Ed.), *Escherichia coli: mechanisms of virulence*. Cambridge University Press : Cambridge, 1997, 281-329.
- MAINIL J. Shiga/verocytotoxins and Shiga/verotoxigenic *Escherichia coli* in animals. *Vet. Res.*, 1999, **30**, 235-257.
- MAINIL J. Le point des connaissances sur les entérites à *Escherichia coli* chez le veau. *Ann. Méd. Vét.*, 2000, **144**, 121-136.
- MAINIL J. Les souches pathogènes d'*Escherichia coli* chez les chiens et les chats: IV) Discussion générale. *Ann. Méd. Vét.*, 2002, **146**, 219-224.
-

- MAINIL J. Virulence factors and specific properties of invasive *Escherichia coli*: I) Adhesins and colonisation factors. *Ann. Méd. Vét.*, 2003, **147**, 105-126.
- MAINIL J., POHL P. Les souches attachantes et effaçantes d'*Echerischia coli* d'origine bovines. *Ann. Méd. Vét.*, 1994, **138**, 419-429.
- MAINIL J.G., DAUBE G. Verotoxigenic *Escherichia coli* from animals, humans and foods: who's who? *J. Appl. Microbiol.*, 2005, **98**, 1332-1344.
- MAINIL J.G., JACQUEMIN E., HERAULT F., OSWALD E. Presence of pap-, sfa-, and afa-related sequences in necrotoxicogenic *Escherichia coli* isolates from cattle: evidence for new variants of the AFA family. *Can. J. Vet. Res.*, 1997, **61**, 193-199.
- MAINIL J.G., JACQUEMIN E.R., KAECKENBEECK A.E., POHL P.H. Association between the effacing (*eae*) gene and the Shiga-like toxin-encoding genes in *Escherichia coli* isolates from cattle. *Am. J. Vet. Res.*, 1993, **54**, 1064-1068.
- MAINIL J.G., JACQUEMIN E., POHL P., KAECKENBEECK A., BENZ I. DNA sequences coding for the F18 fimbriae and AIDA adhesin are localised on the same plasmid in *Escherichia coli* isolates from piglets. *Vet. Microbiol.*, 2002, **86**, 303-311.
- MCCONNELL M.M., CHART H., SCOTLAND S.M., SMITH H.R., WILLSHAW G.A., ROWE B. Properties of adherence factor plasmids of enteropathogenic *Escherichia coli* and the effect of host strain on expression of adherence to HEp-2 cells. *J. Gen. Microbiol.*, 1989, **135**, 1123-1134.
- MCDANIEL T.K., JARVIS K.G., DONNENBERG M.S., KAPER J.B. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc. Natl. Acad. Sci. U. S A*, 1995, **92**, 1664-1668.
- MELLIES J.L., ELLIOTT S.J., SPERANDIO V., DONNENBERG M.S., KAPER J.B. The Per regulon of enteropathogenic *Escherichia coli* : identification of a regulatory cascade and a novel transcriptional activator, the locus of enterocyte effacement (LEE)-encoded regulator (Ler). *Mol. Microbiol.*, 1999, **33**, 296-306.
- MILON A., OSWALD E., DE RYCKE J. Rabbit EPEC: a model for the study of enteropathogenic *Escherichia coli*. *Vet. Res.*, 1999, **30**, 203-219.
- MONNENS, L., SAVAGE, C. O., et TAYLOR, C. M. Pathophysiology of Haemolytic-uremic syndrome. In : Kaper, J. B. and O'Brien, A. D. (Ed.), *Escherichia coli* O157:H7 and Other Shiga Toxin-producing *E. coli* strains. ASM Press : Washington DC, USA, 1998, 287-292.
- MOON H.W., WHIPP S.C., ARGENZIO R.A., LEVINE M.M., GIANNELLA R.A. Attaching and effacing activities of rabbit and human enteropathogenic *Escherichia coli* in pig and rabbit intestines. *Infect. Immun.*, 1983, **41**, 1340-1351.
- MOXLEY R.A. Edema disease. *Vet. Clin. North Am. Food Anim Pract.*, 2000, **16**, 175-185.
- MOXLEY R.A., FRANCIS D.H. Natural and experimental infection with an attaching and effacing strain of *Escherichia coli* in calves. *Infect. Immun.*, 1986, **53**, 339-346.

-
- NAGY B., FEKETE P.Z. Enterotoxigenic *Escherichia coli* (ETEC) in farm animals. *Vet. Res.*, 1999, **30**, 259-284.
- NAGY B., WHIPP S.C., IMBERECHTS H., BERTSCHINGER H.U., DEAN-NYSTROM E.A., CASEY T.A., SALAJKA E. Biological relationship between F18ab and F18ac fimbriae of enterotoxigenic and verotoxigenic *Escherichia coli* from weaned pigs with oedema disease or diarrhoea. *Microb. Pathog.*, 1997, **22**, 1-11.
- NAKAZATO G., GYLES C., ZIEBELL K., KELLER R., TRABULSI L.R., GOMES T.A., IRINO K., DA SILVEIRA W.D., PESTANA DE CASTRO A.F. Attaching and effacing *Escherichia coli* isolated from dogs in Brazil: characteristics and serotypic relationship to human enteropathogenic *E. coli* (EPEC). *Vet. Microbiol.*, 2004, **101**, 269-277.
- NATARO J.P., KAPER J.B. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.*, 1998, **11**, 142-201.
- NEWTON H.J., SLOAN J., BENNETT-WOOD V., ADAMS L.M., ROBINS-BROWNE R.M., HARTLAND E.L. Contribution of long polar fimbriae to the virulence of rabbit-specific enteropathogenic *Escherichia coli*. *Infect. Immun.*, 2004, **72**, 1230-1239.
- NICHOLLS L., GRANT T.H., ROBINS-BROWNE R.M. Identification of a novel genetic locus that is required for in vitro adhesion of a clinical isolate of enterohaemorrhagic *Escherichia coli* to epithelial cells. *Mol. Microbiol.*, 2000, **35**, 275-288.
- NIEWERTH U., FREY A., VOSS T., LE BOUGUENEC C., BALJER G., FRANKE S., SCHMIDT M.A. The AIDA autotransporter system is associated with F18 and stx2e in *Escherichia coli* isolates from pigs diagnosed with edema disease and postweaning diarrhea. *Clin. Diagn. Lab Immunol.*, 2001, **8**, 143-149.
- NGELEKA M., PRITCHARD J., APPELYARD G., MIDDLETON D.M., FAIRBROTHER J.M. Isolation and association of *Escherichia coli* AIDA-I/STb, rather than EAST1 pathotype, with diarrhea in piglets and antibiotic sensitivity of isolates. *J. Vet. Diagn. Invest.*, 2003, **15**, 242-252.
- O'BRIEN A.D., LAVECK G.D., THOMPSON M.R., FORMAL S.B. Production of *Shigella dysenteriae* type 1-like cytotoxin by *Escherichia coli*. *J. Infect. Dis.*, 1982, **146**, 763-769.
- ORSKOV, F. Facultatively anaerobic Gram-negative rods. Family I. *Enterobacteriaceae*. Genus II. *Escherichia coli*. In : Krieg, N. R. and Holt, J. G. (Ed.), *Bergey's Manual of Systematic Bacteriology*, Volume 1. Williams & Wilkins : Baltimore/London, 1984, 420-423.
- ORSKOV F., ORSKOV I. Serotyping of *Escherichia coli*. *Methods Microbiol.*, 1984, **14**, 43-112.
- OSTROFF S.M., TARR P.I., NEILL M.A., LEWIS J.H., HARGRETT-BEAN N., KOBAYASHI J.M. Toxin genotypes and plasmid profiles as determinants of systemic sequelae in *Escherichia coli* O157:H7 infections. *J. Infect. Dis.*, 1989, **160**, 994-998.
-

- OSWALD E., SCHMIDT H., MORABITO S., KARCH H., MARCHES O., CAPRIOLI A. Typing of intimin genes in human and animal enterohemorrhagic and enteropathogenic *Escherichia coli*: characterization of a new intimin variant. *Infect. Immun.*, 2000, **68**, 64-71.
- PALLEN M.J., BEATSON S.A., BAILEY C.M. Bioinformatics analysis of the locus for enterocyte effacement provides novel insights into type-III secretion. *BMC. Microbiol.*, 2005, **5**, 9-29.
- PATON A.W., SRIMANOTE P., WOODROW M.C., PATON J.C. Characterization of *Saa*, a novel autoagglutinating adhesin produced by locus of enterocyte effacement-negative Shiga-toxigenic *Escherichia coli* strains that are virulent for humans. *Infect. Immun.*, 2001, **69**, 6999-7009.
- PATON J.C., PATON A.W. Pathogenesis and diagnosis of Shiga toxin-producing *Escherichia coli* infections. *Clin. Microbiol. Rev.*, 1998, **11**, 450-479.
- PEETERS J. Enteropathogenic strains of *Escherichia coli* in rabbits. *Annales de Medecine Veterinaire*, 1993, **137**, 361-368.
- PERNA N.T., PLUNKETT G., III, BURLAND V., MAU B., GLASNER J.D., ROSE D.J., MAYHEW G.F., EVANS P.S., GREGOR J., KIRKPATRICK H.A., POSFAI G., HACKETT J., KLINK S., BOUTIN A., SHAO Y., MILLER L., GROTEBECK E.J., DAVIS N.W., LIM A., DIMALANTA E.T., POTAMOUCIS K.D., APODACA J., ANANTHARAMAN T.S., LIN J., YEN G., SCHWARTZ D.C., WELCH R.A., BLATTNER F.R. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature*, 2001, **409**, 529-533.
- POHL P. Les souches pathogènes d'*Escherichia coli*, *histoire et classification* . *Ann. Méd. Vét.*, 1993, **137**, 325-333.
- POLOTSKY Y.E., DRAGUNSKAYA E.M., SELIVERSTOVA V.G., AVDEEVA T.A., CHAKHUTINSKAYA M.G., KÉTYI I., VERTÉNYI A., RALOVICH B., EMODY L., MÁLOVICS I., SAFONOVA N.V., SNIGIREVSKAYA E.S., KARYAGINA E.I. Pathogenic effect of enterotoxigenic *Escherichia coli* and *Escherichia coli* causing infantile diarrhoea. *Acta Microbiol. Acad. Sci. Hung.*, 1977, **24**, 221-236.
- PRITCHARD J., NGELEKA M., MIDDLETON D.M. In vivo and in vitro colonization patterns of AIDA-I-positive *Escherichia coli* isolates from piglets with diarrhea. *J. Vet. Diagn. Invest.*, 2004, **16**, 108-115.
- PRUIMBOOM-BREES I.M., MORGAN T.W., ACKERMANN M.R., NYSTROM E.D., SAMUEL J.E., CORNICK N.A., MOON H.W. Cattle lack vascular receptors for *Escherichia coli* O157:H7 Shiga toxins. *Proc. Natl. Acad. Sci. U. S A*, 2000, **97**, 10325-10329.
- QADRI F., SVENNERHOLM A.M., FARUQUE A.S., SACK R.B. Enterotoxigenic *Escherichia coli* in developing countries: epidemiology, microbiology, clinical features, treatment, and prevention. *Clin. Microbiol. Rev.*, 2005, **18**, 465-483.
- RIPPINGER P., BERTSCHINGER H.U., IMBERECHTS H., NAGY B., SORG I., STAMM M., WILD P., WITTIG W. Designations F18ab and F18ac for the related fimbrial

- types F107, 2134P and 8813 of *Escherichia coli* isolated from porcine postweaning diarrhoea and from oedema disease. *Vet. Microbiol.*, 1995, **45**, 281-295.
- ROSE, P., CHANT, I. Haematology of Hemolytic-Uremic syndrome. In : Kaper, J. B. and O'Brien, A. D. (Ed.), *Escherichia coli* O157:H7 and Other Shiga Toxin-producing *E. coli* strains. ASM Press : Washington DC, USA, 1998, 293-303.
- ROSENSHINE I., RUSCHKOWSKI S., FINLAY B.B. Expression of attaching/effacing activity by enteropathogenic *Escherichia coli* depends on growth phase, temperature, and protein synthesis upon contact with epithelial cells. *Infect. Immun.*, 1996, **64**, 966-973.
- ROTHBAUM R., MCADAMS A.J., GIANNELLA R., PARTIN J.C. A clinicopathologic study of enterocyte-adherent *Escherichia coli*: a cause of protracted diarrhea in infants. *Gastroenterology*, 1982, **83**, 441-454.
- ROWE, B., GROSS, R. J. Facultatively anaerobic Gram-negative rods. Family I. *Enterobacteriaceae*. Genus II. *Shigella*. In : Krieg, N. R. and Holt, J. G. (Ed.), *Bergey's Manual of Systematic Bacteriology*, Volume 1. Williams & Wilkins : Baltimore/London, 1984, 423-427.
- SANCAK A.A., RUTGERS H.C., HART C.A., BATT R.M. Prevalence of enteropathic *Escherichia coli* in dogs with acute and chronic diarrhoea. *Vet. Rec.*, 2004, **154**, 101-106.
- SANDVIG K. Shiga toxins. *Toxicon*, 2001, **39**, 1629-1635.
- SCALETSKY I.C., SILVA M.L., TRABULSI L.R. Distinctive patterns of adherence of enteropathogenic *Escherichia coli* to HeLa cells. *Infect. Immun.*, 1984, **45**, 534-536.
- SCHEUTZ, F., BEUTIN, L., PIERARD, D., et SMITH, H. R. Nomenclature of Verocytotoxins. In : Duffy, G., Garvey, P., and Mc Dowell, D. A. Eds (Ed.), *Verocytotoxigenic Escherichia coli*. Food & Nutrition Press Inc. Trumbull, 2001, 447-452.
- SCHMIDT, H, BITZAN, M., et KARCH, H. Pathogenic aspects of Shiga toxin-producing *E. coli* infections in humans. In : Duffy, G., Garvey, P., and Mc Dowell, D. A. Eds (Ed.), *Verocytotoxigenic Escherichia coli*. Food & Nutrition Press Inc. Trumbull, 2001, 241-262.
- SCHMIDT H., BEUTIN L., KARCH H. Molecular analysis of the plasmid-encoded hemolysin of *Escherichia coli* O157:H7 strain EDL 933. *Infect. Immun.*, 1995, **63**, 1055-1061.
- SCHMIDT H., KARCH H. Enterohemolytic phenotypes and genotypes of shiga toxin-producing *Escherichia coli* O111 strains from patients with diarrhea and hemolytic-uremic syndrome. *J. Clin. Microbiol.*, 1996, **34**, 2364-2367.
- SCHMIDT H., KERNBACH C., KARCH H. Analysis of the EHEC hly operon and its location in the physical map of the large plasmid of enterohaemorrhagic *Escherichia coli* O157:H7. *Microbiology*, 1996, **142 (Pt 4)**, 907-914.
-

-
- SCOTLAND S.M., SMITH H.R., ROWE B. Two distinct toxins active on Vero cells from *Escherichia coli* O157. *Lancet*, 1985, **2**, 885-886.
- SHERLOCK O., SCHEMBRI M.A., REISNER A., KLEMM P. Novel roles for the AIDA adhesin from diarrheagenic *Escherichia coli*: cell aggregation and biofilm formation. *J. Bacteriol.*, 2004, **186**, 8058-8065.
- SMITH H.W., GREEN P., PARSELL Z. Vero cell toxins in *Escherichia coli* and related bacteria: transfer by phage and conjugation and toxic action in laboratory animals, chickens and pigs. *J. Gen. Microbiol.*, 1983, **129**, 3121-3137.
- SOJKA, W.J. *Escherichia coli* in domestic animals and poultry. Commonwealth Agricultural Bureaux: Farnham Royal : 1965.
- SONNTAG A.K., BIELASZEWSKA M., MELLMANN A., DIERKSEN N., SCHIERACK P., WIELER L.H., SCHMIDT M.A., KARCH H. Shiga toxin 2e-producing *Escherichia coli* isolates from humans and pigs differ in their virulence profiles and interactions with intestinal epithelial cells. *Appl. Environ. Microbiol.*, 2005, **71**, 8855-8863.
- STALEY T.E., JONES K., CORLEY L.D. Attachment and penetration of *Escherichia coli* into intestinal epithelium of the ileum in newborn pigs. *Am. J. Pathol.*, 1969, **56**, 371-392.
- STEVENS M.P., VAN DIEMEN P.M., FRANKEL G., PHILLIPS A.D., WALLIS T.S. Efa1 influences colonization of the bovine intestine by shiga toxin-producing *Escherichia coli* serotypes O5 and O111. *Infect. Immun.*, 2002, **70**, 5158-5166.
- STEVENSON G., ANDRIANOPOULOS K., HOBBS M., REEVES P.R. Organization of the *Escherichia coli* K-12 gene cluster responsible for production of the extracellular polysaccharide colanic acid. *J. Bacteriol.*, 1996, **178**, 4885-4893.
- STORDEUR P., CHINA B., CHARLIER G., ROELS S., MAINIL J. Clinical signs, reproduction of attaching/effacing lesions, and enterocyte invasion after oral inoculation of an O118 enterohaemorrhagic *Escherichia coli* in neonatal calves. *Microbes Infect.*, 2000, **2**, 17-24.
- STROCKBINE N.A., MARQUES L.R., NEWLAND J.W., SMITH H.W., HOLMES R.K., O'BRIEN A.D. Two toxin-converting phages from *Escherichia coli* O157:H7 strain 933 encode antigenically distinct toxins with similar biologic activities. *Infect. Immun.*, 1986, **53**, 135-140.
- TAKAO T., TANABE T., HONG Y.M., SHIMONISHI Y., KURAZONO H., YUTSUDO T., SASAKAWA C., YOSHIKAWA M., TAKEDA Y. Identity of molecular structure of Shiga-like toxin I (VT1) from *Escherichia coli* O157:H7 with that of Shiga toxin. *Microb. Pathog.*, 1988, **5**, 57-69.
- TARR P.I., BILGE S.S., VARY J.C., JR., JELACIC S., HABEEB R.L., WARD T.R., BAYLOR M.R., BESSER T.E. Iha: a novel *Escherichia coli* O157:H7 adherence-conferring molecule encoded on a recently acquired chromosomal island of conserved structure. *Infect. Immun.*, 2000, **68**, 1400-1407.
-

-
- TATSUNO I., HORIE M., ABE H., MIKI T., MAKINO K., SHINAGAWA H., TAGUCHI H., KAMIYA S., HAYASHI T., SASAKAWA C. *tox*B gene on pO157 of enterohemorrhagic *Escherichia coli* O157:H7 is required for full epithelial cell adherence phenotype. *Infect. Immun.*, 2001, **69**, 6660-6669.
- TATSUNO I., MUNDY R., FRANKEL G., CHONG Y., PHILLIPS A.D., TORRES A.G., KAPER J.B. The *lpf* gene cluster for long polar fimbriae is not involved in adherence of enteropathogenic *Escherichia coli* or virulence of *Citrobacter rodentium*. *Infect. Immun.*, 2006, **74**, 265-272.
- TOBE T., SASAKAWA C. Role of bundle-forming pilus of enteropathogenic *Escherichia coli* in host cell adherence and in microcolony development. *Cell Microbiol.*, 2001, **3**, 579-585.
- TOBE T., SASAKAWA C. Species-specific cell adhesion of enteropathogenic *Escherichia coli* is mediated by type IV bundle-forming pili. *Cell Microbiol.*, 2002, **4**, 29-42.
- TOBE T., SCHOOLNIK G.K., SOHEL I., BUSTAMANTE V.H., PUENTE J.L. Cloning and characterization of *bfp*TVW, genes required for the transcriptional activation of *bfp*A in enteropathogenic *Escherichia coli*. *Mol. Microbiol.*, 1996, **21**, 963-975.
- TOMA C., HIGA N., IYODA S., RIVAS M., IWANAGA M. The long polar fimbriae genes identified in Shiga toxin-producing *Escherichia coli* are present in other diarrheagenic *E. coli* and in the standard *E. coli* collection of reference (ECOR) strains. *Res. Microbiol.*, 2005, **157**, 153-161.
- TOMA C., MARTINEZ E.E., SONG T., MILIWEBSKY E., CHINEN I., IYODA S., IWANAGA M., RIVAS M. Distribution of putative adhesins in different seropathotypes of Shiga toxin-producing *Escherichia coli*. *J. Clin. Microbiol.*, 2004, **42**, 4937-4946.
- TORRES A.G., GIRON J.A., PERNA N.T., BURLAND V., BLATTNER F.R., AVELINO-FLORES F., KAPER J.B. Identification and characterization of *lpf*ABCC'DE, a fimbrial operon of enterohemorrhagic *Escherichia coli* O157:H7. *Infect. Immun.*, 2002, **70**, 5416-5427.
- TORRES A.G., ZHOU X., KAPER J.B. Adherence of diarrheagenic *Escherichia coli* strains to epithelial cells. *Infect. Immun.*, 2005, **73**, 18-29.
- TOTH I., SCHMIDT H., DOW M., MALIK A., OSWALD E., NAGY B. Transduction of porcine enteropathogenic *Escherichia coli* with a derivative of a shiga toxin 2-encoding bacteriophage in a porcine ligated ileal loop system. *Appl. Environ. Microbiol.*, 2003, **69**, 7242-7247.
- TOZZI A.E., CAPRIOLI A., MINELLI F., GIANVITI A., DE PETRIS L., EDEFONTI A., MONTINI G., FERRETTI A., DE PALO T., GAIDO M., RIZZONI G. Shiga toxin-producing *Escherichia coli* infections associated with hemolytic uremic syndrome, Italy, 1988-2000. *Emerg. Infect. Dis.*, 2003, **9**, 106-108.
- TOZZOLI R., CAPRIOLI A., MORABITO S. Detection of *tox*B, a plasmid virulence gene of *Escherichia coli* O157, in enterohemorrhagic and enteropathogenic *E. coli*. *J. Clin. Microbiol.*, 2005, **43**, 4052-4056.
-

- TRABULSI L.R., KELLER R., TARDELLI GOMES T.A. Typical and atypical enteropathogenic *Escherichia coli*. *Emerg. Infect. Dis.*, 2002, **8**, 508-513.
- TREVENA W.B., HOOPER R.S., WRAY C., WILLSHAW G.A., CHEASTY T., DOMINGUE G. Vero cytotoxin-producing *Escherichia coli* O157 associated with companion animals. *Vet. Rec.*, 1996, **138**, 400-400.
- ULSHEN M.H., ROLLO J.L. Pathogenesis of *Escherichia coli* gastroenteritis in man--another mechanism. *N. Engl. J. Med.*, 1980, **302**, 99-101.
- WELCH, R. A. Phylogenetic analyses of RTX family. In : Roth, J. A., Bolin, C. A., Brogden, K. A., Minion, F. C., and Wannemuehler, M. J. (Ed.), Virulence mechanisms of bacterial pathogens. ASM Press : Washington, 1995, 195-206.
- WIELER L.H., BAUERFEIND R. STEC as a veterinary problem. Diagnostics and prophylaxis in animals. *Methods Mol. Med.*, 2003, **73**, 75-89.
- WITTIG W., PRAGER R., STAMM M., STRECKEL W., TSCHAPE H. Expression and plasmid transfer of genes coding for the fimbrial antigen F107 in porcine *Escherichia coli* strains. *Zentralbl. Bakteriol.*, 1994, **281**, 130-139.
- WOLF M.K., ANDREWS G.P., FRITZ D.L., SJOGREN R.W., JR., BOEDEKER E.C. Characterization of the plasmid from *Escherichia coli* RDEC-1 that mediates expression of adhesin AF/R1 and evidence that AF/R1 pili promote but are not essential for enteropathogenic disease. *Infect. Immun.*, 1988, **56**, 1846-1857.
- WOLF M.K., BOEDEKER E.C. Cloning of the genes for AF/R1 pili from rabbit enteroadherent *Escherichia coli* RDEC-1 and DNA sequence of the major structural subunit. *Infect. Immun.*, 1990, **58**, 1124-1128.
- WRAY C., MCLAREN I., PEARSON G.R. Occurrence of 'attaching and effacing' lesions in the small intestine of calves experimentally infected with bovine isolates of verocytotoxic *E. coli*. *Vet. Rec.*, 1989, **125**, 365-368.
- ZHU C., HAREL J., JACQUES M., DESAUTELS C., DONNENBERG M.S., BEAUDRY M., FAIRBROTHER J.M. Virulence properties and attaching-effacing activity of *Escherichia coli* O45 from swine postweaning diarrhea. *Infect. Immun.*, 1994, **62**, 4153-4159.
- ZUNIGA A., YOKOYAMA H., ALBICKER-RIPPINGER P., EGGENBERGER E., BERTSCHINGER H.U. Reduced intestinal colonisation with F18-positive enterotoxigenic *Escherichia coli* in weaned pigs fed chicken egg antibody against the fimbriae. *FEMS Immunol. Med. Microbiol.*, 1997, **18**, 153-161.