
Bibliographie

A

- Abu-Ghazaleh R.I., Gleich G.J., Prendergasft G. Interaction of eosinophil granule major basic protein with synthetic lipid bilayers : A mechanism for toxicity. *J. Membr. Biol.*, 1992, **128**, 153-167.
- Adachi T., Choudhury B.K., Stafford S., Sur S., Alam R. The differential role of extracellular signal-regulated kinases and p38 mitogen-activated protein kinase in eosinophil functions. *J. Immunol.*, 2000, **165**, 2198-2204.
- Adam-Klages S., Adam D., Wiegmann K., Struve S., Kolanus W., Schneider-Mergener J., Kronke M. FAN, a novel WD-repeat protein, couples the p55 TNF-receptor to neutral sphingomyelinase. *Cell*, 1996, **86**, 937-947.
- Adam-Klages S., Schwandner R., Luschen S., Ussat S., Kreder D., Kronke M. Caspase-mediated inhibition of human cytosolic phospholipase A2 during apoptosis. *J. Immunol.*, 1998, **161**, 5687-5694.
- Akyurek N., Ren Y., Rassidakis G.Z., Schlette E.J., Medeiros L.J. Expression of inhibitor of apoptosis proteins in B-cell non-Hodgkin and Hodgkin lymphomas. *Cancer*, 2006, **107**, 1844-1851.
- Alessenko A., Chatterjee S. Neutral sphingomyelinase: localization in rat liver nuclei and involvement in regeneration/proliferation. *Mol. Cell. Biochem.*, 1995, **143**, 169-174.
- Allan L.A., Morrice N., Brady S., Magee G., Pathak S., Clarke P.R. Inhibition of caspase-9 through phosphorylation at Thr 125 by ERK MAPK. *Nat. Cell Biol.*, 2003, **5**, 647-654.
- Allen R.T., Hunter W.J. 3rd, Agrawal D.K. Morphological and biochemical characterization and analysis of apoptosis. *J. Pharmacol. Toxicol. Methods*, 1997, **37**, 215-228.
- Alnemri E.S., Livingston D.J., Nicholson D.W., Salvesen G., Thornberry N.A., Wong W.W., Yuan J. Human ICE/CED-3 protease nomenclature. *Cell*, 1996, **87**, 171.
- Alon R., Fuhlbrigge R.C., Finger E.B., Springer T.A. Interactions through L-selectin between leukocytes and adherent leukocytes nucleate rolling adhesions on selectins and VCAM-1 in shear flow. *J. Cell Biol.*, 1996, **135**, 849-865.
- Al-Shami A., Mahanna W., Naccache P.H. Granulocyte-macrophage colony-stimulating factor-activated signaling pathways in human neutrophils. Selective activation of Jak2, Stat3, and Stat5b. *J. Biol. Chem.*, 1998, **273**, 1058-1063.
- Altzner F., Conus S., Cavalli A., Folkers G., Simon H.-U. Calpain-1 regulates Bax and subsequent Smac-dependent caspase-3 activation in neutrophil apoptosis. *J. Biol. Chem.*, 2004, **279**, 5947-5957.
- Alvarado-Kristensson M., Melander F., Leandersson K., Ronnstrand L., Wernstedt C., Andersson T. p38-MAPK signals survival by phosphorylation of caspase-8 and caspase-3 in human neutrophils. *J. Exp. Med.*, 2004, **199**, 449-458.
- Alvarado-Kristensson M., Andersson T. Protein phosphatase 2A regulates apoptosis in neutrophils by dephosphorylating both p38 MAPK and its substrate caspase 3. *J. Biol Chem.*, 2005, **280**, 6238-6244.
- Amarante-Mendes G.P., Green D.R. The regulation of apoptotic cell death. *Braz. J. Med. Biol. Res.*, 1999, **32**, 1053-1061.
- Ambrosini G., Adida C., Altieri D.C. A novel anti-apoptosis gene, survivin, expressed in cancer and lymphoma. *Nat. Med.*, 1997, **3**, 917-921.
- Ameisen J.C., Estaquier J., Idziorek T., De Bels F. The relevance of apoptosis to AIDS pathogenesis. *Trends Cell. Biol.*, 1995, **5**, 27-32.
- Anderson P. Kinase cascades regulating entry into apoptosis. *Microbiol. Mol. Biol. Rev.*, 1997, **61**, 33-46.

- Anto J.M., Vermeire P., Sunyer J. Chronic obstructive pulmonary disease. In: European Respiratory Monograph, Volume 5, Monograph 15 : Respiratory Epidemiology in Europe, Annesi-Maesano I, Gulsvik A, Viegi G (Eds), Sheffield, UK : European Respiratory Society Journals Ltd, 2000, 1-22.
- Antonsson B., Montessuit S., Lauper S., Eskes R., Martinou J.C. Bax oligomerization is required for channel-forming activity in liposomes and to trigger cytochrome c release from mitochondria. *Biochem. J.*, 2000, **345**, 271-278.
- Antonsson B., Montessuit S., Sanchez B., Martinou J.C. Bax is present as a high molecular weight oligomer/complex in the mitochondrial membrane of apoptotic cells. *J. Biol. Chem.*, 2001, **276**, 11615-11623.
- Athens J.W., Haab O.P., Raab S.O., Mauer A.M., Ashenbrucker H., Cartwright G.E., Wintrobe M.M. Leukokinetic studies. IV. The total blood, circulating and marginal granulocyte pools and the granulocyte turnover rate in normal subjects. *J. Clin. Invest.*, 1961, **40**, 989-995.
- Auge N., Escargueil-Blanc I., Lajoie-Mazenc I., Suc I., Andrieu-Abadie N., Pieraggi M.T., Chatelut M., Thiers J.C., Jaffrezou J.P., Laurent G., Levade T., Negre-Salvayre A., Salvayre R. Potential role for ceramide in mitogen-activated protein kinase activation and proliferation of vascular smooth muscle cells induced by oxidized low density lipoprotein. *J. Biol. Chem.*, 1998, **273**, 12893-12900.

B

- Babior B.M. The respiratory burst of phagocytes. *J. Clin. Invest.*, 1984, **73**, 599-601.
- Baggiolini M., Dewald B., Moser B. Interleukin-8 and related chemotactic cytokines--CXC and CC chemokines. *Adv. Immunol.*, 1994, **55**, 97-179.
- Banchereau J., Bazan F., Blanchard D., Briere F., Galizzi J.P., van Kooten C., Liu Y.J., Rousset F., Saeland S. The CD40 antigen and its ligand. *Annu. Rev. Immunol.*, 1994, **12**, 881-922.
- Bargatze R.F., Kurk S., Butcher E.C., Jutila M.A. Neutrophils roll on adherent neutrophils bound to cytokine-induced endothelial cells via L-selectin on the rolling cells. *J. Exp. Med.*, 1994, **180**, 1785-1792.
- Barker R.L., Gleich G.J., Pease L.R. Acidic precursor revealed in human eosinophil granule major basic protein cDNA. *J. Exp. Med.*, 1988, **168**, 1493-1498.
- Barkett M., Xue D., Horvitz H.R., Gilmore T.D. Phosphorylation of IkappaB-alpha inhibits its cleavage by caspase CPP32 *in vitro*. *J. Biol. Chem.*, 1997, **272**, 29419-29422.
- Barnes P.J. Cytokine-directed therapies for the treatment of chronic airway diseases. *Cytokine Growth Factor. Rev.*, 2003, **14**, 511-522.
- Bates M.E., Busse W.W., Bertics P.J. Interleukin 5 signals through Shc and Grb2 in human eosinophils. *Am. J. Respir. Cell. Mol. Biol.*, 1998, **18**, 75-83.
- Bektas M., Spiegel S. Glycosphingolipids and cell death. *Glycoconj. J.*, 2004, **20**, 39-47.
- Belaouaj A., McCarthy R., Baumann M., Gao Z., Ley T.J., Abraham S.N., Shapiro S.D. Mice lacking neutrophil elastase reveal impaired host defense against gram negative bacterial sepsis. *Nat. Med.*, 1998, **4**, 615-618.
- Belvisi M.G., Bottomley K.M. The role of matrix metalloproteinases (MMPs) in the pathophysiology of chronic obstructive pulmonary disease (COPD): a therapeutic role for inhibitors of MMPs ? *Inflamm. Res.*, 2003, **52**, 95-100.
- Belvisi M.G. Regulation of inflammatory cell function by corticosteroids. *Proc. Am. Thorac. Soc.*, 2004, **1**, 207-214.
- Benelli R., Albin A., Noonan D. Neutrophils and angiogenesis: potential initiators of the angiogenic cascade. *Chem. Immunol. Allergy*, 2003, **83**, 167-181.

- Bertolotto C., Ricci J.E., Luciano F., Mari B., Chambard J.C., Auberger P. Cleavage of the serum response factor during death receptor-induced apoptosis results in an inhibition of the c-FOS promoter transcriptional activity. *J. Biol. Chem.*, 2000, **275**, 12941-12947.
- Bezombes C., Segui B., Cuvillier O., Bruno A.P., Uro-Coste E., Gouaze V., Andrieu-Abadie N., Carpentier S., Laurent G., Salvayre R., Jaffrezou J.P., Levade T. Lysosomal sphingomyelinase is not solicited for apoptosis signaling. *FASEB J.*, 2001, **15**, 297-299.
- Bhunia A.K., Han H., Snowden A., Chatterjee S. Lactosylceramide stimulates Ras-GTP loading, kinases (MEK, Raf), p44 mitogen-activated protein kinase, and c-fos expression in human aortic smooth muscle cells. *J. Biol. Chem.*, 1996, **271**, 10660-10666.
- Birbes H., El Bawab S., Hannun Y.A., Obeid L.M. Selective hydrolysis of a mitochondrial pool of sphingomyelin induces apoptosis. *FASEB J.*, 2001, **15**, 2669-2679.
- Black R.A., Kronheim S.R., Merriam J.E., March C.J., Hopp T.P. A pre-aspartate-specific protease from human leukocytes that cleaves pro-interleukin-1 beta. *J. Biol. Chem.*, 1989, **264**, 5323-5326.
- Bleicher R.J., Cabot M.C. Glucosylceramide synthase and apoptosis. *Biochim. Biophys. Acta*, 2002, **1585**, 172-178.
- Blume-Jensen P., Janknecht R., Hunter T. The kit receptor promotes cell survival via activation of PI 3-kinase and subsequent Akt-mediated phosphorylation of Bad on Ser136. *Curr. Biol.*, 1998, **8**, 779-782.
- Bohnsack J.F., Chang J., Zhou X., Yednock T.A. Mechanisms of beta 1 integrin-dependent adherence of granulocytic HL60 to fibronectin. *J. Leukoc. Biol.*, 1995, **57**, 592-599.
- Boland M.P., Foster S.J., O'Neill L.A. Daunorubicin activates NFkappaB and induces kappaB-dependent gene expression in HL-60 promyelocytic and Jurkat T lymphoma cells. *J. Biol. Chem.*, 1997, **272**, 12952-12960.
- Bories D., Raynal M.C., Solomon D.H., Darzynkiewicz Z., Cayre Y.E. Down-regulation of a serine protease, myeloblastin, causes growth arrest and differentiation of promyelocytic leukemia cells. *Cell*, 1989, **59**, 959-968.
- Borner C., Monney L. Apoptosis without caspases: an inefficient molecular guillotine ? *Cell Death Differ.*, 1999, **6**, 497-507.
- Borregaard N., Cowland J.B. Granules of the human neutrophilic polymorphonuclear leukocyte. *Blood*, 1997, **89**, 3503-3521.
- Bose R., Verheij M., Haimovitz-Friedman A., Scotto K., Fuks Z., Kolesnick R. Ceramide synthase mediates daunorubicin-induced apoptosis: an alternative mechanism for generating death signals. *Cell*, 1995, **82**, 405-414.
- Bouchier-Hayes L., Lartigue L., Newmeyer D.D. Mitochondria: pharmacological manipulation of cell death. *J. Clin. Invest.*, 2005, **115**, 2640-2647.
- Bourteele S., Hausser A., Doppler H., Horn-Muller J., Ropke C., Schwarzmann G., Pfizenmaier K., Muller G. Tumor necrosis factor induces ceramide oscillations and negatively controls sphingolipid synthases by caspases in apoptotic Kym-1 cells. *J. Biol. Chem.*, 1998, **273**, 31245-31251.
- Bousquet J., Chanez P., Lacoste J.Y., Barneon G., Ghavanian N., Enander I., Venge P., Ahlstedt S., Simony-Lafontaine J., Godard P., Michel F.B. Eosinophilic inflammation in asthma. *N. Engl. J. Med.*, 1990, **323**, 1033-1039.
- Brach M.A., deVos S., Gruss H.J., Herrmann F. Prolongation of survival of human polymorphonuclear neutrophils by granulocyte-macrophage colony-stimulating factor is caused by inhibition of programmed cell death. *Blood*, 1992, **80**, 2920-2924.
- Brady S.C., Lindsey A.A., Clarke P.R. Regulation of caspase 9 through phosphorylation by protein kinase C zeta in response to hyperosmotic stress. *Mol. Cell. Biol.*, 2005, **25**, 10543-10555.

- Brenner B., Ferlinz K., Grassme H., Weller M., Koppenhoefer U., Dichgans J., Sandhoff K., Lang F., Gulbins E. Fas/CD95/Apo-I activates the acidic sphingomyelinase via caspases. *Cell. Death Differ.*, 1998, **5**, 29-37.
- Brizzi M.F., Aronica M.G., Rosso A., Bagnara G.P., Yarden Y., Pegoraro L. Granulocyte-macrophage colony-stimulating factor stimulates JAK2 signaling pathway and rapidly activates p93fes, STAT1 p91, and STAT3 p92 in polymorphonuclear leukocytes. *J. Biol. Chem.*, 1996, **271**, 3562-3567.
- Broide D.H., Paine M.M., Firestein G.S. Eosinophils express interleukin 5 and granulocyte macrophage-colony-stimulating factor mRNA at sites of allergic inflammation in asthmatics. *J. Clin. Invest.*, 1992, **90**, 1414–1424.
- Brough D., Rothwell N.J. Caspase-1-dependent processing of pro-interleukin-1beta is cytosolic and precedes cell death. *J. Cell Sci.*, 2007, **120**, 772-781.
- Brown D.A., London E. Structure and function of sphingolipid- and cholesterol-rich membrane rafts. *J. Biol. Chem.*, 2000, **275**, 17221-17224.
- Bureau F., Vanderplasschen A., Jaspar F., Minner F., Pastoret P.-P., Merville M.P., Bours V., Lekeux P. Constitutive nuclear factor-kappaB activity preserves homeostasis of quiescent mature lymphocytes and granulocytes by controlling the expression of distinct Bcl-2 family proteins. *Blood*, 2002, **99**, 3683-3691.
- Busse W.W., Sedgwick J.B. Eosinophils in asthma. *Ann. Allergy*, 1991, **68**, 286-290.
- Busse W.W., Lemanske R.F. Asthma. *N. Engl. J. Med.*, 2001, **344**, 350-362.
- Byun Y., Chen F., Chang R., Trivedi M., Green K.J., Cryns V.L. Caspase cleavage of vimentin disrupts intermediate filaments and promotes apoptosis. *Cell. Death Differ.*, 2001, **8**, 443-450.

C

- Candeias L.P., Patel K.B., Stratford M.R., Wardman P. Free hydroxyl radicals are formed on reaction between the neutrophil-derived species superoxide anion and hypochlorous acid. *FEBS Lett.*, 1993, **333**, 151-153.
- Capron M., Tomassini M., Ven der Vorst E., Kusnierz J.P., Pepin K.G., Capron A. Existence et fonctions d'un recepteur pour l'immunoglobuline A sur les eosinophiles humaines. *CRC Acad. Sci. Immunol.*, 1988, **307**, 397– 405.
- Cardone M.H., Roy N., Stennicke H.R., Salvesen G.S., Franke T.F., Stanbridge E., Frisch S., Reed J.C. Regulation of cell death protease caspase-9 by phosphorylation. *Science*, 1998, **282**, 1318-1321.
- Carreras M.C., Pargament G.A., Catz S.D., Poderoso J.J., Boveris A. Kinetics of nitric oxide and hydrogen peroxide production and formation of peroxynitrite during the respiratory burst of human neutrophils. *FEBS Lett.*, 1994, **341**, 65-68.
- Casciola-Rosen L., Nicholson D.W., Chong T., Rowan K.R., Thornberry N.A., Miller D.K., Rosen A. Apopain/CPP32 cleaves proteins that are essential for cellular repair: a fundamental principle of apoptotic death. *J. Exp. Med.*, 1996, **183**, 1957-1964.
- Cassatella M.A. Neutrophil-derived proteins: selling cytokines by the pound. *Adv. Immunol.*, 1999, **73**, 369-509.
- Castillo S.S., Levy M., Wang C., Thaikootathil J.V., Khan E., Goldkorn T. Nitric oxide-enhanced caspase-3 and acidic sphingomyelinase interaction: a novel mechanism by which airway epithelial cells escape ceramide-induced apoptosis. *Exp. Cell. Res.*, 2007, **313**, 816-823.
- Cataldo D.D., Gueders M.M., Rocks N., Sounni N.E., Evrard B., Bartsch P., Louis R., Noel A., Foidart J.M.. Pathogenic role of matrix metalloproteases and their inhibitors in asthma and chronic obstructive pulmonary disease and therapeutic relevance of matrix metalloproteases inhibitors. *Cell. Mol. Biol. (Noisy-le-Grand)*, 2003, **49**, 875-884.

- Caulin C., Salvesen G.S., Oshima R.G. Caspase cleavage of keratin 18 and reorganization of intermediate filaments during epithelial cell apoptosis. *J. Cell. Biol.*, 1997, **138**, 1379-1394.
- Caux C., Burdin N., Galibert L., Hermann P., Renard N., Servet-Delprat C., Banchereau J. Functional CD40 on B lymphocytes and dendritic cells. *Res. Immunol.*, 1994, **145**, 235-249.
- Chai J., Du C., Wu J.W., Kyin S., Wang X., Shi Y. Structural and biochemical basis of apoptotic activation by Smac/Diablo. *Nature*, 2000, **406**, 855-862.
- Chai J., Shiozaki E., Srinivasula S.M., Wu Q., Datta P., Alnemri E.S., Shi Y. Structural basis of caspase-7 inhibition by XIAP. *Cell*, 2001, **104**, 69-80.
- Chang B.S., Minn A.J., Muchmore S.W., Fesik S.W., Thompson C.B. Identification of a novel regulatory domain in Bcl-X(L) and Bcl-2. *EMBO J.*, 1997, **16**, 968-977.
- Chang L., Karin M. Mammalian MAP kinase signalling cascades. *Nature*, 2001, **410**, 37-40.
- Charcot J.M., Robin C. Observation de leucocythémie. *C. R. Mem. Soc. Biol.*, 1853, **5**, 454-450.
- Chen F., Chang R., Trivedi M., Capetanaki Y., Cryns V.L. Caspase proteolysis of desmin produces a dominant-negative inhibitor of intermediate filaments and promotes apoptosis. *J. Biol. Chem.*, 2003, **278**, 6848-6853.
- Chen G., Ray R., Dubik D., Shi L., Cizeau J., Bleackley R.C., Saxena S., Gietz R.D., Greenberg A.H. The E1B 19K/Bcl-2-binding protein Nip3 is a dimeric mitochondrial protein that activates apoptosis. *J. Exp. Med.*, 1997, **186**, 1975-1983.
- Cheng E.H., Kirsch D.G., Clem R.J., Ravi R., Kastan M.B., Bedi A., Ueno K., Hardwick J.M. Conversion of Bcl-2 to a Bax-like death effector by caspases. *Science*, 1997, **278**, 1966-1968.
- Chetta A., Zanini A., Foresi A., D'Ippolito R., Tipa A., Castagnaro A., Baraldo S., Neri M., Saetta M., Olivieri D. Vascular endothelial growth factor up-regulation and bronchial wall remodelling in asthma. *Clin. Exp. Allergy*, 2005, **35**, 1437-1442.
- Chiba R., Nakagawa N., Kurasawa K., Tanaka Y., Saito Y., Iwamoto I. Ligation of CD31 (PECAM-1) on endothelial cells increases adhesive function of α v β 3 integrin and enhances β 1 integrin-mediated adhesion of eosinophils to endothelial cells. *Blood*, 1999, **94**, 1319-1329.
- Chihab R., Porn-Ares M.I., Alvarado-Kristensson M., Andersson T. Sphingosine 1-phosphate antagonizes human neutrophil apoptosis via p38 mitogen-activated protein kinase. *Cell. Mol. Life Sci.*, 2003, **60**, 776-785.
- Chinnaiyan A.M., O'Rourke K., Tewari M., Dixit V.M. FADD, a novel death domain-containing protein, interacts with the death domain of Fas and initiates apoptosis. *Cell*, 1995, **81**, 505-512.
- Clem R.J., Cheng E.H., Karp C.L., Kirsch D.G., Ueno K., Takahashi A., Kastan M.B., Griffin D.E., Earnshaw W.C., Veluona M.A., Hardwick J.M. Modulation of cell death by Bcl-XL through caspase interaction. *Proc. Natl. Acad. Sci. U. S. A.*, 1998, **95**, 554-559.
- Colotta F., Re F., Polentarutti N., Sozzani S., Mantovani A. Modulation of granulocyte survival and programmed cell death by cytokines and bacterial products. *Blood*, 1992, **80**, 2012-2020.
- Colussi P.A., Harvey N.L., Kumar S. Prodomain-dependent nuclear localization of the caspase-2 (Nedd2) precursor. A novel function for a caspase prodomain. *J. Biol. Chem.*, 1998, **273**, 24535-24542.
- Conesa A., Tassinari P., Aldrey O., Taylor P., Bianco N.E., De Sanctis J.B. Interleukin-2 induces peroxide production by primed normodense eosinophils of patients with asthma. *Allergy Asthma Proc.*, 2003, **24**, 27-33.
- Conroy D.M., Williams T.J. Eotaxin and the attraction of eosinophils to the asthmatic lung. *Respir. Res.*, 2001, **2**, 150-156.

- Coroneos E., Martinez M., McKenna S., Kester M. Differential regulation of sphingomyelinase and ceramidase activities by growth factors and cytokines. Implications for cellular proliferation and differentiation. *J. Biol. Chem.*, 1995, **270**, 23305-23309.
- Coroneos E., Wang Y., Panuska J.R., Templeton D.J., Kester M. Sphingolipid metabolites differentially regulate extracellular signal-regulated kinase and stress-activated protein kinase cascades. *Biochem. J.*, 1996, **316**, 13-17.
- Cowburn A.S., Cadwallader K.A., Reed B.J., Farahi N., Chilvers E.R. Role of PI3-kinase-dependent Bad phosphorylation and altered transcription in cytokine-mediated neutrophil survival. *Blood*, 2002, **100**, 2607-2616.
- Cox G. Glucocorticoid treatment inhibits apoptosis in human neutrophils. Separation of survival and activation outcomes. *J. Immunol.*, 1995, **154**, 4719-4725.
- Craxton A., Shu G., Graves J.D., Saklatvala J., Krebs E.G., Clark E.A. p38 MAPK is required for CD40-induced gene expression and proliferation in B lymphocytes. *J. Immunol.*, 1998, **161**, 3225-3236.
- Crompton M. The mitochondrial permeability transition pore and its role in cell death. *Biochem. J.*, 1999, **341**, 233-249.
- Crook N.E., Clem R.J., Miller L.K. An apoptosis-inhibiting baculovirus gene with a zinc finger-like motif. *J. Virol.*, 1993, **67**, 2168-2174.
- Cuvillier O., Pirianov G., Kleuser B., Vanek P.G., Coso O.A., Gutkind S., Spiegel S. Suppression of ceramide-mediated programmed cell death by sphingosine-1-phosphate. *Nature*, 1996, **381**, 800-803.
- Cuvillier O., Rosenthal D.S., Smulson M.E., Spiegel S. Sphingosine 1-phosphate inhibits activation of caspases that cleave poly(ADP-ribose) polymerase and lamins during Fas-and ceramide-mediated apoptosis in Jurkat T lymphocytes. *J. Biol. Chem.*, 1998, **273**, 2910-2916.
- Cuvillier O., Levade T. Sphingosine 1-phosphate antagonizes apoptosis of human leukemia cells by inhibiting release of cytochrome c and Smac/DIABLO from mitochondria. *Blood*, 2001, **98**, 2828-2836.

D

- Daeron M. Fc Receptor biology. *Annu. Rev. Immunol.*, 1997, **15**, 203-234.
- Daigle I., Simon H.U. Critical role for caspases 3 and 8 in neutrophil but not eosinophil apoptosis. *Int. Arch. Allergy Immunol.*, 2001a, **126**, 147-156.
- Daigle I., Simon H.U. Alternative functions for TRAIL receptors in eosinophils and neutrophils. *Swiss. Med. Wkly.*, 2001b, **131**, 231-237.
- Daniel N.N., Korsmeyer S.J. Cell death: critical control points. *Cell*, 2004, **116**, 205-219.
- Dancey J.T., Deubelbeiss K.A., Harker L.A., Finch C.A. Neutrophil kinetics in man. *J. Clin. Invest.*, 1976, **58**, 705-715.
- Datta S.R., Brunet A., Greenberg M.E. Cellular survival: a play in three Acts. *Genes Dev.*, 1999, **13**, 2905-2927.
- Daugas E., Susin S.A., Zamzami N., Ferri K.F., Irinopoulou T., Larochette N., Prevost M.C., Leber B., Andrews D., Penninger J., Kroemer G. Mitochondrio-nuclear translocation of AIF in apoptosis and necrosis. *FASEB J.*, 2000, **14**, 729-739.
- Dbaibo G.S., Pushkareva M.Y., Rachid R.A., Alter N., Smyth M.J., Obeid L.M., Hannun Y.A. p53-dependent ceramide response to genotoxic stress. *J. Clin. Invest.*, 1998, **102**, 329-339.
- De Maria R., Lenti L., Malisan F., d'Agostino F., Tomassini B., Zeuner A., Rippo M.R., Testi R. Requirement for GD3 ganglioside in CD95- and ceramide-induced apoptosis. *Science*, 1997, **277**, 1652-1655.
- del Peso L., Gonzalez-Garcia M., Page C., Herrera R., Nunez G. Interleukin-3-induced phosphorylation of BAD through the protein kinase Akt. *Science*, 1997, **278**, 687-689.

- Del Pozo V., De Andres B., Martin E., Cardaba B., Fernandez J.C., Gallardo S., Tramon P., Leyva-Cobian F., Palomino P., Lahoz C. Eosinophil as antigen-presenting cell: activation of T cell clones and T cell hybridoma by eosinophils after antigen processing. *Eur. J. Immunol.*, 1992, **22**, 1919-25.
- DeLeo F.R., Quinn M.T. Assembly of the phagocyte NADPH oxidase: molecular interaction of oxidase proteins. *J. Leukoc. Biol.*, 1996, **60**, 677-691.
- Delhalle S. Rôle du facteur de transcription NF- κ B dans l'apoptose des cellules cancéreuses. Thèse de Doctorat. Université de Liège (Faculté de Médecine) : Liège, 2002, 184p.
- Denburg J.A. The inflammatory response. *Am. J. Respir. Crit. Care Med.*, 1996, **153**, S11-S13.
- Desreumaux P., Janin A., Colombel J.F., Prin L., Plumas J., Emilie D., Torpier G., Capron A., Capron M. Interleukin 5 messenger RNA expression by eosinophils in the intestinal mucosa of patients with coeliac disease. *J. Exp. Med.*, 1992, **175**, 293-296.
- Deveraux Q.L., Reed J.C. IAP family proteins--suppressors of apoptosis. *Genes Dev.*, 1999, **13**, 239-252.
- Dewson G., Walsh G.M., Wardlaw A.J. Expression of Bcl-2 and its homologues in human eosinophils. Modulation by interleukin-5. *Am. J. Respir. Cell. Mol. Biol.*, 1999, **20**, 720-728.
- Dewson G., Cohen G.M., Wardlaw A.J. Interleukin-5 inhibits translocation of Bax to the mitochondria, cytochrome c release, and activation of caspases in human eosinophils. *Blood*, 2001, **98**, 2239-2247.
- Dibbert B., Daigle I., Braun D., Schranz C., Weber M., Blaser K., Zangemeister-Wittke U., Akbar A.N., Simon H.U. Role for Bcl-xL in delayed eosinophil apoptosis mediated by granulocyte-macrophage colony-stimulating factor and interleukin-5. *Blood*, 1998, **92**, 778-783.
- Dibbert B., Weber M., Nikolaizik W.H., Vogt P., Schöni M.H., Blaser K., Simon H.-U. Cytokine-mediated Bax deficiency and consequent delayed neutrophil apoptosis: a general mechanism to accumulate effector cells in inflammation. *Proc. Natl Acad. Sci. U. S. A.*, 1999, **96**, 13330-13335.
- Dobrowsky R.T., Kamibayashi C., Mumby M.C., Hannun Y.A. Ceramide activates heterotrimeric protein phosphatase 2A. *J. Biol. Chem.*, 1993, **268**, 15523-15530.
- Dobrowsky R.T. Sphingolipid signalling domains floating on rafts or buried in caves ? *Cell. Signal.*, 2000, **12**, 81-90.
- Dombrowicz D., Capron M. Eosinophils, allergy and parasites. *Curr. Opin. Immunol.*, 2001, **13**, 716-720.
- Doyle N.A., Bhagwan S.D., Meek B.B., Kutkoski G.J., Steeber D.A., Tedder T.F., Doerschuk C.M. Neutrophil margination, sequestration, and emigration in the lungs of L-selectin-deficient mice. *J. Clin. Invest.*, 1997, **99**, 526-533.
- Dransfield I., Buckle A.M., Savill J.S., McDowall A., Haslett C., Hogg N. Neutrophil apoptosis is associated with a reduction in CD16 (Fc gamma RIII) expression. *J. Immunol.*, 1994, **153**, 1254-1263.
- Droin N., Bichat F., Rebe C., Wotawa A., Sordet O., Hammann A., Bertrand R., Solary E. Involvement of caspase-2 long isoform in Fas-mediated cell death of human leukemic cells. *Blood*, 2001, **97**, 1835-1844.
- Druilhe A., Arock M., Le Goff L., Pretolani M. Human eosinophils express bcl-2 family proteins: modulation of Mcl-1 expression by IFN-gamma. *Am. J. Respir. Cell. Mol. Biol.*, 1998a, **18**, 315-322.
- Druilhe A., Wallaert B., Tsicopoulos A., Lapa e Silva J.R., Tillie-Leblond I., Tonnel A.B., Pretolani M. Apoptosis, proliferation, and expression of Bcl-2, Fas, and Fas ligand in bronchial biopsies from asthmatics. *Am. J. Respir. Cell. Mol. Biol.*, 1998b, **19**, 747-757.
- Druilhe A., Letuve S., Pretolani M. Eosinophil apoptosis in asthma. *Pathol. Biol. (Paris)*, 2000, **48**, 566-573.
- Du C., Fang M., Li Y., Li L., Wang X. Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. *Cell*, 2000, **102**, 33-42.

- Duan H., Dixit V.M. RAIDD is a new 'death' adaptor molecule. *Nature*, 1997, **385**, 86-89.
- Dubois P., Degraeve E., Vandenplas O. Asthma and airway hyperresponsiveness among Belgian conscripts, 1978-91. *Thorax*, 1998, **53**, 101-105.
- Duncan C.J., Lawrie A., Blaylock M.G., Douglas J.G., Walsh G.M. Reduced eosinophil apoptosis in induced sputum correlates with asthma severity. *Eur. Respir. J.*, 2003, **22**, 484-490.
- Durham S.R., Church M.K. Principles of allergy diagnosis. In: Allergy, 2nd Edn, Holgate S.T., Church M.K., Lichtenstein L.M. (Eds), London, UK : Mosby, 2001, pp. 3-16.

E

- Earnshaw W.C., Martins L.M., Kaufmann S.H. Mammalian caspases: structure, activation, substrates, and functions during apoptosis. *Annu. Rev. Biochem.*, 1999, **68**, 383-424.
- Edwards S.W., Hallett M.B. Seeing the wood for the trees: the forgotten role of neutrophils in rheumatoid arthritis. *Immunol. Today*, 1997, **18**, 320-324.
- Elsbach P. The bactericidal/permeability-increasing protein (BPI) in antibacterial host defense. *J. Leukoc. Biol.*, 1998, **64**, 14-18.
- Endo K., Oki E., Biedermann V., Kojima H., Yoshida K., Johannes F.J., Kufe D., Datta R. Proteolytic cleavage and activation of protein kinase C mu by caspase-3 in the apoptotic response of cells to 1-beta-D-arabinofuranosylcytosine and other genotoxic agents. *J. Biol. Chem.*, 2000, **275**, 18476-18481.
- Epling-Burnette P.K., Zhong B., Bai F., Jiang K., Bailey R.D., Garcia R., Jove R., Djeu J.Y., Loughran T.P. Jr, Wei S. Cooperative regulation of Mcl-1 by Janus kinase/STAT and phosphatidylinositol 3-kinase contribute to granulocyte-macrophage colony-stimulating factor-delayed apoptosis in human neutrophils. *J. Immunol.*, 2001, **166**, 7486-7495.
- Erhlich P. Über die spezifischen granulation des Blutes. *Arch. Anat. Physiol. Lpz. Physiol. Abt.*, 1879, **3**, 571-579.
- Erjefalt J.S., Uller L., Malm-Erjefalt M., Persson C.G. Rapid and efficient clearance of airway tissue granulocytes through transepithelial migration. *Thorax*, 2004, **59**, 136-143.

F

- Fadeel B., Ahlin A., Henter J.I., Orrenius S., Hampton M.B. Involvement of caspases in neutrophil apoptosis: regulation by reactive oxygen species. *Blood*, 1998, **92**, 4808-4818.
- Fadok V.A., Savill J.S., Haslett C., Bratton D.L., Doherty D.E., Campbell P.A., Henson P.M. Different populations of macrophages use either the vitronectin receptor or the phosphatidylserine receptor to recognize and remove apoptotic cells. *J. Immunol.*, 1992, **149**, 4029-4035.
- Fanger N.A., Liu C., Guyre P.M., Wardwell K., O'Neil J., Guo T.L., Christian T.P., Mudzinski S.P., Gosselin E.J. Activation of human T cells by major histocompatibility complex class II expressing neutrophils: proliferation in the presence of superantigen, but not tetanus toxoid. *Blood*, 1997, **89**, 4128-35.
- Fantuzzi G., Puren A.J., Harding M.W., Livingston D.J., Dinarello C.A. Interleukin-18 regulation of interferon gamma production and cell proliferation as shown in interleukin-1beta-converting enzyme (caspase-1)-deficient mice. *Blood*, 1998, **91**, 2118-2125.
- Farrell A.M., Uchida Y., Nagiec M.M., Harris I.R., Dickson R.C., Elias P.M., Holleran W.M. UVB irradiation up-regulates serine palmitoyltransferase in cultured human keratinocytes. *J. Lipid Res.*, 1998, **39**, 2031-2038.

- Fecho K., Cohen P.L. Fas ligand (gld)- and Fas (lpr)-deficient mice do not show alterations in the extravasation or apoptosis of inflammatory neutrophils. *J. Leukoc. Biol.*, 1998, **64**, 373-383.
- Ferland C., Guilbert M., Davoine F., Flamand N., Chakir J., Laviolette M. Eotaxin promotes eosinophil transmigration via the activation of the plasminogen-plasmin system. *J. Leukoc. Biol.*, 2001, **69**, 772-778.
- Fiévez L. Rôles du facteur de transcription STAT5 dans l'homéostasie des neutrophiles et l'inflammation. Presse de la Faculté de Médecine Vétérinaire, ULg, Liège, Belgique, 2006.
- Fillet M., Van Heugen J.C., Servais A.C., De Graeve J., Crommen J. Separation, identification and quantitation of ceramides in human cancer cells by liquid chromatography-electrospray ionization tandem mass spectrometry. *J. Chromatogr. A*, 2002, **949**, 225-233.
- Fisher G.H., Rosenberg F.J., Straus S.E., Dale J.K., Middleton L.A., Lin A.Y., Strober W., Lenardo M.J., Puck J.M. Dominant interfering Fas gene mutations impair apoptosis in a human autoimmune lymphoproliferative syndrome. *Cell*, 1995, **81**, 935-946.
- Flemming W. Über die Bildung von Richtungsfiguren in Säugethiereiern beim Untergang Graffscher Follikel. *Archiv. Anat. Entwicklungsgeschichte. Arch Anat. Physiol.*, 1885, 221-244.

G

- Gagnon E., Duclos S., Rondeau C., Chevet E., Cameron P.H., Steele-Mortimer O., Paiement J., Bergeron J.J., Desjardins M. Endoplasmic reticulum-mediated phagocytosis is a mechanism of entry into macrophages. *Cell*, 2002, **110**, 119-131.
- Ganz T., Selsted M.E., Szklarek D., Harwig S.S.L., Daher K., Bainton D.F., Lehrer R.I. Defensins. Natural peptide antibiotics of human neutrophils. *J. Clin. Invest.*, 1985, **76**, 1427-1435.
- Ganz T., Selsted M.E., Lehrer R.I. Defensins. *Eur. J. Haematol.*, 1990, **44**, 1-8.
- Gao G., Dou Q.P. N-terminal cleavage of bax by calpain generates a potent proapoptotic 18-kDa fragment that promotes bcl-2-independent cytochrome C release and apoptotic cell death. *J. Cell. Biochem.*, 2000, **80**, 53-72.
- Garcia-Ruiz C., Colell A., Mari M., Morales A., Fernandez-Checa J.C. Direct effect of ceramide on the mitochondrial electron transport chain leads to generation of reactive oxygen species. Role of mitochondrial glutathione. *J. Biol. Chem.*, 1997, **272**, 11369-11377.
- Gasmi L., McLennan A.G., Edwards S.W. The diadenosine polyphosphates Ap3A and Ap4A and adenosine triphosphate interact with granulocyte-macrophage colony-stimulating factor to delay neutrophil apoptosis: implications for neutrophil-platelet interactions during inflammation. *Blood*, 1996, **87**, 3442-3449.
- Gaulfield J.P., Korman G., Butterworth A.E., Hogan M., David J.R. The adherence of human neutrophils and eosinophils to schistosomula: Evidence for membrane fusion between cells and parasites. *J. Cell. Biol.*, 1980, **86**, 46-63.
- Geijsen N., Uings I.J., Pals C., Armstrong J., McKinnon M., Raaijmakers J.A., Lammers J.W., Koenderman L., Coffey P.J. Cytokine-specific transcriptional regulation through an IL-5R α interacting protein. *Science*, 2001, **293**, 1136-1138.
- Gelfand E.W. Inflammatory mediators in allergic rhinitis. *J. Allergy Clin. Immunol.*, 2004, **114**, S135-S138.
- Gentile V., Thomazy V., Piacentini M., Fesus L., Davies P.J. Expression of tissue transglutaminase in Balb-C 3T3 fibroblasts: effects on cellular morphology and adhesion. *J. Cell Biol.*, 1992, **119**, 463-474.

- Ghayur T., Hugunin M., Talanian R.V., Ratnofsky S., Quinlan C., Emoto Y., Pandey P., Datta R., Huang Y., Kharbanda S., Allen H., Kamen R., Wong W., Kufe D. Proteolytic activation of protein kinase C delta by an ICE/CED 3-like protease induces characteristics of apoptosis. *J. Exp. Med.*, 1996, **184**, 2399-2404.
- Giembycz M.A., Lindsay M.A. Pharmacology of the eosinophil. *Pharmacol. Rev.*, 1999, **51**, 213-340.
- Gilliet M., Soumelis V., Watanabe N., Hanabuchi S., Antonenko S., de Waal-Malefyt R., Liu Y.J. Human dendritic cells activated by TSLP and CD40L induce proallergic cytotoxic T cells. *J. Exp. Med.*, 2003, **197**, 1059-1063.
- Ginis I., Faller D.V. Protection from apoptosis in human neutrophils is determined by the surface of adhesion. *Am. J. Physiol.*, 1997, **272**, C295-C309.
- Girard D., Paquet M.E., Paquin R., Beaulieu A.D. Differential effects of interleukin-15 (IL-15) and IL-2 on human neutrophils: modulation of phagocytosis, cytoskeleton rearrangement, gene expression, and apoptosis by IL-15. *Blood*, 1996, **88**, 3176-3184.
- Girard D., Paquin R., Beaulieu A.D. Responsiveness of human neutrophils to interleukin-4: induction of cytoskeletal rearrangements, de novo protein synthesis and delay of apoptosis. *Biochem. J.*, 1997, **325**, 147-153.
- Glucksmann A. Cell deaths in normal vertebrate ontogeny. *Biol. Rev.*, 1951, **26**, 59-86.
- Goepel F., Weinmann P., Schymeinsky J., Walzog B. Identification of caspase-10 in human neutrophils and its role in spontaneous apoptosis. *J. Leukoc. Biol.*, 2004, **75**, 836-843.
- Goldkorn T., Balaban N., Shannon M., Chea V., Matsukuma K., Gilchrist D., Wang H., Chan C. H₂O₂ acts on cellular membranes to generate ceramide signaling and initiate apoptosis in tracheobronchial epithelial cells. *J. Cell. Sci.*, 1998, **111**, 3209-3220.
- Gouni A.S., Lamkhioued B., Ochiai K., Tanaka Y., Delaporte E., Capron A., Kinet J.P., Capron M. High-affinity IgE receptor on eosinophils is involved in defence against parasites. *Nature*, 1994, **367**, 183-186.
- Gouon-Evans V., Pollard J.W. Eotaxin is required for eosinophil homing into the stroma of the pubertal and cycling uterus. *Endocrinology*, 2001, **142**, 4515-4521.
- Goyal L. Cell death inhibition: keeping caspases in check. *Cell*, 2001, **104**, 805-808.
- Grandgirard D., Studer E., Monney L., Belser T., Fellay I., Borner C., Michel M.R. Alphaviruses induce apoptosis in Bcl-2-overexpressing cells: evidence for a caspase-mediated, proteolytic inactivation of Bcl-2. *EMBO J.*, 1998, **17**, 1268-1278.
- Grassme H., Cremesti A., Kolesnick R., Gulbins E. Ceramide-mediated clustering is required for CD95-DISC formation. *Oncogene*, 2003, **22**, 5457-5470.
- Graves J.D., Gotoh Y., Draves K.E., Ambrose D., Han D.K., Wright M., Chernoff J., Clark E.A., Krebs E.G. Caspase-mediated activation and induction of apoptosis by the mammalian Ste20-like kinase Mst1. *EMBO J.*, 1998, **17**, 2224-2234.
- Green D.R., Reed J.C. Mitochondria and apoptosis. *Science.*, 1998, **281**, 1309-1312.
- Grewal I.S., Flavell R.A. CD40 and CD154 in cell-mediated immunity. *Annu. Rev. Immunol.*, 1998, **16**, 111-135.
- Gross A., Jockel J., Wei M.C., Korsmeyer S.J. Enforced dimerization of BAX results in its translocation, mitochondrial dysfunction and apoptosis. *EMBO J.*, 1998, **17**, 3878-3885.
- Gross A., McDonnell J.M., Korsmeyer S.J. BCL-2 family members and the mitochondria in apoptosis. *Genes Dev.*, 1999, **13**, 1899-1911.
- Gu M., Kerwin J.L., Watts J.D., Aebersold R. Ceramide profiling of complex lipid mixtures by electrospray ionization mass spectrometry. *Anal. Biochem.*, 1997, **244**, 347-356.

- Guilbert M., Ferland C., Bossé M., Flamand N., Lavigne S., Laviolette M. 5-Oxo-6,8,11,14-eicosatetraenoic acid induces important eosinophil transmigration through basement membrane components. Comparison of normal and asthmatic eosinophils. *Am. J. Respir. Cell. Mol. Biol.*, 1999, **21**, 97-104.
- Gulbins E., Kolesnick R. Raft ceramide in molecular medicine. *Oncogene*, 2003, **22**, 7070-7077.
- Guthrie L.A., McPhail L.C., Henson P.M., Johnston R.B.Jr. Priming of neutrophils for enhanced release of oxygen metabolites by bacterial lipopolysaccharide. Evidence for increased activity of the superoxide-producing enzyme. *J. Exp. Med.*, 1984, **160**, 1656-1671.

H

- Haldar S., Jena N., Croce C.M. Inactivation of Bcl-2 by phosphorylation. *Proc. Natl. Acad. Sci. U. S. A.*, 1995, **92**, 4507-4511.
- Hall S.E., Savill J.S., Henson P.M., Haslett C. Apoptotic neutrophils are phagocytosed by fibroblasts with participation of the fibroblast vitronectin receptor and involvement of a mannose/fucose-specific lectin. *J. Immunol.*, 1994, **153**, 3218-3227.
- Hampton M.B., Kettle A.J., Winterbourn C.C. Inside the neutrophil phagosome: oxidants, myeloperoxidase, and bacterial killing. *Blood*, 1998a, **92**, 3007-3017.
- Hampton M.B., Fadeel B., Orrenius S. Redox regulation of the caspases during apoptosis. *Ann. N. Y. Acad. Sci.*, 1998b, **854**, 328-335.
- Hannah S., Mecklenburgh K., Rahman I., Bellingan G.J., Greening A., Haslett C., Chilvers E.R. Hypoxia prolongs neutrophil survival *in vitro*. *FEBS Lett.*, 1995, **372**, 233-237.
- Hannah S., Nadra I., Dransfield I., Pryde J.G., Rossi A.G., Haslett C. Constitutive neutrophil apoptosis in culture is modulated by cell density independently of $\beta 2$ integrin-mediated adhesion. *FEBS Lett.*, 1998, **421**, 141-146.
- Hannun Y.A. Functions of ceramide in coordinating cellular responses to stress. *Science*, 1996, **274**, 1855-1859.
- Harada H., Becknell B., Wilm M., Mann M., Huang L.J., Taylor S.S., Scott J.D., Korsmeyer S.J. Phosphorylation and inactivation of BAD by mitochondria-anchored protein kinase A. *Mol. Cell.*, 1999, **3**, 413-422.
- Harter L., Keel M., Hentze H., Steckholzer U., Ungethüm U., Trentz O., Ertel W. Spontaneous in contrast to CD95-induced neutrophil apoptosis is independent of caspase activity. *J. Trauma*, 2001, **50**, 982-988.
- Harwig S.S., Tan L., Qu X.D., Cho Y., Eisenhauer P.B., Lehrer R.I. Bactericidal properties of murine intestinal phospholipase A2. *J. Clin. Invest.*, 1995, **95**, 603-610.
- Haslett C., Savill J.S., Whyte M.K., Stern M., Dransfield I., Meagher L.C. Granulocyte apoptosis and the control of inflammation. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 1994, **345**, 327-333.
- Hebestreit H., Yousefi S., Balatti I., Weber M., Cramer R., Simon D., Hartung K., Schapowal A., Blaser K., Simon H.U. Expression and function of the Fas receptor on human blood and tissue eosinophils. *Eur. J. Immunol.*, 1996, **26**, 1775-1780.
- Hebestreit H., Dibbert B., Balatti I., Braun D., Schapowal A., Blaser K., Simon H.U. Disruption of fas receptor signaling by nitric oxide in eosinophils. *J. Exp. Med.*, 1998, **187**, 415-425.
- Hegde R., Srinivasula S.M., Zhang Z., Wassell R., Mukattash R., Cilenti L., DuBois G., Lazebnik Y., Zervos A.S., Fernandes-Alnemri T., Alnemri E.S. Identification of Omi/HtrA2 as a mitochondrial apoptotic serine protease that disrupts inhibitor of apoptosis protein-caspase interaction. *J. Biol. Chem.*, 2002, **277**, 432-438.

- Hendey B., Lawson M., Marcantonio E.E., Maxfield F.R. Intracellular calcium and calcineurin regulate neutrophil motility on vitronectin through a receptor identified by antibodies to integrins α v and β 3. *Blood*, 1996, **87**, 2038-2048.
- Herren B., Levkau B., Raines E.W., Ross R. Cleavage of beta-catenin and plakoglobin and shedding of VE-cadherin during endothelial apoptosis: evidence for a role for caspases and metalloproteinases. *Mol. Biol. Cell.*, 1998, **9**, 1589-1601.
- Hiraguri M., Miike S., Sano H., Kurasawa K., Saito Y., Iwamoto I. Granulocyte-macrophage colony-stimulating factor and IL-5 activate mitogen-activated protein kinase through Jak2 kinase and phosphatidylinositol 3-kinase in human eosinophils. *J. Allergy Clin. Immunol.*, 1997, **100**, S45-S51.
- Hogan S.P., Mould A., Kikutani H., Ramsay A.J., Foster P.S. Aeroallergen-induced eosinophilic inflammation, lung damage, and airways hyperreactivity in mice can occur independently of IL-4 and allergen-specific immunoglobulins. *J. Clin. Invest.*, 1997, **99**, 1329-1339.
- Hogan S.P., Mishra A., Brandt E.B., Foster P.S., Rothenberg M.E. A critical role for eotaxin in experimental oral antigen-induced eosinophilic gastrointestinal allergy. *Proc. Natl. Acad. Sci. U. S. A.*, 2000, **97**, 6681-6686.
- Holgate S.T. The epidemic of allergy and asthma. *Nature*, 1999, **402**, B2-B4.
- Hsu H., Xiong J., Goeddel D.V. The TNF receptor 1-associated protein TRADD signals cell death and NF-kappa B activation. *Cell*, 1995, **81**, 495-504.
- Hsu Y.T., Wolter K.G., Youle R.J. Cytosol-to-membrane redistribution of Bax and Bcl-X(L) during apoptosis. *Proc. Natl Acad. Sci. U. S. A.*, 1997, **94**, 3668-3672.

I

- Iking-Konert C., Cseko C., Wagner C., Stegmaier S., Andrassy K., Hansch G.M. Transdifferentiation of polymorphonuclear neutrophils: acquisition of CD83 and other functional characteristics of dendritic cells. *J. Mol. Med.*, 2001, **79**, 464-474.
- Iking-Konert C., Ostendorf B., Sander O., Jost M., Wagner C., Joosten L., Schneider M., Hansch G.M. Transdifferentiation of polymorphonuclear neutrophils to dendritic-like cells at the site of inflammation in rheumatoid arthritis: evidence for activation by T cells. *Ann. Rheum. Dis.*, 2005, **64**, 1436-1442.
- Ishihara K., Satoh I., Mue S., Ohuchi K. Possible participation of a JAK2 signaling pathway in recombinant rat interleukin-5-induced prolongation of rat eosinophil survival. *Biochim. Biophys. Acta*, 2001, **1536**, 73-84.
- Ito T., Deng X., Carr B., May W.S. Bcl-2 phosphorylation required for anti-apoptosis function. *J. Biol. Chem.*, 1997, **272**, 11671-11673.
- Iwabuchi K., Nagaoka I. Lactosylceramide-enriched glycosphingolipid signaling domain mediates superoxide generation from human neutrophils. *Blood*, 2002, **100**, 1454-1464.
- Iwai K., Kondo T., Watanabe M., Yabu T., Kitano T., Taguchi Y., Umehara H., Takahashi A., Uchiyama T., Okazaki T. Ceramide increases oxidative damage due to inhibition of catalase by caspase-3-dependent proteolysis in HL-60 cell apoptosis. *J. Biol. Chem.*, 2003, **278**, 9813-9822.

J

- Janicke R.U., Ng P., Sprengart M.L., Porter A.G. Caspase-3 is required for alpha-fodrin cleavage but dispensable for cleavage of other death substrates in apoptosis. *J. Biol. Chem.*, 1998, **273**, 15540-15545.
- Jayadev S., Hayter H.L., Andrieu N., Gamard C.J., Liu B., Balu R., Hayakawa M., Ito F., Hannun Y.A. Phospholipase A2 is necessary for tumor necrosis factor alpha-induced ceramide generation in L929 cells. *J. Biol. Chem.*, 1997, **272**, 17196-17203.
- Jones W.T. The blood-corpuscle considered in its different phases of development in the animal series. Memoir I Vertebrata. *Phil. Trans. R. Soc. Lond.*, 1846, **136**, 63-87.

K

- Kane D.J., Sarafian T.A., Anton R., Hahn H., Gralla E.B., Valentine J.S., Ord T., Bredesen D.E. Bcl-2 inhibition of neural death: decreased generation of reactive oxygen species. *Science*, 1993, **262**, 1274-1277.
- Kaneto H., Fujii J., Myint T., Miyazawa N., Islam K.N., Kawasaki Y., Suzuki K., Nakamura M., Tatsumi H., Yamasaki Y., Taniguchi N. Reducing sugars trigger oxidative modification and apoptosis in pancreatic beta-cells by provoking oxidative stress through the glycation reaction. *Biochem. J.*, 1996, **320**, 855-863.
- Kanfer J.N., Young O.M., Shapiro D., Brady R.O. The metabolism of sphingomyelin. I. Purification and properties of a sphingomyelin-cleaving enzyme from rat liver tissue. *J. Biol. Chem.*, 1966, **241**, 1081-1084.
- Kang J.J., Schaber M.D., Srinivasula S.M., Alnemri E.S., Litwack G., Hall D.J., Bjornsti M.A. Cascades of mammalian caspase activation in the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 1999, **274**, 3189-3198.
- Kankaanranta H., De Souza P.M., Barnes P.J., Salmon M., Giembycz M.A., Lindsay M.A. SB 203580, an inhibitor of p38 mitogen-activated protein kinase, enhance constitutive apoptosis of cytokine-deprived human eosinophils. *J. Pharmacol. Exp. Ther.*, 1999, **290**, 621-628.
- Kankaanranta H., Lindsay M.A., Giembycz M.A., Zhang, X., Moilanen E., Barnes P.J. Delayed eosinophil apoptosis in asthma. *J. Allergy Clin. Immunol.*, 2000, **106**, 77-83.
- Kanto T., Kalinski P., Hunter O.C., Lotze M.T., Amoscato A.A. Ceramide mediates tumor-induced dendritic cell apoptosis. *J. Immunol.*, 2001, **16**, 3773-3784.
- Kato M., Kephart G.M., Talley N.J., Wagner J.M., Sarr M.G., Bonno M., McGovern T.W., Gleich G.J. Eosinophil infiltration and degranulation in normal human tissue. *Anat. Rec.*, 1998, **252**, 418-425.
- Kato M., Kephart G.M., Morikawa A., Gleich G.J. Eosinophil infiltration and degranulation in normal human tissues: evidence for eosinophil degranulation in normal gastrointestinal tract. *Int. Arch. Allergy Immunol.*, 2001, **125 Suppl. 1**, 55-58.
- Keel M., Ungethum U., Steckholzer U., Niederer E., Hartung T., Trentz O., Ertel W. Interleukin-10 counterregulates proinflammatory cytokine-induced inhibition of neutrophil apoptosis during severe sepsis. *Blood*, 1997, **90**, 3356-3363.
- Kehry M.R. CD40-mediated signaling in B cells. Balancing cell survival, growth, and death. *J. Immunol.*, 1996, **156**, 2345-2348.
- Kelekar A., Thompson C.B. Bcl-2-family proteins: the role of the BH3 domain in apoptosis. *Trends Cell. Biol.*, 1998, **8**, 324-330.
- Kerr J.F., Wyllie A.H., Currie A.R. Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br. J. Cancer*, 1972, **26**, 239-257.
- Kerr J.F. History of the events leading to the formulation of the apoptosis concept. *Toxicology*, 2002, **181-182**, 471-474.

- Kilpatrick L.E., Lee J.Y., Haines K.M., Campbell D.E., Sullivan K.E., Korchak H.M. A role for PKC-delta and PI 3-kinase in TNF-alpha-mediated antiapoptotic signaling in the human neutrophil. *Am. J. Physiol. Cell. Physiol.*, 2002, **283**, C48-C57.
- Kimani G., Tonnesen M.G., Henson P.M. Stimulation of eosinophil adherence to human vascular endothelial cells *in vitro* by platelet-activating factor. *J. Immunol.*, 1988, **140**, 3161-3166.
- King P., Goodbourn S. STAT1 is inactivated by a caspase. *J. Biol. Chem.*, 1998, **273**, 8699-8704.
- Kirschnek S., Paris F., Weller M., Grassme H., Ferlinz K., Riehle A., Fuks Z., Kolesnick R., Gulbins E. CD95-mediated apoptosis *in vivo* involves acid sphingomyelinase. *J. Biol. Chem.*, 2000, **275**, 27316-27323.
- Kischkel F.C., Hellbardt S., Behrmann I., Germer M., Pawlita M., Krammer P.H., Peter M.E. Cytotoxicity-dependent APO-1 (Fas/CD95)-associated proteins form a death-inducing signaling complex (DISC) with the receptor. *EMBO J.*, 1995, **14**, 5579-5588.
- Kischkel F.C., Lawrence D.A., Tinel A., LeBlanc H., Virmani A., Schow P., Gazdar A., Blenis J., Arnott D., Ashkenazi A. Death receptor recruitment of endogenous caspase-10 and apoptosis initiation in the absence of caspase-8. *J. Biol. Chem.*, 2001, **276**, 46639-46646.
- Kisseleva T., Bhattacharya S., Braunstein J., Schindler C.W. Signaling through the JAK/STAT pathway, recent advances and future challenges. *Gene*, 2002, **285**, 1-24.
- Kita H., Gleich G.J. Eosinophils and IgE receptors: a continuing controversy. *Blood*, 1997, **89**, 3497-3501.
- Kitanaka C., Kuchino Y. Caspase-independent programmed cell death with necrotic morphology. *Cell Death Differ.*, 1999, **6**, 508-515.
- Klebanoff S.J., Olszowski S., Van Voorhis W.C., Ledbetter J.A., Waltersdorph A.M., Schlechte K.G. Effects of gamma-interferon on human neutrophils: protection from deterioration on storage. *Blood*, 1992, **80**, 225-234.
- Kluck R.M., Bossy-Wetzel E., Green D.R., Newmeyer D.D. The release of cytochrome c from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science*, 1997, **275**, 1132-1136.
- Knepper-Nicolai B., Savill J., Brown S.B. Constitutive apoptosis in human neutrophils requires synergy between calpains and the proteasome downstream of caspases. *J. Biol. Chem.*, 1998, **273**, 30530-30536.
- Knol E.F., Roos D. Mechanisms regulating eosinophil extravasation in asthma. *Eur. Respir. J.*, 1996, **9**, 136S-140S.
- Ko Y.G., Liu P., Pathak R.K., Craig L.C., Anderson R.G. Early effects of pp60(v-src) kinase activation on caveolae. *J. Cell. Biochem.*, 1998, **71**, 524-535.
- Kolesnick R.N. 1,2-Diacylglycerols but not phorbol esters stimulate sphingomyelin hydrolysis in GH3 pituitary cells. *J. Biol. Chem.*, 1987, **262**, 16759-16762.
- Kolesnick R.N., Krönke M. Regulation of ceramide production and apoptosis. *Annu. Rev. Physiol.*, 1998, **60**, 643-665.
- Kolesnick R. The therapeutic potential of modulating the ceramide/sphingomyelin pathway. *J. Clin. Invest.*, 2002, **110**, 3-8.
- Kolesnick R., Fuks Z. Radiation and ceramide-induced apoptosis. *Oncogene*, 2003, **22**, 5897-5906.
- Kothakota S., Azuma T., Reinhard C., Klippel A., Tang J., Chu K., McGarry T.J., Kirschner M.W., Kohts K., Kwiatkowski D.J., Williams L.T. Caspase-3-generated fragment of gelsolin: effector of morphological change in apoptosis. *Science*, 1997, **278**, 294-308.
- Krajewska M., Wang H.G., Krajewski S., Zapata J.M., Shabaik A., Gascoyne R., Reed J.C. Immunohistochemical analysis of *in vivo* patterns of expression of CPP32 (Caspase-3), a cell death protease. *Cancer Res.*, 1997, **57**, 1605-1613.

- Krajewski S., Tanaka S., Takayama S., Schibler M.J., Fenton W., Reed J.C. Investigation of the subcellular distribution of the bcl-2 oncoprotein: residence in the nuclear envelope, endoplasmic reticulum, and outer mitochondrial membranes. *Cancer Res.*, 1993, **53**, 4701-4714.
- Kroegel C., Costabel U., Matthys H. Mechanism of membrane damage mediated by eosinophil major basic protein. *Lancet*, 1987, **1**, 1380-1381.
- Kroegel C., Giembycz M.A., Barnes P.J. Characterization of eosinophil cell activation by peptides. Differential effects of substance P, melittin, and FMET-Leu-Phe. *J. Immunol.*, 1990, **145**, 2581-2587.
- Kroegel C., Virchow J.C. Jr., Luttmann W., Walker C., Warner J.A. Pulmonary immune cells in health and disease: the eosinophil leucocyte (Part I). *Eur. Respir. J.*, 1994a, **7**, 519-543.
- Kroegel C., Warner J.A., Virchow J.C. Jr., Matthys H. Pulmonary immune cells in health and disease: the eosinophil leucocyte (Part II). *Eur. Respir. J.*, 1994b, **7**, 743-760.
- Kurt-Jones E.A., Mandell L., Whitney C., Padgett A., Gosselin K., Newburger P.E., Finberg R.W. Role of toll-like receptor 2 (TLR2) in neutrophil activation: GM-CSF enhances TLR2 expression and TLR2-mediated interleukin 8 responses in neutrophils. *Blood*, 2002, **100**, 1860-1868.
- Kuwana T., Bouchier-Hayes L., Chipuk J.E., Bonzon C., Sullivan B.A., Green D.R., Newmeyer D.D. BH3 domains of BH3-only proteins differentially regulate Bax-mediated mitochondrial membrane permeabilization both directly and indirectly. *Mol. Cell.*, 2005, **17**, 525-535.

L

- Lauffenburger D.A., Horwitz A.F. Cell migration: a physically integrated molecular process. *Cell*, 1996, **84**, 359-369.
- Lavrik I.N., Golks A., Krammer P.H. Caspases: pharmacological manipulation of cell death. *J. Clin. Invest.*, 2005, **115**, 2665-2672.
- Lee A., Whyte M.K.B., Haslett C. Inhibition of apoptosis and prolongation of neutrophil functional longevity by inflammatory mediators. *J. Leukoc. Biol.*, 1993, **54**, 283-289.
- Lee H.H., Dadgostar H., Cheng Q., Shu J., Cheng G. NF-kappaB-mediated up-regulation of Bcl-x and Bfl-1/A1 is required for CD40 survival signaling in B lymphocytes. *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 9136-9141.
- Lee J.Y., Hannun Y.A., Obeid L.M. Ceramide inactivates cellular protein kinase Calpha. *J. Biol. Chem.*, 1996, **271**, 13169-13174.
- Lehrer R.I., Ganz T., Selsted M.E., Babior B.M., Curnutte J.T. Neutrophils and host defense. *Ann. Intern. Med.*, 1988, **109**, 127-142.
- Lei X.F., Ohkawara Y., Stampfli M.R., Mastruzzo C., Marr R.A., Snider D., Xing Z., Jordana M. Disruption of antigen-induced inflammatory responses in CD40 ligand knockout mice. *J. Clin. Invest.*, 1998, **101**, 1342-1353.
- Lendahl U., Orrenius S. [Sydney Brenner, Robert Horvitz and John Sulston. Winners of the 2002 Nobel Prize in medicine or physiology. Genetic regulation of organ development and programmed cell death]. *Lakartidningen*, 2002, **99**, 4026-4032.
- Lepine J., Messner H.A. Pluripotent hemopoietic progenitors (CFU-GEMM) in chronic myelogenous leukemia. *Int. J. Cell. Cloning*, 1983, **1**, 230-239.
- Letuve S., Druilhe A., Grandsaigne M., Aubier M., Pretolani M. Involvement of caspases and of mitochondria in Fas ligation-induced eosinophils apoptosis: modulation by interleukin-5 and interferon-gamma. *J. Leukoc. Biol.*, 2001, **70**, 767-775.

- Leuenroth S.J., Grutkoski P.S., Ayala A., Simms H.H. The loss of Mcl-1 expression in human polymorphonuclear leukocytes promotes apoptosis. *J. Leukoc. Biol.*, 2000a, **68**, 158-166.
- Leuenroth S.J., Grutkoski P.S., Ayala A., Simms H.H. Suppression of PMN apoptosis by hypoxia is dependent on Mcl-1 and MAPK activity. *Surgery*, 2000b, **128**, 171-177.
- Levkau B., Herren B., Koyama H., Ross R., Raines E.W. Caspase-mediated cleavage of focal adhesion kinase pp125FAK and disassembly of focal adhesions in human endothelial cell apoptosis. *J. Exp. Med.*, 1998, **187**, 579-586.
- Leyden E. Zur Kenntnis des bronchial-asthma. *Arch. Pathol. Anat.*, 1872, **54**, 324-352.
- Li H., Zhu H., Xu C.J., Yuan J. Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell*, 1998, **94**, 491-501.
- Li P., Nijhawan D., Budihardjo I., Srinivasula S.M., Ahmad M., Alnemri E.S., Wang X. Cytochrome c and dATP-dependent formation of Apaf-1/caspase-9 complex initiates an apoptotic protease cascade. *Cell*, 1997, **91**, 479-489.
- Li L.Y., Luo X., Wang X. Endonuclease G is an apoptotic DNase when released from mitochondria. *Nature*, 2001, **412**, 95-99.
- Liles W.C., Kiener P.A., Ledbetter J.A., Aruffo A., Klebanoff S.J. Differential expression of Fas (CD95) and Fas ligand on normal human phagocytes: implications for the regulation of apoptosis in neutrophils. *J. Exp. Med.*, 1996, **184**, 429-440.
- Liston P., Fong W.G., Kelly N.L., Toji S., Miyazaki T., Conte D., Tamai K., Craig C.G., McBurney M.W., Korneluk R.G. Identification of XAF1 as an antagonist of XIAP anti-Caspase activity. *Nat. Cell. Biol.*, 2001, **3**, 128-133.
- Liu B., Hannun Y.A. Inhibition of the neutral magnesium-dependent sphingomyelinase by glutathione. *J. Biol. Chem.*, 1997, **272**, 16281-16287.
- Liu L.Y., Mathur S.K., Sedgwick J.B., Jarjour N.N., Busse W.W., Kelly E.A., Padigel U.M., Lee J.J., Nolan T.J., Schad G.A., Abraham D. Human airway and peripheral blood eosinophils enhance Th1 and Th2 cytokine secretion. *Allergy*, 2006, **61**, 589-597.
- Liu P., Anderson R.G. Compartmentalized production of ceramide at the cell surface. *J. Biol. Chem.*, 1995, **270**, 27179-27185.
- Loike J.D., Cao L., Budhu S., Marcantonio E.E., El Khoury J., Hoffman S., Yednock T.A., Silverstein S.C. Differential regulation of beta1 integrins by chemoattractants regulates neutrophil migration through fibrin. *J. Cell. Biol.*, 1999, **144**, 1047-1056.
- Longthorne V.L., Williams G.T. Caspase activity is required for commitment to Fas-mediated apoptosis. *EMBO J.*, 1997, **16**, 3805-3812.
- Lopez A.D., Murray C.C. The global burden of disease, 1990-2020. *Nat. Med.*, 1998, **4**, 1241-1243.
- Lukacs N.W., Strieter R.M., Kunkel S.L. Leukocyte infiltration in allergic airway inflammation. *Am. J. Respir. Cell Mol. Biol.*, 1995, **13**, 1-6.

M

- MacKinnon A.C., Buckley A., Chilvers E.R., Rossi A.G., Haslett C., Sethi T. Sphingosine kinase: a point of convergence in the action of diverse neutrophil priming agents. *J. Immunol.*, 2002, **169**, 6394-6400.
- Maianski N.A., Roos D., Kuijpers T.W. Tumor necrosis factor alpha induces a caspase-independent death pathway in human neutrophils. *Blood*, 2003, **101**, 1987-1995.

- Malisan F., Testi R. GD3 ganglioside and apoptosis. *Biochim. Biophys. Acta*, 2002, **1585**, 179-187.
- Mancini M., Nicholson D.W., Roy S., Thornberry N.A., Peterson E.P., Casciola-Rosen L.A., Rosen A. The caspase-3 precursor has a cytosolic and mitochondrial distribution: implications for apoptotic signaling. *J. Cell Biol.*, 1998, **140**, 1485-1495.
- Mansat V., Laurent G., Levade T., Bettaieb A., Jaffrezou J.P. The protein kinase C activators phorbol esters and phosphatidylserine inhibit neutral sphingomyelinase activation, ceramide generation, and apoptosis triggered by daunorubicin. *Cancer Res.*, 1997, **57**, 5300-5304.
- Marchesini N., Hannun Y.A. Acid and neutral sphingomyelinases: roles and mechanisms of regulation. *Biochem. Cell Biol.*, 2004, **82**, 27-44.
- Martin M.C., Allan L.A., Lickrish M., Sampson C., Morrice N., Clarke P.R. Protein kinase A regulates caspase-9 activation by Apaf-1 downstream of cytochrome c. *J. Biol. Chem.*, 2005, **280**, 15449-15455.
- Martinou I., Desagher S., Eskes R., Antonsson B., Andre E., Fakan S., Martinou J.C. The release of cytochrome c from mitochondria during apoptosis of NGF-deprived sympathetic neurons is a reversible event. *J. Cell Biol.*, 1999, **144**, 883-889.
- Martins L.M., Kottke T.J., Kaufmann S.H., Earnshaw W.C. Phosphorylated forms of activated caspases are present in cytosol from HL-60 cells during etoposide-induced apoptosis. *Blood*, 1998, **92**, 3042-3049.
- Marzo I., Brenner C., Zamzami N., Jurgensmeier J.M., Susin S.A., Vieira H.L., Prevost M.C., Xie Z., Matsuyama S., Reed J.C., Kroemer G. Bax and adenine nucleotide translocator cooperate in the mitochondrial control of apoptosis. *Science*, 1998a, **281**, 2027-2031.
- Marzo I., Brenner C., Zamzami N., Susin S.A., Beutner G., Brdiczka D., Remy R., Xie Z.H., Reed J.C., Kroemer G. The permeability transition pore complex: a target for apoptosis regulation by caspases and bcl-2-related proteins. *J. Exp. Med.*, 1998b, **187**, 1261-1271.
- Mashima T., Naito M., Tsuruo T. Caspase-mediated cleavage of cytoskeletal actin plays a positive role in the process of morphological apoptosis. *Oncogene*, 1999, **18**, 2423-2430.
- Mathias S., Dressler K.A., Kolesnick R.N. Characterization of a ceramide-activated protein kinase: stimulation by tumor necrosis factor alpha. *Proc. Natl. Acad. Sci. U. S. A.*, 1991, **88**, 10009-10013.
- Mathias S., Pena L.A., Kolesnick R.N. Signal transduction of stress via ceramide. *Biochem. J.*, 1998, **335**, 465-480.
- Matsuda H., Suda T., Sato J., Nagata T., Koide Y., Chida K., Nakamura H. alpha-Galactosylceramide, a ligand of natural killer T cells, inhibits allergic airway inflammation. *Am. J. Respir. Cell. Mol. Biol.*, 2005, **33**, 22-31.
- McConkey D.J. Calcium-dependent, interleukin 1-converting enzyme inhibitor-insensitive degradation of lamin B1 and DNA fragmentation in isolated thymocyte nuclei. *J. Biol. Chem.*, 1996, **271**, 22398-22406.
- McGinnis K.M., Whitton M.M., Gnegy M.E., Wang K.K. Calcium/calmodulin-dependent protein kinase IV is cleaved by caspase-3 and calpain in SH-SY5Y human neuroblastoma cells undergoing apoptosis. *J. Biol. Chem.*, 1998, **273**, 19993-20000.
- MacKinnon A.C., Buckley A., Chilvers E.R., Rossi A.G., Haslett C., Sethi T. Sphingosine kinase: a point of convergence in the action of diverse neutrophil priming agents. *J. Immunol.*, 2002, **169**, 6394-6400.
- Meagher L.C., Cousin J.M., Seckl J.R., Haslett C. Opposing effects of glucocorticoids on the rate of apoptosis in neutrophilic and eosinophilic granulocytes. *J. Immunol.*, 1996, **156**, 4422-4428.
- Mecklenburgh K.I., Walmsley S.R., Cowburn A.S., Wiesener M., Reed B.J., Upton P.D., Deighton J., Greening A.P., Chilvers E.R. Involvement of a ferroprotein sensor in hypoxia-mediated inhibition of neutrophil apoptosis. *Blood*, 2002, **100**, 3008-3016.

- Meeusen E.N., Balic A. Do eosinophils have a role in the killing of helminth parasites ? *Parasitol. Today*, 2000, **16**, 95-101.
- Mehlhop P.D., van de Rijn M., Brewer J.P., Kisselgof A.B., Geha R.S., Oettgen H.C., Martin T.R. CD40L, but not CD40, is required for allergen-induced bronchial hyperresponsiveness in mice. *Am. J. Respir. Cell. Mol. Biol.*, 2000, **23**, 646-651.
- Michel C., van Echten-Deckert G., Rother J., Sandhoff K., Wang E., Merrill A.H.Jr. Characterization of ceramide synthesis. A dihydroceramide desaturase introduces the 4,5-trans-double bond of sphingosine at the level of dihydroceramide. *J. Biol. Chem.*, 1997, **272**, 22432-22437.
- Miike S., Nakao A., Hiraguri M., Kurasawa K., Saito Y., Iwamoto I. Involvement of JAK2, but not PI 3-kinase/Akt and MAP kinase pathways, in anti-apoptotic signals of GM-CSF in human eosinophils. *J. Leukoc. Biol.*, 1999, **65**, 700-706.
- Mizgerd J.P., Meek B.B., Kutkoski G.J., Bullard D.C., Beaudet A.L., Doerschuk C.M. Selectins and neutrophil traffic: margination and Streptococcus pneumoniae-induced emigration in murine lungs. *J. Exp. Med.*, 1996, **184**, 639-645.
- Moore K.L., Patel K.D., Bruehl R.E., Li F., Johnson D.A., Lichenstein H.S., Cummings R.D., Bainton D.F., McEver R.P. P-selectin glycoprotein ligand-1 mediates rolling of human neutrophils on P-selectin. *J. Cell. Biol.*, 1995, **128**, 661-671.
- Morishima Y., Ishii Y., Kimura T., Shibuya A., Shibuya K., Hegab A.E., Iizuka T., Kiwamoto T., Matsuno Y., Sakamoto T., Nomura A., Taniguchi M., Sekizawa K. Suppression of eosinophilic airway inflammation by treatment with alpha-galactosylceramide. *Eur. J. Immunol.*, 2005, **35**, 2803-2814.
- Mosimann S.C., Newton D.L., Youle R.J., James M.N. X-ray crystallographic structure of recombinant eosinophil-derived neurotoxin at 1.83 Å resolution. *J. Mol. Biol.*, 1996, **260**, 540-552.
- Moulding D.A., Quayle J.A., Hart C.A., Edwards S.W. Mcl-1 expression in human neutrophils : regulation by cytokines and correlation with cell survival. *Blood*, 1998, **92**, 2495-2502.
- Moulding D.A., Giles R.V., Spiller D.G., White M.R., Tidd D.M., Edwards S.W. Apoptosis is rapidly triggered by antisense depletion of MCL-1 in differentiating U937 cells. *Blood*, 2000, **96**, 1756-1763.
- Moulding D.A., Akgul C., Derouet M., White M.R.H., Edwards S.W. Bcl-2 family expression in human neutrophils during delayed and accelerated apoptosis. *J. Leukoc. Biol.*, 2001, **70**, 783-792.
- Murray J., Barbara J.A., Dunkley S.A., Lopez A.F., Van Ostade X., Condliffe A.M., Dransfield I., Haslett C., Chilvers E.R. Regulation of neutrophil apoptosis by tumor necrosis factor-alpha: requirement for TNFR55 and TNFR75 for induction of apoptosis *in vitro*. *Blood*, 1997, **90**, 2772-2783.
- Muzio M., Salvesen G.S., Dixit V.M. FLICE induced apoptosis in a cell-free system. Cleavage of caspase zymogens. *J. Biol. Chem.*, 1997, **272**, 2952-2956.

N

- Nagata S. Apoptosis by death factor. *Cell*, 1997, **88**, 355-365.
- Nathan C.F. Neutrophil activation on biological surfaces. Massive secretion of hydrogen peroxide in response to products of macrophages and lymphocytes. *J. Clin. Invest.*, 1987, **80**, 1550-1560.
- Nathan R.A. The burden of allergic rhinitis. *Allergy Asthma Proc.*, 2007, **28**, 3-9.
- Nechushtan A., Smith C.L., Hsu Y.T., Youle R.J. Conformation of the Bax C-terminus regulates subcellular location and cell death. *EMBO J.*, 1999, **18**, 2330-2341.

Nhan T.Q., Liles W.C., Schwartz S.M. Physiological functions of caspases beyond cell death. *Am. J. Pathol.*, 2006, **169**, 729-737.

Nobel C.S., Kimland M., Nicholson D.W., Orrenius S., Slater A.F. Disulfiram is a potent inhibitor of proteases of the caspase family. *Chem. Res. Toxicol.*, 1997, **10**, 1319-1324.

Nojiri H., Takaku F., Ohta M., Miura Y., Saito M. Changes in glycosphingolipid composition during differentiation of human leukemic granulocytes in chronic myelogenous leukemia compared with in vitro granulocytic differentiation of human promyelocytic leukemia cell line HL-60. *Cancer Res.*, 1985, **45**, 6100-6106.

O

Obeid L.M., Linardic C.M., Karolak L.A., Hannun Y.A. Programmed cell death induced by ceramide. *Science*, 1993, **259**, 1769-1771.

Ochiai K., Kagami M., Matsumura R., Tomioka H. IL-5 but not interferon-gamma (IFN-gamma) inhibits eosinophil apoptosis by up-regulation of bcl-2 expression. *Clin. Exp. Immunol.*, 1997, **107**, 198-204.

Oehler L., Majdic O., Pickl W.F., Stockl J., Riedl E., Drach J., Rappersberger K., Geissler K., Knapp W. Neutrophil granulocyte-committed cells can be driven to acquire dendritic cell characteristics. *J. Exp. Med.*, 1998, **187**, 1019-1028.

Ogata N., Kikuchi Y., Kouro T., Tomonaga M., Takatsu K. The activation of the JAK2/STAT5 pathway is commonly involved in signalling through the human IL-5 receptor. *Int. Arch. Allergy Immunol.*, 1997, **114**, 24-27.

Ogretmen B., Hannun Y.A. Updates on functions of ceramide in chemotherapy-induced cell death and in multidrug resistance. *Drug Resist. Updat.*, 2001, **4**, 368-377.

Ogretmen B., Hannun Y.A. Biologically active sphingolipids in cancer pathogenesis and treatment. *Nat. Rev. Cancer*, 2004, **4**, 604-616.

Ohkawara Y., Lim K.G., Xing Z., Glibetic M., Nakano K., Dolovich J., Croitoru K., Weller P.F., Jordana M. CD40 expression by human peripheral blood eosinophils. *J. Clin. Invest.*, 1996, **97**, 1761-1766.

Oltvai Z.N., Milliman C.L., Korsmeyer S.J. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell.*, 1993, **74**, 609-619.

Oppenheim J.J., Zachariae C.O., Mukaida N., Matsushima K. Properties of the novel proinflammatory supergene "intercrine" cytokine family. *Annu. Rev. Immunol.*, 1991, **9**, 617-648.

Oyadomari S., Araki E., Mori M. Endoplasmic reticulum stress-mediated apoptosis in pancreatic beta-cells. *Apoptosis*, 2002, **7**, 335-345.

P

Park H.Y., Song M.G., Lee J.S., Kim J.W., Jin J.O., Park J.I., Chang Y.C., Kwak J.Y. Apoptosis of human neutrophils induced by protein phosphatase 1/2A inhibition is caspase-independent and serine protease-dependent. *J. Cell. Physiol.*, 2007, [Epub ahead of print].

Parkos C.A. Molecular events in neutrophil transepithelial migration. *Bioessays*, 1997, **19**, 865-873.

Paumen M.B., Ishida Y., Muramatsu M., Yamamoto M., Honjo T. Inhibition of carnitine palmitoyltransferase I augments sphingolipid synthesis and palmitate-induced apoptosis. *J. Biol. Chem.*, 1997, **272**, 3324-3329.

- Pazdrak K., Schreiber D., Forsythe P., Justement L., Alam R. The intracellular signal transduction mechanism of interleukin 5 in eosinophils: the involvement of lyn tyrosine kinase and the Ras-Raf-1-MEK-microtubule-associated protein kinase pathway. *J. Exp. Med.*, 1995, **181**, 1827-1834.
- Pazdrak K., Adachi T., Alam R. Src homology 2 protein tyrosine phosphatase (SHPTP2)/Src homology 2 phosphatase 2 (SHP2) tyrosine phosphatase is a positive regulator of the interleukin 5 receptor signal transduction pathways leading to the prolongation of eosinophils survival. *J. Exp. Med.*, 1997, **186**, 561-568.
- Peacock C.D., Misso N.L., Watkins D.N., Thompson P.J. PGE 2 and dibutyryl cyclic adenosine monophosphate prolong eosinophil survival in vitro. *J. Allergy Clin. Immunol.*, 1999, **104**, 153-162.
- Pena L.A., Fuks Z., Kolesnick R. Stress-induced apoptosis and the sphingomyelin pathway. *Biochem. Pharmacol.*, 1997, **53**, 615-621.
- Perianin A., Snyderman R. Mastoparan, a wasp venom peptide, identifies two discrete mechanisms for elevating cytosolic calcium and inositol trisphosphates in human polymorphonuclear leukocytes. *J. Immunol.*, 1989, **143**, 1669-1673.
- Pericle F., Liu J.H., Diaz J.I., Blanchard D.K., Wei S., Forni G., Djeu J.Y. Interleukin-2 prevention of apoptosis in human neutrophils. *Eur. J. Immunol.*, 1994, **24**, 440-444.
- Perry D.K., Hannun Y.A. The role of ceramide in cell signaling. *Biochim. Biophys. Acta*, 1998, **1436**, 233-243.
- Perry D.K., Carton J., Shah A.K., Meredith F., Uhlinger D.J., Hannun Y.A. Serine palmitoyltransferase regulates de novo ceramide generation during etoposide-induced apoptosis. *J. Biol. Chem.*, 2000, **275**, 9078-9084.
- Perry D.K., Kolesnick R.N. Ceramide and sphingosine 1-phosphate in anti-cancer therapies. *Cancer. Treat. Res.*, 2003, **115**, 345-354.
- Peter M.E., Heufelder A.E., Hengartner M.O. Advances in apoptosis research. *Proc. Natl Acad. Sci. U. S. A.*, 1997, **94**, 12736-12737.
- Petit P.X., Zamzami N., Vayssiere J.L., Mignotte B., Kroemer G., Castedo M. Implication of mitochondria in apoptosis. *Mol. Cell. Biochem.*, 1997, **174**, 185-188.
- Pitti R.M., Marsters S.A., Ruppert S., Donahue C.J., Moore A., Ashkenazi A. Induction of apoptosis by Apo-2 ligand, a new member of the tumor necrosis factor cytokine family. *J. Biol. Chem.*, 1996, **271**, 12687-12690.
- Plo I., Ghandour S., Feutz A.C., Clanet M., Laurent G., Bettaieb A. Involvement of de novo ceramide biosynthesis in lymphotoxin-induced oligodendrocyte death. *Neuroreport*, 1999, **10**, 2373-2376.
- Poommipanit P.B., Chen B., Oltvai Z.N. Interleukin-3 induces the phosphorylation of a distinct fraction of bcl-2. *J. Biol. Chem.*, 1999, **274**, 1033-1039.
- Puthalakath H., Huang D.C., O'Reilly L.A., King S.M., Strasser A. The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol. Cell.*, 1999, **3**, 287-296.
- Puxeddu I., Ribatti D., Crivellato E., Levi-Schaffer F. Mast cells and eosinophils: a novel link between inflammation and angiogenesis in allergic diseases. *J. Allergy Clin. Immunol.*, 2005, **116**, 531-536.
- Pyne S. Cellular signaling by sphingosine and sphingosine 1-phosphate. Their opposing roles in apoptosis. *Subcell. Biochem.*, 2002, **36**, 245-268.
- Pyne S., Pyne N.J. Sphingosine 1-phosphate signalling and termination at lipid phosphate receptors. *Biochim. Biophys. Acta*, 2002, **1582**, 121-131.

Qin Z.H., Wang Y., Kikly K.K., Sapp E., Kegel K.B., Aronin N., DiFiglia M. Pro-caspase-8 is predominantly localized in mitochondria and released into cytoplasm upon apoptotic stimulation. *J. Biol. Chem.*, 2001, **276**, 8079-8086.

R

Rao L., Perez D., White E. Lamin proteolysis facilitates nuclear events during apoptosis. *J. Cell. Biol.*, 1996, **135**, 1441-1455.

Ravi R., Bedi A., Fuchs E.J., Bedi A. CD95 (Fas)-induced caspase-mediated proteolysis of NF-kappaB. *Cancer Res.*, 1998, **58**, 882-886.

Reed C.J. Apoptosis and cancer: strategies for integrating programmed cell death. *Semin. Hematol.*, 2000, **37**, 9-16.

Reed J.C. Splicing and dicing apoptosis genes. *Nat. Biotechnol.*, 1999, **17**, 1064-1065.

Reynolds C.P., Maurer B.J., Kolesnick R.N. Ceramide synthesis and metabolism as a target for cancer therapy. *Cancer Lett.*, 2004, **206**, 169-180.

Rippo M.R., Malisan F., Ravagnan L., Tomassini B., Condo I., Costantini P., Susin S.A., Rufini A., Todaro M., Kroemer G., Testi R. GD3 ganglioside directly targets mitochondria in a bcl-2-controlled fashion. *FASEB J.*, 2000, **14**, 2047-2054.

Rosenberg H.F., Tenen D.G., Ackerman S.J. Molecular cloning of the human eosinophil-derived neurotoxin: a member of the ribonuclease gene family. *Proc. Natl. Acad. Sci. U. S. A.*, 1989, **86**, 4460-4464.

Rothe M., Pan M.G., Henzel W.J., Ayres T.M., Goeddel D.V. The TNFR2-TRAF signaling complex contains two novel proteins related to baculoviral inhibitor of apoptosis proteins. *Cell*, 1995, **83**, 1243-1252.

Rothenberg M.E. Eosinophilia. *N. Engl. J. Med.*, 1998, **338**, 1592-1600.

Rothenberg M.E., Mishra A., Brandt E.B., Hogan S.P. Gastrointestinal eosinophils. *Immunol. Rev.*, 2001, **179**, 139-155.

Rotrosen D. The respiratory burst oxidase. In: Inflammation: Basic principles and clinical correlates, Gallin J.I., Goldstein I.M., Snyderman R. (Eds), New York : Raven Press, 1992, p 589.

Roussel E., Gingras M.C. Transendothelial migration induces rapid expression on neutrophils of granule-release VLA6 used for tissue infiltration. *J. Leukoc. Biol.*, 1997, **62**, 356-362.

Roy N., Mahadevan M.S., McLean M., Shutler G., Yaraghi Z., Farahani R., Baird S., Besner-Johnston A., Lefebvre C., Kang X., et al. The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. *Cell*, 1995, **80**, 167-178.

Rudel T., Bokoch G.M. Membrane and morphological changes in apoptotic cells regulated by caspase-mediated activation of PAK2. *Science*, 1997, **276**, 1571-1574.

Ruvolo P.P., Deng X., Carr B.K., May W.S. A functional role for mitochondrial protein kinase Calpha in Bcl2 phosphorylation and suppression of apoptosis. *J. Biol. Chem.*, 1998, **273**, 25436-25442.

Ruvolo P.P., Deng X., Ito T., Carr B.K., May W.S. Ceramide induces Bcl2 dephosphorylation via a mechanism involving mitochondrial PP2A. *J. Biol. Chem.*, 1999, **274**, 20296-20300.

S

Sakahira H., Enari M., Nagata S. Cleavage of CAD inhibitor in CAD activation and DNA degradation during apoptosis. *Nature*, 1998, **391**, 96-99.

- Salamone G., Giordano M., Trevani A.S., Gamberale R., Vermeulen M., Schettinni J., Geffner J.R. Promotion of neutrophil apoptosis by TNF-alpha. *J. Immunol.*, 2001, **166**, 3476-3483.
- Salinas M., Lopez-Valdaliso R., Martin D., Alvarez A., Cuadrado A. Inhibition of PKB/Akt1 by C2-ceramide involves activation of ceramide-activated protein phosphatase in PC12 cells. *Mol. Cell. Neurosci.*, 2000, **15**, 156-169.
- Salvesen G., Farley D., Shuman J., Przybyla A., Reilly C., Travis J. Molecular cloning of human cathepsin G: structural similarity to mast cell and cytotoxic T lymphocyte proteinases. *Biochemistry*, 1987, **26**, 2289-2293.
- Salvesen G.S., Duckett C.S. IAP proteins : blocking the road to death's door. *Mol. Cell. Biol.*, 2002, **3**, 401-410.
- Samejima K., Tone S., Earnshaw W.C. CAD/DFF40 nuclease is dispensable for high molecular weight DNA cleavage and stage I chromatin condensation in apoptosis. *J. Biol. Chem.*, 2001, **276**, 45427-45432.
- Santana P., Pena L.A., Haimovitz-Friedman A., Martin S., Green D., McLoughlin M., Cordon-Cardo C., Schuchman E.H., Fuks Z., Kolesnick R. Acid sphingomyelinase-deficient human lymphoblasts and mice are defective in radiation-induced apoptosis. *Cell*, 1996, **86**, 189-199.
- Santoro M.F., Annand R.R., Robertson M.M., Peng Y.W., Brady M.J., Mankovich J.A., Hackett M.C., Ghayur T., Walter G., Wong W.W., Giegel D.A. Regulation of protein phosphatase 2A activity by caspase-3 during apoptosis. *J. Biol. Chem.*, 1998, **273**, 13119-13128.
- Savill J.S., Wyllie A.H., Henson J.E., Walport M.J., Henson P.M., Haslett C. Macrophage phagocytosis of aging neutrophils in inflammation. Programmed cell death in the neutrophil leads to its recognition by macrophages. *J. Clin. Invest.*, 1989, **83**, 865-875.
- Savill J., Dransfield I., Hogg N., Haslett C. Vitronectin receptor-mediated phagocytosis of cells undergoing apoptosis. *Nature*, 1990, **343**, 170-173.
- Savill J., Fadok V., Henson P., Haslett C. Phagocyte recognition of cells undergoing apoptosis. *Immunol. Today*, 1993, **14**, 131-136.
- Savill J., Haslett C. Granulocyte clearance by apoptosis in the resolution of inflammation. *Semin. Cell. Biol.*, 1995, **6**, 385-393.
- Savill J., Dransfield I., Gregory C., Haslett C. A blast from the past : clearance of apoptotic cells regulates immune responses. *Nat. Rev. Immunol.*, 2002, **2**, 965-975.
- Sawai H., Okazaki T., Yamamoto H., Okano H., Takeda Y., Tashima M., Sawada H., Okuma M., Ishikura H., Umehara H., Domae N. Requirement of AP-1 for ceramide-induced apoptosis in human leukemia HL-60 cells. *J. Biol. Chem.*, 1995, **270**, 27326-27331.
- Scheel-Toellner D., Wang K., Craddock R., Webb P.R., McGettrick H.M., Assi L.K., Parkes N., Clough L.E., Gulbins E., Salmon M., Lord J.M. Reactive oxygen species limit neutrophil life span by activating death receptor signaling. *Blood*, 2004a, **104**, 2557-2564.
- Scheel-Toellner D., Wang K.Q., Webb P.R., Wong S.H., Craddock R., Assi L.K., Salmon M., Lord J.M. Early events in spontaneous neutrophil apoptosis. *Biochem. Soc. Trans.*, 2004b, **32**, 461-464.
- Schimmer A.D. Inhibitor of apoptosis proteins: translating basic knowledge into clinical practice. *Cancer Res.*, 2004, **64**, 7183-7190.
- Schmidt H.H., Seifert R., Bohme E. Formation and release of nitric oxide from human neutrophils and HL-60 cells induced by a chemotactic peptide, platelet activating factor and leukotriene B4. *FEBS Lett.*, 1989, **244**, 357-360.
- Schwandner R., Wiegmann K., Bernardo K., Kreder D., Kronke M. TNF receptor death domain-associated proteins TRADD and FADD signal activation of acid sphingomyelinase. *J. Biol. Chem.*, 1998, **273**, 5916-5922.

- Scott D.W., Livnat D., Pennell C.A., Keng P. Lymphoma models for B cell activation and tolerance. III. Cell cycle dependence for negative signalling of WEHI-231 B lymphoma cells by anti- μ . *J. Exp. Med.*, 1986, **164**, 156-164.
- Sebbagh M., Renvoize C., Hamelin J., Riche N., Bertoglio J., Breard J. Caspase-3-mediated cleavage of ROCK I induces MLC phosphorylation and apoptotic membrane blebbing. *Nat. Cell. Biol.*, 2001, **3**, 346-352.
- Seely A.J., Pascual J.L., Christou N.V. Science review: Cell membrane expression (connectivity) regulates neutrophil delivery, function and clearance. *Crit. Care.*, 2003, **7**, 291-307.
- Seminario M.C., Gleich G.J. The role of eosinophils in the pathogenesis of asthma. *Curr. Opin. Immunol.*, 1994, **6**, 860-864.
- Sexton D.W., Al-Rabia M., Blaylock M.G., Walsh G.M. Phagocytosis of apoptotic eosinophils but not neutrophils by bronchial epithelial cells. *Clin. Exp. Allergy*, 2004, **34**, 1514-1524.
- Sgorbissa A., Benetti R., Marzinotto S., Schneider C., Brancolini C. Caspase-3 and caspase-7 but not caspase-6 cleave Gas2 in vitro: implications for microfilament reorganization during apoptosis. *J. Cell Sci.*, 1999, **112**, 4475-4482.
- Shapiro S.D. Matrix metalloproteinase degradation of extracellular matrix: biological consequences. *Curr. Opin. Cell Biol.*, 1998, **10**, 602-608.
- Shen Y.H., Godlewski J., Zhu J., Sathyanarayana P., Leaner V., Birrer M.J., Rana A., Tzivion G. Cross-talk between JNK/SAPK and ERK/MAPK pathways: sustained activation of JNK blocks ERK activation by mitogenic factors. *J. Biol. Chem.*, 2003, **278**, 26715-26721.
- Sher A., Coffman R.L., Hieny S., Cheever A.W. Ablation of eosinophil and IgE responses with anti-IL-5 or anti-IL-4 antibodies fails to affect immunity against *Schistosoma mansoni* in the mouse. *J. Immunol.*, 1990, **145**, 3911-3916.
- Shibasaki F., Kondo E., Akagi T., McKeon F. Suppression of signalling through transcription factor NF-AT by interactions between calcineurin and Bcl-2. *Nature*, 1997, **386**, 728-731.
- Shimabukuro M., Higa M., Zhou Y.T., Wang M.Y., Newgard C.B., Unger R.H. Lipoapoptosis in beta-cells of obese prediabetic fa/fa rats. Role of serine palmitoyltransferase overexpression. *J. Biol. Chem.*, 1998a, **273**, 32487-32490.
- Shimabukuro M., Zhou Y.T., Levi M., Unger R.H. Fatty acid-induced beta cell apoptosis: a link between obesity and diabetes. *Proc. Natl Acad. Sci. U. S. A.*, 1998b, **95**, 2498-2502.
- Shimizu S., Ide T., Yanagida T., Tsujimoto Y. Electrophysiological study of a novel large pore formed by Bax and the voltage-dependent anion channel that is permeable to cytochrome c. *J. Biol. Chem.*, 2000, **275**, 12321-12325.
- Silverstein S.C., Steinman R.M., Cohn Z.A. Endocytosis. *Annu. Rev. Biochem.*, 1977, **46**, 669-722.
- Simon H.U., Blaser K. Inhibition of programmed eosinophil death: a key pathogenic event for eosinophilia ? *Immunol. Today*, 1995, **16**, 53-55.
- Simon H.U., Yousefi S., Dibbert B., Hebestreit H., Weber M., Branch D.R., Blaser K., Levi-Schaffer F., Anderson G.P. Role for tyrosine phosphorylation and Lyn tyrosine kinase in fas receptor-mediated apoptosis in eosinophils. *Blood*, 1998, **92**, 547-557.
- Simon H.U. Neutrophil apoptosis pathways and their modifications in inflammation. *Immunol. Rev.*, 2003, **193**, 101-110.
- Singh I., Pahan K., Khan M., Singh A.K. Cytokine-mediated induction of ceramide production is redox-sensitive. Implications to proinflammatory cytokine-mediated apoptosis in demyelinating diseases. *J. Biol. Chem.*, 1998, **273**, 20354-20362.
- Slee E.A., Harte M.T., Kluck R.M., Wolf B.B., Casiano C.A., Newmeyer D.D., Wang H.G., Reed J.C., Nicholson D.W., Alnemri E.S., Green D.R., Martin S.J. Ordering the cytochrome c-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. *J. Cell Biol.*, 1999, **144**, 281-292.
- Smith C.W. Introduction: functional polarity of motile neutrophils. *Blood*, 2000, **95**, 2459-2461.

- Smolenska-Sym G., Spsychalska J., Zdebska E., Wozniak J., Traczyk Z., Pszenna E., Maj S., Danikiewicz W., Bienkowski T., Koscielak J. Ceramides and glycosphingolipids in maturation process: leukemic cells as an experimental model *Blood Cells Mol. Dis.*, 2004, **33**, 68-76.
- So T., Lee S.W., Croft M. Tumor necrosis factor/tumor necrosis factor receptor family members that positively regulate immunity. *Int. J. Hematol.*, 2006, **83**, 1-11.
- Sorrentino S., Glitz D.G., Hamann K.J., Loegering D.A., Checkel J.L., Gleich G.J. Eosinophil-derived neurotoxin and human liver ribonuclease. Identity of structure and linkage of neurotoxicity to nuclease activity. *J. Biol. Chem.*, 1992, **267**, 14859-14865.
- Squier M.K., Sehnert A.J., Sellins K.S., Malkinson A.M., Takano E., Cohen J.J. Calpain and calpastatin regulate neutrophil apoptosis. *J. Cell. Physiol.*, 1999, **178**, 311-319.
- Srinivasula S.M., Ahmad M., Fernandes-Alnemri T., Litwack G., Alnemri E.S. Molecular ordering of the Fas-apoptotic pathway: the Fas/APO-1 protease Mch5 is a CrmA-inhibitable protease that activates multiple Ced-3/ICE-like cysteine proteases. *Proc. Natl Acad. Sci. U. S. A.*, 1996, **93**, 14486-14491.
- Srinivasula S.M., Ahmad M., Fernandes-Alnemri T., Alnemri E.S. Autoactivation of procaspase-9 by Apaf-1-mediated oligomerization. *Mol. Cell.*, 1998, **1**, 949-957.
- Srinivasula S.M., Ahmad M., Guo Y., Zhan Y., Lazebnik Y., Fernandes-Alnemri T., Alnemri E.S. Identification of an endogenous dominant-negative short isoform of caspase-9 that can regulate apoptosis. *Cancer Res.*, 1999, **59**, 999-1002.
- Stanger B.Z., Leder P., Lee T.H., Kim E., Seed B. RIP: a novel protein containing a death domain that interacts with Fas/APO-1 (CD95) in yeast and causes cell death. *Cell*, 1995, **81**, 513-523.
- Steehmaier M., Borges E., Berger J., Schwarz H., Vestweber D. The E-selectin-ligand ESL-1 is located in the Golgi as well as on microvilli on the cell surface. *J. Cell. Sci.*, 1997, **110**, 687-694.
- Steinbach K.H., Schick P., Trepel F., Raffler H., Dohrmann J., Heilgeist G., Heltzel W., Li K., Past W., van der Woerd de Lange J.A., Theml H., Fliedner T.M., Begemann H. Estimation of kinetic parameters of neutrophilic, eosinophilic, and basophilic granulocytes in human blood. *Blut*, 1979, **39**, 27-38.
- Stennicke H.R., Jurgensmeier J.M., Shin H., Deveraux Q., Wolf B.B., Yang X., Zhou Q., Ellerby H.M., Ellerby L.M., Bredesen D., Green D.R., Reed J.C., Froelich C.J., Salvesen G.S. Pro-caspase-3 is a major physiologic target of caspase-8. *J. Biol. Chem.*, 1998, **273**, 27084-27090.
- Stern M., Meagher L., Savill J., Haslett C. Apoptosis in human eosinophils. Programmed cell death in the eosinophil leads to phagocytosis by macrophages and is modulated by IL-5. *J. Immunol.*, 1992, **148**, 3543-3549.
- Stern M., Savill J., Haslett C. Human monocyte-derived macrophage phagocytosis of senescent eosinophils undergoing apoptosis. Mediation by alpha v beta 3/CD36/thrombospondin recognition mechanism and lack of phlogistic response. *Am. J. Pathol.*, 1996, **149**, 911-921.
- Stie J., Jesaitis A.J. Reorganization of the human neutrophil plasma membrane is associated with functional priming: implications for neutrophil preparations. *J. Leukoc. Biol.*, 2007, **81**, 672-685.
- Strasser A., Puthalakath H., Bouillet P., Huang D.C., O'Connor L., O'Reilly L.A., Cullen L., Cory S., Adams J.M. The role of bim, a proapoptotic BH3-only member of the Bcl-2 family in cell-death control. *Ann. N. Y. Acad. Sci.*, 2000, **917**, 541-548.
- Strasser A. The role of BH3-only proteins in the immune system. *Nat. Rev. Immunol.*, 2005, **5**, 189-200.
- Strek M.E., Leff A.R. Eosinophils. In: Asthma. Barnes P.J., Grunstein M.M., Leff A.R., Woolcock A.J. (Eds), Chapter 30. Lippincott-Raven: Philadelphia, 1997.

- Strelow A., Bernardo K., Adam-Klages S., Linke T., Sandhoff K., Kronke M., Adam D. Overexpression of acid ceramidase protects from tumor necrosis factor-induced cell death. *J. Exp. Med.*, 2000, **192**, 601-612.
- Stridh H., Kimland M., Jones D.P., Orrenius S., Hampton M.B. Cytochrome c release and caspase activation in hydrogen peroxide- and tributyltin-induced apoptosis. *FEBS Lett.*, 1998, **429**, 351-355.
- Stringer R.E., Hart C.A., Edwards S.W. Sodium butyrate delays neutrophil apoptosis: role of protein biosynthesis in neutrophil survival. *Br. J. Haematol.*, 1996, **92**, 169-175.
- Stuart L.M., Ezekowitz R.A. Phagocytosis: elegant complexity. *Immunity*, 2005, **22**, 539-550.
- Susin S.A., Zamzami N., Castedo M., Hirsch T., Marchetti P., Macho A., Daugas E., Geuskens M., Kroemer G. Bcl-2 inhibits the mitochondrial release of an apoptogenic protease. *J. Exp. Med.*, 1996, **184**, 1331-1341.
- Susin S.A., Lorenzo H.K., Zamzami N., Marzo I., Brenner C., Larochette N., Prevost M.C., Alzari P.M., Kroemer G. Mitochondrial release of caspase-2 and -9 during the apoptotic process. *J. Exp. Med.*, 1999a, **189**, 381-394.
- Susin S.A., Lorenzo H.K., Zamzami N., Marzo I., Snow B.E., Brothers G.M., Mangion J., Jacotot E., Costantini P., Loeffler M., Larochette N., Goodlett D.R., Aebersold R., Siderovski D.P., Penninger J.M., Kroemer G. Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature*, 1999b, **397**, 441-446.
- Suzuki M., Youle R.J., Tjandra N. Structure of Bax: coregulation of dimer formation and intracellular localization. *Cell*, 2000, **103**, 645-654.
- Suzuki Y., Imai Y., Nakayama H., Takahashi K., Takio K., Takahashi R. A serine protease, HtrA2, is released from the mitochondria and interacts with XIAP, inducing cell death. *Mol. Cell.*, 2001, **8**, 613-621.
- Swanson J.A., Hoppe A.D. The coordination of signaling during Fc receptor-mediated phagocytosis. *J. Leukoc. Biol.*, 2004, **76**, 1093-1103.

T

- Taha T.A., Mullen T.D., Obeid L.M. A house divided: ceramide, sphingosine, and sphingosine-1-phosphate in programmed cell death. *Biochim. Biophys. Acta*, 2006, **1758**, 2027-2036.
- Tai P.C., Spry C.J., Peterson C., Venge P., Olsson I. Monoclonal antibodies distinguish between storage and secreted forms of eosinophil cationic protein. *Nature*, 1984, **309**, 182-184.
- Tai P.C., Sun L., Spry C.J. Effects of IL-5, granulocyte/macrophage colony-stimulating factor (GM-CSF) and IL-3 on the survival of human blood eosinophils *in vitro*. *Clin. Exp. Immunol.*, 1991, **85**, 312-316.
- Taimen P., Kallajoki M. NuMA and nuclear lamins behave differently in Fas-mediated apoptosis. *J. Cell Sci.*, 2003, **116**, 571-583.
- Takahashi H., Nukiwa T., Yoshimura K., Quick C.D., States D.J., Holmes M.D., Whang-Peng J., Knutsen T., Crystal R.G. Structure of the human neutrophil elastase gene. *J. Biol. Chem.*, 1988, **263**, 14739-14747.
- Takahashi M., Mukai H., Toshimori M., Miyamoto M., Ono Y. Proteolytic activation of PKN by caspase-3 or related protease during apoptosis. *Proc. Natl. Acad. Sci. U. S. A.*, 1998, **95**, 11566-11571.
- Takeda Y., Watanabe H., Yonehara S., Yamashita T., Saito S., Sendo F. Rapid acceleration of neutrophil apoptosis by tumor necrosis factor-alpha. *Int. Immunol.*, 1993, **5**, 691-694.
- Tamiya-Koizumi K., Umekawa H., Yoshida S., Kojima K. Existence of Mg²⁺-dependent, neutral sphingomyelinase in nuclei of rat ascites hepatoma cells. *J. Biochem. (Tokyo)*, 1989, **106**, 593-598.
- Tammariello S.P., Quinn M.T., Estus S. NADPH oxidase contributes directly to oxidative stress and apoptosis in nerve growth factor-deprived sympathetic neurons. *J. Neurosci.*, 2000, **20**, RC53.

- Taverne J., Bradley J.E. Immunity to Protozoa and Worms. In: Roit I., Brostoff J., Male D. (Eds.), *Immunology*. Chapter 18. Mosby: London, 1998.
- Teixeira M.M., Williams T.J., Hellewell P.G. Mechanisms and pharmacological manipulation of eosinophil accumulation *in vivo*. *Trends Pharmacol. Sci.*, 1995, **16**, 418-423.
- Temkin V., Levi-Schaffer F. Mechanism of tumour necrosis factor alpha mediated eosinophil survival. *Cytokine*, 2001, **15**, 20-26.
- Tepper A.D., Diks S.H., van Blitterswijk W.J., Borst J. Glucosylceramide synthase does not attenuate the ceramide pool accumulating during apoptosis induced by CD95 or anti-cancer regimens. *J. Biol. Chem.*, 2000, **275**, 34810-34817.
- Thome M., Schneider P., Hofmann K., Fickenscher H., Meinel E., Neipel F., Mattmann C., Burns K., Bodmer J.L., Schroter M., Scaffidi C., Krammer P.H., Peter M.E., Tschopp J. Viral FLICE-inhibitory proteins (FLIPs) prevent apoptosis induced by death receptors. *Nature*, 1997, **386**, 517-521.
- Torrego A., Hew M., Oates T., Sukkar M., Fan Chung K. Expression and activation of TGF-beta isoforms in acute allergen-induced remodelling in asthma. *Thorax*, 2007, **62**, 307-313.
- Tsujimoto Y., Finger L.R., Yunis J., Nowell P.C., Croce C.M. Cloning of the chromosome breakpoint of neoplastic B cells with the t(14;18) chromosome translocation. *Science*, 1984, **226**, 1097-1099.
- Tsukahara K., Nakao A., Hiraguri M., Miike S., Mamura M., Saito Y., Iwamoto I. Tumor necrosis factor-alpha mediates antiapoptotic signals partially via p38 MAPkinase activation in human eosinophils. *Int. Arch. Allergy Immunol.*, 1999, **120**, 54-59.
- Tsuyuki S., Bertrand C., Erard F., Trifilieff A., Tsuyuki J., Wesp M., Anderson G.P., Coyle A.J. Activation of the Fas receptor on lung eosinophils leads to apoptosis and the resolution of eosinophilic inflammation of the airways. *J. Clin. Invest.*, 1995, **96**, 2924-2931.
- Turlej R.K., Fiévez L., Sandersen C.F., Dogné S., Kirschvink N., Lekeux P., Bureau F. Enhanced survival of lung granulocytes in an animal model of asthma : evidence for a role of GM-CSF-activated STAT5 signaling pathway. *Thorax*, 2001, **56**, 696-702.
- Turner J., Cho Y., Dinh N.N., Waring A.J., Lehrer R.I. Activities of LL-37, a cathelin-associated antimicrobial peptide of human neutrophils. *Antimicrob. Agents Chemother.*, 1998, **42**, 2206-2214.
- Turzanski J., Grundy M., Shang S., Russell N., Pallis M. P-glycoprotein is implicated in the inhibition of ceramide-induced apoptosis in TF-1 acute myeloid leukemia cells by modulation of the glucosylceramide synthase pathway. *Exp. Hematol.*, 2005, **33**, 62-72.

U

- Ucker D.S., Obermiller P.S., Eckhart W., Apgar J.R., Berger N.A., Meyers J. Genome digestion is a dispensable consequence of physiological cell death mediated by cytotoxic T lymphocytes. *Mol. Cell Biol.*, 1992, **12**, 3060-3069.
- Ullman M.D., Radin N.S. The enzymatic formation of sphingomyelin from ceramide and lecithin in mouse liver. *J. Biol. Chem.*, 1974, **249**, 1506-1512.

V

- Van den Berg J.M., Weyer S., Weening J.J., Roos D., Kuijpers T.W. Divergent effects of tumor necrosis factor- α on apoptosis of human neutrophils. *J. Leukoc. Biol.*, 2001, **69**, 467-473.

- van der Bruggen T., Caldenhoven E., Kanters D., Coffey P., Raaijmakers J.A., Lammers J.W., Koenderman L. Interleukin-5 signaling in human eosinophils involves JAK2 tyrosine kinase and Stat1 alpha. *Blood*, 1995, **85**, 1442-1448.
- van der Bruggen T., Koenderman L. Signal transduction in eosinophils. *Clin. Exp. Allergy*, 1996, **26**, 880-891.
- van Gisbergen K.P., Sanchez-Hernandez M., Geijtenbeek T.B., van Kooyk Y. Neutrophils mediate immune modulation of dendritic cells through glycosylation-dependent interactions between Mac-1 and DC-SIGN. *J. Exp. Med.*, 2005, **201**, 1281-1292.
- Vander Heiden M.G., Chandel N.S., Schumacker P.T., Thompson C.B. Bcl-xL prevents cell death following growth factor withdrawal by facilitating mitochondrial ATP/ADP exchange. *Mol. Cell.*, 1999, **3**, 159-167.
- Vassina E.M., Yousefi S., Simon D., Zwicky C., Conus S., Simon H.U. cIAP-2 and survivin contribute to cytokine-mediated delayed eosinophils apoptosis. *Eur. J. Immunol.*, 2006, **36**, 1975-1984.
- Vaux D.L. Apoptosis timeline. *Cell Death Differ.*, 2002, **9**, 349-354.
- Venge P. Human eosinophil granule proteins: structure, function and release. In: Immunopharmacology of Eosinophils. Smith H., Cook R.M. (Eds.), Academic Press, London, 1993, 43-55.
- Verhagen A.M., Coulson E.J., Vaux D.L. Inhibitor of apoptosis proteins and their relatives: IAPs and other BIRPs. *Genome Biol.*, 2001, **2**, S3009-S3016.
- Verheij M., Bose R., Lin X.H., Yao B., Jarvis W.D., Grant S., Birrer M.J., Szabo E., Zon L.I., Kyriakis J.M., Haimovitz-Friedman A., Fuks Z., Kolesnick R.N. Requirement for ceramide-initiated SAPK/JNK signalling in stress-induced apoptosis. *Nature*, 1996, **380**, 75-79.
- Viegi G., Annesi I., Matteelli G. Epidemiology of asthma. In: European Respiratory Monograph, Volume 8, Monograph 23 : Asthma, Chung F, Fabbri LM (Eds), Sheffield, UK : European Respiratory Society Journals Ltd, 2003, 1-25.
- Vignola A.M., Chanez P., Chiappara G., Siena L., Merendino A., Reina C., Gagliardo R., Profita M., Bousquet J., Bonsignore G. Evaluation of apoptosis of eosinophils, macrophages, and T lymphocytes in mucosal biopsy specimens of patients with asthma and chronic bronchitis. *J. Allergy Clin. Immunol.*, 1999, **103**, 563-573.
- Vijayanand P., Seumois G., Pickard C., Powell R.M., Angco G., Sammut D., Gadola S.D., Friedmann P.S., Djukanovic R. Invariant natural killer T cells in asthma and chronic obstructive pulmonary disease. *N. Engl. J. Med.*, 2007, **356**, 1410-22.
- Vogt C. Untersuchungen über die Entwicklungsgeschichte der Geburtshelerkroete (*Alytes obstetricians*), Solothurn: Jent und Gassman, 1842, pp 130.

W

- Walker C., Virchow J.C. Jr, Bruijnzeel P.L., Blaser K. T cells and asthma. II. Regulation of the eosinophilia of asthma by T cell cytokines. *Int. Arch. Allergy Appl. Immunol.*, 1991, **94**, 248-250.
- Walker R.I., Willemze R. Neutrophil kinetics and the regulation of granulopoiesis. *Rev. Infect. Dis.*, 1980, **2**, 282-292.
- Wang H.G., Miyashita T., Takayama S., Sato T., Torigoe T., Krajewski S., Tanaka S., Hovey L. 3rd, Troppmair J., Rapp U.R., et al. Apoptosis regulation by interaction of Bcl-2 protein and Raf-1 kinase. *Oncogene*, 1994, **9**, 2751-2756.
- Wang H.G., Pathan N., Ethell I.M., Krajewski S., Yamaguchi Y., Shibasaki F., McKeon F., Bobo T., Franke T.F., Reed J.C. Ca²⁺-induced apoptosis through calcineurin dephosphorylation of BAD. *Science*, 1999, **284**, 339-343.

- Wang K.K., Posmantur R., Nadimpalli R., Nath R., Mohan P., Nixon R.A., Talanian R.V., Keegan M., Herzog L., Allen H. Caspase-mediated fragmentation of calpain inhibitor protein calpastatin during apoptosis. *Arch. Biochem. Biophys.*, 1998, **356**, 187-196.
- Ward C., Chilvers E.R., Lawson M.F., Pryde J.G., Fujihara S., Farrow S.N., Haslett C., Rossi A.G. NF-kappaB activation is a critical regulator of human granulocyte apoptosis in vitro. *J. Biol. Chem.*, 1999, **274**, 4309-4318.
- Wardlaw A.J., Moqbel R., Kay A.B. Eosinophils: biology and role in disease. *Adv. Immunol.*, 1995, **60**, 151-266.
- Wasmoen T.L., Bell M.P., Loegering D.A., Gleich G.J., Prendergast F.G., McKean D.J. Biochemical and amino acid sequence analysis of human eosinophil granule major basic protein. *J. Biol. Chem.*, 1988, **263**, 12559-12563.
- Watanabe M., Nittoh T., Suzuki T., Kitoh A., Mue S., Ohuchi K. Isolation and partial characterization of eosinophil granule proteins in rats—Eosinophil cationic protein and major basic protein. *Int. Arch. Allergy Immunol.*, 1995, **108**, 11-18.
- Watson R.W., Redmond H.P., Wang J.H., Bouchier-Hayes D. Bacterial ingestion, tumor necrosis factor-alpha, and heat induce programmed cell death in activated neutrophils. *Shock*, 1996, **5**, 47-51.
- Watts J.D., Gu M., Polverino A.J., Patterson S.D., Aebersold R. Fas-induced apoptosis of T cells occurs independently of ceramide generation. *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 7292-7296.
- Webb S.J., Harrison D.J., Wyllie A.H. Apoptosis: an overview of the process and its relevance in disease. *Adv. Pharmacol.*, 1997, **41**, 1-34.
- Weersink A.J., van Kessel K.P., van den Tol M.E., van Strijp J.A., Torensma R., Verhoef J., Elsbach P., Weiss J. Human granulocytes express a 55-kDa lipopolysaccharide-binding protein on the cell surface that is identical to the bactericidal/permeability-increasing protein. *J. Immunol.*, 1993, **150**, 253-263.
- Wei S., Liu J.H., Epling-Burnette P.K., Gamero A.M., Ussery D., Pearson E.W., Elkabani M.E., Diaz J.I., Djeu J.Y. Critical role of Lyn kinase in inhibition of neutrophil apoptosis by granulocyte-macrophage colony-stimulating factor. *J. Immunol.*, 1996, **157**, 5155-5162.
- Weinmann P., Gaetgens P., Walzog B. Bcl-xL and Bax- alpha-mediated regulation of apoptosis of human neutrophils via caspase-3. *Blood*, 1999, **93**, 3106-3115.
- Weinrauch Y., Elsbach P., Madsen L.M., Foreman A., Weiss J. The potent anti-Staphylococcus aureus activity of a sterile rabbit inflammatory fluid is due to a 14-kD phospholipase A2. *J. Clin. Invest.*, 1996, **97**, 250-257.
- Weiss S.J., Peppin G., Ortiz X., Ragsdale C., Test S.T. Oxidative autoactivation of latent collagenase by human neutrophils. *Science*, 1985, **227**, 747-749.
- Weiss S.J., Peppin G.J. Collagenolytic metalloenzymes of the human neutrophil. Characteristics, regulation and potential function *in vivo*. *Biochem. Pharmacol.*, 1986, **35**, 3189-3197.
- Weissmann G. Inflammation: historical perspective. In: Inflammation, Gallin J.I., Goldstein I.M., Snyderman R. (Eds), New York : Raven Press, 1988, pp 5-8.
- Weller P.F., Bach D., Austen K.F. Human eosinophil lysophospholipase: The sole protein component of Charcot-Leyden crystals. *J. Immunol.*, 1982, **128**, 1346-1349.
- Weller P.F., Bach D.S., Austen K.F. Biochemical characterization of human eosinophil Charcot-Leyden crystal protein (lysophospholipase). *J. Biol. Chem.*, 1984, **259**, 15100-15105.
- Westwick J.K., Bielawska A.E., Dbaibo G., Hannun Y.A., Brenner D.A. Ceramide activates the stress-activated protein kinases. *J. Biol. Chem.*, 1995, **270**, 22689-22692.
- Widlak P., Li L.Y., Wang X., Garrard W.T. Action of recombinant human apoptotic endonuclease G on naked DNA and chromatin substrates: cooperation with exonuclease and DNase I. *J. Biol. Chem.*, 2001, **276**, 48404-48409.

- Widmann C., Gerwins P., Johnson N.L., Jarpe M.B., Johnson G.L. MEK kinase 1, a substrate for DEVD-directed caspases, is involved in genotoxin-induced apoptosis. *Mol. Cell. Biol.*, 1998, **18**, 2416-2429.
- Wieder T., Orfanos C.E., Geilen C.C. Induction of ceramidemediated apoptosis by the anticancer phospholipid analog, hexadecylphosphocholine. *J. Biol. Chem.*, 1998, **273**, 11025–11031.
- Wiegmann K., Schutze S., Machleidt T., Witte D., Kronke M. Functional dichotomy of neutral and acidic sphingomyelinases in tumor necrosis factor signaling. *Cell*, 1994, **78**, 1005-1015.
- Wiegmann K., Schwandner R., Krut O., Yeh W.C., Mak T.W., Kronke M. Requirement of FADD for tumor necrosis factor-induced activation of acid sphingomyelinase. *J. Biol. Chem.*, 1999, **274**, 5267-5270.
- Williams M.A., Solomkin J.S. Integrin-mediated signaling in human neutrophil functioning. *J. Leukoc. Biol.*, 1999, **65**, 725-736.
- Woerly G., Roger N., Loiseau S., Dombrowicz D., Capron A., Capron M. Expression of CD28 and CD86 by human eosinophils and role in the secretion of type 1 cytokines (interleukin 2 and interferon gamma): inhibition by immunoglobulin A complexes. *J. Exp. Med.*, 1999, **190**, 487-495.
- Wolter K.G., Hsu Y.T., Smith C.L., Nechushtan A., Xi X.G., Youle R.J. Movement of Bax from the cytosol to mitochondria during apoptosis. *J. Cell. Biol.*, 1997, **139**, 1281-1292.
- Wong C.K., Zhang J., Ip W.K., Lam C.W. Intracellular signal transduction in eosinophils and its clinical significance. *Immunopharmacol. Immunotoxicol.*, 2002, **24**, 165-186.
- Woolley K.L., Gibson P.G., Carty K., Wilson A.J., Twaddell S.H., Woolley M.J. Eosinophil apoptosis and the resolution of airway inflammation in asthma. *Am. J. Respir. Crit. Care Med.*, 1996, **154**, 237-243.
- Worthen G.S., Avdi N., Vukajlovich S., Tobias P.S. Neutrophil adherence induced by lipopolysaccharide *in vitro*. Role of plasma component interaction with lipopolysaccharide. *J. Clin. Invest.*, 1992, **90**, 2526-2535.
- Wu Y.C., Stanfield G.M., Horvitz H.R. NUC-1, a caenorhabditis elegans DNase II homolog, functions in an intermediate step of DNA degradation during apoptosis. *Genes Dev.*, 2000, **14**, 536-548.
- Wyllie A.H. Glucocorticoid-induced thymocyte apoptosis is associated with endogenous endonuclease activation. *Nature*, 1980, **284**, 555-556.

X

- Xia Z., Dickens M., Raingeaud J., Davis R.J., Greenberg M.E. Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science*, 1995, **270**, 1326-1331.
- Xu J., Yeh C.H., Chen S., He L., Sensi S.L., Canzoniero L.M., Choi D.W., Hsu C.Y. Involvement of de novo ceramide biosynthesis in tumor necrosis factor- α /cycloheximide-induced cerebral endothelial cell death. *J. Biol. Chem.*, 1998, **273**, 16521–16526.

Y

- Yamaguchi K., Nishio K., Sato N., Tsumura H., Ichihara A., Kudo H., Aoki T., Naoki K., Suzuki K., Miyata A., Suzuki Y., Morooka S. Leukocyte kinetics in the pulmonary microcirculation: observations using real-time confocal luminescence microscopy coupled with high-speed video analysis. *Lab. Invest.*, 1997, **76**, 809-822.

- Yamashita K., Takahashi A., Kobayashi S., Hirata H., Mesner P.W. Jr., Kaufmann S.H., Yonehara S., Yamamoto K., Uchiyama T., Sasada M. Caspases mediate tumor necrosis factor-alpha-induced neutrophil apoptosis and downregulation of reactive oxygen production. *Blood*, 1999, **93**, 674-685.
- Yan L., Vandivier R.W., Suffredini A.F., Danner R.L. Human polymorphonuclear leukocytes lack detectable nitric oxide synthase activity. *J. Immunol.*, 1994, **153**, 1825-1834.
- Ye Y.L., Huang W.C., Lee Y.L., Chiang B.L. Interleukin-12 inhibits eotaxin secretion of cultured primary lung cells and alleviates airway inflammation in vivo. *Cytokine*, 2002, **19**, 76-84.
- Ye H., Cande C., Stephanou N.C., Jiang S., Gurbuxani S., Larochette N., Daugas E., Garrido C., Kroemer G., Wu H. DNA binding is required for the apoptogenic action of apoptosis inducing factor. *Nat. Struct. Biol.*, 2002, **9**, 680-684.
- Yoder M.C., Checkley L.L., Giger U., Hanson W.L., Kirk K.R., Capen R.L., Wagner W.W. Jr. Pulmonary microcirculatory kinetics of neutrophils deficient in leukocyte adhesion-promoting glycoproteins. *J. Appl. Physiol.*, 1990, **69**, 207-213.
- Yousefi S., Green D.R., Blaser K., Simon H.U. Protein-tyrosine phosphorylation regulates apoptosis in human eosinophils and neutrophils. *Proc. Natl. Acad. Sci. U. S. A.*, 1994, **91**, 10868-10872.
- Yousefi S., Blaser K., Simon H.U. Activation of signaling pathways and prevention of apoptosis by cytokines in eosinophils. *Int. Arch. Allergy Immunol.*, 1997, **112**, 9-12.

Z

- Zanetti M., Gennaro R., Romeo D. The cathelicidin family of antimicrobial peptide precursors: a component of the oxygen-independent defense mechanisms of neutrophils. *Ann. N. Y. Acad. Sci.*, 1997, **832**, 147-162.
- Zangrilli J., Robertson N., Shetty A., Wu J., Hastie A., Fish J.E., Litwack G., Peters S.P. Effect of IL-5, glucocorticoid, and Fas ligation on Bcl-2 homologue expression and caspase activation in circulating human eosinophils. *Clin. Exp. Immunol.*, 2000, **120**, 12-21.
- Zeibecoglou K., Ying S., Meng Q., Kay A.B., Robinson D.S., Papageorgiou N. Expression of eotaxin in induced sputum of atopic and nonatopic asthmatics. *Allergy*, 2000, **55**, 1042-1048.
- Zha J., Harada H., Yang E., Jockel J., Korsmeyer S.J. Serine phosphorylation of death agonist BAD in response to survival factor results in binding to 14-3-3 not BCL-X(L). *Cell*, 1996, **87**, 619-628.
- Zhang X., Moilanen E., Kankaanranta H. Enhancement of human eosinophil apoptosis by fluticasone propionate, budesonide, and beclomethasone. *Eur. J. Pharmacol.*, 2000, **406**, 325-332.
- Zhang X., Moilanen E., Lahti A., Hamalainen M., Giembycz M.A., Barnes P.J., Lindsay M.A., Kankaanranta H. Regulation of eosinophil apoptosis by nitric oxide: Role of c-Jun-N-terminal kinase and signal transducer and activator of transcription 5. *J. Allergy Clin. Immunol.*, 2003, **112**, 93-101.
- Zhang Y., Center D.M., Wu D.M., Cruikshank W.W., Yuan J., Andrews D.W., Kornfeld H. Processing and activation of pro-interleukin-16 by caspase-3. *J. Biol. Chem.*, 1998, **273**, 1144-1149.
- Zhu D., Hattori H., Jo H., Jia Y., Subramanian K.K., Loison F., You J., Le Y., Honczarenko M., Silberstein L., Luo H.R. Deactivation of phosphatidylinositol 3,4,5-trisphosphate/Akt signaling mediates neutrophil spontaneous death. *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 14836-14841.
- Zhu L., Ling S., Yu X.D., Venkatesh L.K., Subramanian T., Chinnadurai G., Kuo T.H. Modulation of mitochondrial Ca(2+) homeostasis by Bcl-2. *J. Biol. Chem.*, 1999, **274**, 33267-33273.

- Zou H., Henzel W.J., Liu X., Lutschg A., Wang X. Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in cytochrome c-dependent activation of caspase-3. *Cell*, 1997, **90**, 405-413.
- Zou H., Li Y., Liu X., Wang X. An APAF-1.cytochrome c multimeric complex is a functional apoptosome that activates procaspase-9. *J. Biol. Chem.*, 1999, **274**, 11549-11556.
- Zundel W., Giaccia A. Inhibition of the anti-apoptotic PI(3)K/Akt/Bad pathway by stress. *Genes Dev.*, 1998, **12**, 1941-1946.
- Zundel W., Swiersz L.M., Giaccia A. Caveolin 1-mediated regulation of receptor tyrosine kinase-associated phosphatidylinositol 3-kinase activity by Ceramide. *Mol. Cell. Biol.*, 2000, **20**, 1507-1514.