

## Motivations

Anticoherent states are maximally entangled symmetric multiqubit states, optimal for rotation and magnetic field sensing.<sup>1</sup> Despite growing interest<sup>2</sup>, a general experimental protocol for their generation has remained elusive, until now.

## Anticoherent states

An anticoherent state to order  $t$  is an isotropic state for which the moments of  $\mathbf{J} \cdot \mathbf{n}$  up to order  $t$  are independent of the direction  $\mathbf{n}$ , i.e.

$$\text{Tr}[\rho(\mathbf{J} \cdot \mathbf{n})^k] \neq f(\mathbf{n}) \text{ for } k = 1, 2, \dots, t$$

We can quantify the degree of anticoherence of any state  $|\psi\rangle$  through the measure of anticoherence<sup>3</sup>

$$\mathcal{A}_t^{\text{Bures}} = 1 - \sqrt{\frac{\sqrt{t+1} - \sum_{i=1}^{t+1} \sqrt{\lambda_i}}{\sqrt{t+1} - 1}} \quad \text{5-anticoherent state}$$



where the  $\lambda_i$  are the Schmidt coefficients of  $|\psi\rangle$ . This measure can serve as a figure of merit for numerical optimisation.

Any symmetric multiqubit state can be expanded in the tensor multipolar basis

$$\rho = \sum_{L=0}^N \sum_{M=-L}^L \rho_{LM} T_{LM}.$$

In this manner, a state is anticoherent of order  $t$  iff

$$\rho_{LM} = 0 \text{ for } L = 1, 2, \dots, t \text{ and } M = -L, \dots, L$$

Hence, in order to generate anticoherent states, the goal is to completely depopulate the lower state multipoles  $\rho_{LM}$ .

## Pulse-based protocol


Our protocol consists of a sequence of  $n_C$  cycles in which we apply first a rotation followed by a squeezing<sup>4</sup>

$$\Pi_y(\theta) = e^{-iJ_y\theta} \quad \Pi_{z2}(\eta) = e^{-iJ_z^2\eta}$$

such that the final state is

$$|\psi_{n_C}\rangle = \left( \prod_{i=1}^{n_C} \Pi_{z2}(\eta_i) \Pi_y(\theta_i) \right) |\psi_0\rangle$$

where  $\theta_1 = 0$  and the initial state is  $|\psi_0\rangle = e^{-iJ_x\pi/2}|0\rangle^{\otimes N}$  where  $|0\rangle$  is the qubit ground state.




From a physical point of view, our protocol exploits the fact that the rotations (resp. squeezings) only couple multipoles with the same  $L$  (resp.  $M$ ). Moreover, during a squeezing along  $z$ , the transitions for  $M = 0$  are forbidden.

For different number of qubits  $N$  and anticoherence order  $t$ , we use the Nelder-Mead algorithm to find the  $2n_C$  parameters  $\theta_i$  and  $\eta_i$  that maximises  $\mathcal{A}_t^{\text{Bures}}$ .


## Numerical results

For all  $N \leq 48$ , the optimised parameters are able to generate the highest order anticoherent state allowed.

(State multipoles evolution for  $N = 6$  and  $t = 3$ )



(Achieved anticoherence for a fixed number of cycles  $n_C$ )




For each anticoherence order ( $t = 2, 3$  and  $4$ ), there is a threshold number of cycles above which the state is perfectly anticoherent (up to numerical errors).

For  $N = 48$ , we obtain a 9-anticoherent state, the maximal order for this  $N$ , verifying  $\mathcal{A}_9^{\text{Bures}} > 0.99$  with  $n_C = 14$ .

## High-dimensional systems

We observe that, for even  $N$ , in order to generate an anticoherent state of order 2, we can take  $\eta_1 = \frac{\pi}{2}$ ,  $\theta_2 = -\frac{\pi}{2N}$  and  $\theta_3 = \frac{\pi}{2}$  and only optimise over  $\eta_2$  and  $\eta_3$ .



It allows us to obtain the scaling laws  $\eta_2 = \frac{3}{4\sqrt{N}}$  and  $\eta_3 = \frac{5}{2\sqrt{N}}$  that we can exploit to generate anticoherent states with  $N = 10000$  verifying  $1 - \mathcal{A}_2^{\text{Bures}} < 10^{-3}$ .

## Perspectives

- Test the protocol in experiments with spin or Bose-Einstein condensate systems.
- Use anticoherent states for improved gyroscopy or magnetometry applications in the lab.

## References

1. J. Martin, S. Weigert and O. Giraud, *Quantum* **4**, 285 (2022)
2. M. Rudzinski, A. Burchardt and K. Zyczkowski, *Quantum* **8**, 1234 (2024)
3. D. Baguette and J. Martin, *Phys. Rev. A* **96**, 032304 (2017)
4. S. C. Carrasco *et al.*, *Phys. Rev. Applied* **17**, 064050 (2022)