



Preservation modes of exceptionally preserved fossils from the Early Ordovician Cabrières Biota, France

FARID SALEH, PIERRE GUERIAU, CRISTINA MARTIN-OLMOS, ROMAIN GOUGEON, ROMAIN VAUCHER, PRESCOTT VAYDA, VALENTIN JAMART, NORA CORTHÉSY, SINÉAD LYNCH, GAËTAN J.-M. POTIN, HARRIET B. DRAGE, DAMIEN PAS, ALEXANDRE TORCHET, JONATHAN POPE, MERLE GREIF, ENZO BIROLINI, CHRISTOPHE DUPICHAUD, SOLINE MICHEL, PAULINE GUENSER, ERIC MONCERET AND SYLVIE MONCERET

LETHAIA



The Cabrières Biota is a recently discovered Early Ordovician fossil assemblage from southern France, comprising a diverse range of organisms and recording a palaeopolar ecosystem. Despite its ecological significance and due to its recent discovery, research on this biota is still in its early stages, and the specific taphonomic pathway behind its preservation remains poorly understood. In this study, we use scanning electron microscopy to investigate the taphonomy of soft-bodied fossils from the Cabrières Biota. Our results show that soft-bodied fossils were predominantly pyritized under sulphate-reducing conditions during early diagenesis. This was accompanied by the dissolution of biogenic silicon in skeletal elements of organisms such as sponges, with the released silica likely contributing to the partial silicification of other soft tissues. In fossil soft parts where little to no pyrite or silica enrichment is present, carbon is observed. All exceptionally preserved fossils exhibit signs of significant post-early diagenetic weathering, which altered the taphonomic modes. Pyrite crystals have undergone dissolution and reprecipitation into poorly crystalline iron oxide phases, while carbon has often been partly leached away. Thus, the Cabrières Biota represents a rare case in which a highly reactive early diagenetic environment, coupled with a complex history of maturation and surface weathering, has produced a mosaic of preservation modes even within individual fossils.

□ Floian, exceptional preservation, taphonomy, Palaeozoic.

Farid Saleh ✉ [farid.nassim.saleh@gmail.com], Pierre Gueriau, Valentin Jamart, Nora Corthésy, Sinéad Lynch, Gaëtan J.-M. Potin, Harriet B. Drage and Alexandre Torchet, Institute of Earth Sciences, University of Lausanne, Lausanne, Switzerland; Damien Pas, Institute of Earth Sciences, University of Lausanne, Lausanne, Switzerland, and SediCClim, Geology Department, Liège University, Liège, Belgium; Cristina Martin-Olmos, Center for Advanced Surface Analysis, Institute of Earth Sciences, University of Lausanne, Switzerland, and School of Architecture, Civil and Environmental Engineering, EPFL, Lausanne, Switzerland; Romain Gougeon, Geo-Ocean, University of Brest, CNRS, Ifremer, UMR 6538, F-29280 Plouzané, France; Romain Vaucher, College of Science and Engineering, James Cook University, Townsville, QLD, 4811, Australia; Prescott Vayda, Department of Geosciences, Virginia Tech, Blacksburg, Virginia 24061, USA; Jonathan Pople and Merle Greif, Department of Palaeontology, University of Zurich, Karl-Schmid-Strasse 4, Zurich 8006, Switzerland; Enzo Birolini, Christophe Dupichaud, Soline Michel, and Pauline Guenser, Université de Lyon, Université Claude Bernard Lyon 1, École Normale Supérieure de Lyon, CNRS, UMR5276, LGL-TPE, Villeurbanne, France; Eric Monceret and Sylvie Monceret, Société d'Études Scientifiques de l'Aude, Carcassonne, France; manuscript received on 29/08/2025; manuscript accepted on 29/11/2025; manuscript published on 19/02/2026 in *Lethaia* 59(2).

The Ordovician is widely known for experiencing two major events that profoundly shaped life on Earth: the Ordovician Radiation(s) or Great Ordovician Biodiversification Event, during which most animal phyla underwent a more or less rapid diversification; and the Late Ordovician Glaciation, which caused a sharp decline in biodiversity over a relatively short timespan (Harper *et al.* 2014; Servais & Harper 2018; Rasmussen *et al.* 2019; Stigall *et al.* 2019; Servais *et al.* 2021; Saleh *et al.* 2023a). For decades, our understanding of these diversification and extinction events has primarily relied on the fossil record

of biomineralized, or 'shelly', organisms, such as brachiopods, which preserve well throughout geological timescales (e.g. Harper *et al.* 2015; Stigall *et al.* 2023).

However, shelly organisms represent only a fraction of ancient ecosystems. Many animals are either partially or entirely soft-bodied, having decay-prone tissues that do not fossilize easily (e.g. Parry *et al.* 2018; Purnell *et al.* 2018). Studying these soft-bodied organisms is crucial for capturing a more complete picture of ancient ecosystems and for accurately assessing biodiversity dynamics through deep time (Seilacher *et al.* 1985; Kimmig & Schiffbauer 2024). Soft tissues

are only preserved in sites with exceptional fossil preservation, known as *Konservat-Lagerstätten* (Seilacher *et al.* 1985; Kimmig & Schiffbauer 2024). These sites provide invaluable glimpses into past ecosystems.

One such site for the Ordovician is the Fezouata Shale (Tremadocian–Floian, Morocco; Van Roy *et al.* 2010; Martin *et al.* 2016; Saleh *et al.* 2021a). This Lagerstätte has yielded critical palaeontological (e.g. Vinther *et al.* 2008, 2017; Polechová 2016; Van Iten *et al.* 2016; Gutiérrez-Marco *et al.* 2017; Saleh *et al.* 2018, 2020a, 2021b, 2022a; Lefebvre *et al.* 2019; Pérez-Peris *et al.* 2021; Drage *et al.* 2023; Dupichaud *et al.* 2023; Laibl *et al.* 2023a, b; Potin *et al.* 2023; Lustri *et al.* 2024), sedimentological (e.g. Vaucher *et al.* 2016, 2017; Saleh *et al.* 2019, 2020b, 2022b), and taphonomic (Saleh *et al.* 2020c, 2021c, 2023b, 2024a) insights that have reshaped our understanding of the Cambrian–Ordovician transition. Fossils from Fezouata revealed that many animal groups considered extinct by the end of the Cambrian survived well into the Ordovician (e.g. Van Roy *et al.* 2010; Laibl *et al.* 2023a). This discovery suggested that the Cambrian Explosion and the Ordovician Radiation(s) rather represent a continuous interval of diversification (Van Roy *et al.* 2010, Saleh *et al.* 2020c). The discovery of the Fezouata Biota underscored the significance of any new Early Ordovician site preserving soft-bodied organisms, as such discoveries have the potential to fundamentally advance our understanding of early animal evolution.

The recently discovered Cabrières Biota from southern France is an exceptionally preserved assemblage that includes a diverse array of organisms, such as algae, sponges, cnidarians, arthropods, vermiform organisms, molluscs, brachiopods, echinoderms, and hemichordates (Saleh *et al.* 2024b, c). Remarkably, this site lies closer to the palaeo-South Pole than any other known Ordovician *Konservat-Lagerstätte* (Saleh *et al.* 2024b). Its high fossil diversity amid elevated Early Ordovician sea-surface temperatures suggests that high-latitude regions may have served as refugia for marine life escaping more extreme equatorial conditions (Saleh *et al.* 2024b).

Despite its significance, research on the Cabrières Biota is still in its infancy owing to its recent discovery (Saleh *et al.* 2024b, c), and little is known about the taphonomy of this unique assemblage. Since taphonomic investigations are key for subsequent palaeontological interpretations, here we aim to comprehensively investigate the mode of preservation of soft-bodied organisms from the Cabrières Biota. This is complemented by other studies in this special volume investigating sedimentary processes,

ichnological evidence, and palaeontological occurrences in this *Konservat-Lagerstätte* (Caillaud *et al.* this volume; Daley *et al.* in this volume; Gougeon *et al.* this volume; Gutiérrez-Marco *et al.* in this volume; Harper *et al.* this volume; Lefebvre *et al.* this volume; Li *et al.* this volume; Monceret *et al.* this volume; Polechová *et al.* this volume; Van Iten *et al.* this volume; Vannier *et al.* this volume; Vaucher *et al.* this volume; Vayda *et al.* this volume). The analyses presented here also complement the preliminary scanning electron microscopy and synchrotron X-ray analyses in Saleh *et al.* (2024b). Our findings demonstrate that the Cabrières Biota exhibits a wide range of taphonomic modes, including pyritization, silicification, and preservation of organic material. This diversity of taphonomic modes is likely linked to the unique geological history of Early Ordovician deposits in southern France.

Material and methods

A field campaign was conducted in April 2024, during which hundreds of specimens were collected from the study area. All collected specimens were obtained from surface outcrops, as no drilling was conducted to obtain fresh rock samples. The material was then examined for fossil content, colouration, and general morphological features at the Institute of Earth Sciences of the University of Lausanne, using a standard optical microscope.

Preliminary observations revealed that some fossils exhibit little colour contrast relative to the surrounding sedimentary matrix while others display distinct hues, ranging from yellow, orange, brown, and occasionally, black (Fig. 1). From this initial screening, twelve representative fossils exhibiting varying types of preservation were selected for more detailed investigation using scanning electron microscopy (SEM)-based imaging and spectroscopy. More than 300 energy-dispersive X-ray spectroscopy (EDS) spectra were acquired across these twelve samples to characterize their elemental composition. Out of these, three specimens (sponge spicules UCBL-FSL717878; filamentous algae UCBL-FSL717888a, b; vermiform organisms UCBL-FSL717862) were chosen for detailed EDS mapping based on their contrasting spectral data.

SEM analyses were carried out using a Zeiss GeminiSEM 500 instrument operated under variable pressure conditions. The instrument is a joint facility of the École Polytechnique Fédérale de Lausanne (EPFL) and the University of Lausanne (UNIL), and is housed at the Géopolis building on the University




Fig. 1. Fossils from the Cabrières Biota: sponges (A: UCBL-FSL713619; B: UCBL-FSL717878), algae (C, D: UCBL-FSL717888a, b), and vermiform organisms (E: UCBL-FSL713622; F: UCBL-FSL717862). Note the variation in preservation, with specimen colours ranging from matrix-like green-blue to yellow, orange, red, brown, and even black. UCBL: University Claude Bernard Lyon 1 collections.

of Lausanne campus, Switzerland. Samples were mounted on standard SEM stubs using carbon adhesive tape and were not coated with any conductive metal layer to preserve surface integrity. Imaging and analysis were conducted in a chamber pressure of 40 Pa, using an accelerating voltage of 10 kV and a 30 μ m aperture. A backscattered electron detector (BSD X-Max 150) was employed to enhance material contrast. EDS mapping was performed using the Oxford AZtec Microanalysis System, version 4.2 SP1.

All material is housed in the public collections of the Claude Bernard University Lyon 1 (France). The specimens were legally collected during the field campaign from public land owned by the Cabrières municipality, which signed an agreement authorizing palaeontological exploration and rock material collection on its land. This agreement was made with the Claude Bernard University Lyon 1, the University of Brest (France), and the University of Lausanne (Switzerland). The signed convention, along with the associated documentation

and precise geographical coordinates of the collection sites, is stored with the fossil material at the Claude Bernard University Lyon 1.

Results

The analysed algae from the Cabrières Biota (Fig. 1C) show an enrichment in carbon (Fig. 2A, C), particularly along their marginal parts (Fig. 3A). In some areas, the algae exhibit silicon enrichment compared to other regions, particularly in the central portions of the fossils (Fig. 3A). Note that the silicon enrichment in the fossils differs from that observed in the matrix, as silicon in the matrix is associated with aluminium, unlike in the fossils (Fig. 3A). Where no enrichment in silicon or carbon is observed, iron can be found instead (Fig. 3A, D). Hence, the algae exhibit multiple preservation modes, consistent with the data and taphonomic models for fossil algae presented by Vayda *et al.* (this volume).

The analysed sponge from the Cabrières Biota (Fig. 1B) shows no carbonaceous remains and, interestingly, no silicon enrichment, despite the siliceous nature of its spicules (Li & Reitner in this volume). The original siliceous material of these spicules has been replaced by iron oxide minerals (Fig. 2C–E). These iron oxide minerals are generally euhedral in shape and may be coated by smaller, poorly crystallized minerals on their surfaces (Fig. 2E). In some cases, the euhedral minerals were completely dissolved, leaving negative imprints where they were originally present (Fig. 2E). It is also possible for some of these iron-rich minerals to show some enrichment in elements like cobalt (Fig. 3C).

The analysed vermiform organism from the Cabrières Biota (Fig. 1F) also shows a comparable iron enrichment (Fig. 3F), manifested as small euhedral minerals and larger framboids (Fig. 2F–H). No carbonaceous material or silicon enrichment is associated with these vermiform fossils. Interestingly, in some cases, soft-bodied fossils from the Cabrières Biota show no visible contrast from the surrounding matrix, even to the naked eye (Fig. 1E), suggesting an extensive degree of weathering. Nevertheless, some specimens retain minute morphological details at the micron scale, as revealed by electron microscopy data (Vannier *et al.* in this volume).

Discussion

The investigated fossils from the Cabrières Biota exhibit a variety of preservation modes (Figs 2, 3).

While all fossils contain iron oxide minerals of varying morphologies (Fig. 2C–H), only a limited number also show evidence of carbon and silicon (Fig. 3A–C).

The presence of euhedral and framboid-like minerals (Fig. 2E, H) suggests that iron oxides might result from early diagenetic pyrite precipitation (Gabbott *et al.* 2004; Zhu *et al.* 2005; Farrell 2014; Ma *et al.* 2015; Saleh *et al.* 2020d). Pyrite precipitation is commonly induced by the decay of soft tissues under sulphate-reducing conditions, leading to the replication of organic matter by pyrite crystals that are geologically more stable than soft tissues (Farrell 2014). This mode of preservation is the principal mechanism at certain early Palaeozoic sites, such as the Cambrian Chengjiang Biota (China; Gabbott *et al.* 2004; Zhu *et al.* 2005; Lei *et al.* 2024), the Ordovician Fezouata Biota (Morocco; Saleh *et al.* 2020d; 2021a), and the Beecher's trilobite bed (USA; Briggs *et al.* 1991). It also occurs as an accessory preservation mode in other Lagerstätten, including the Cambrian Burgess Shale Biota (Canada; Gaines *et al.* 2008), and several Cretaceous deposits such as the Crato Biota (Brazil; Barling *et al.* 2023), and the Lagerstätten of Haqel, Hjoula, and Sahel Alma (Lebanon; George *et al.* 2024), amongst others. The replication of fossils by pyrite is even associated with some of the oldest known evidence of life on Earth (Palaeoproterozoic, Gabon; Marin-Carbonne *et al.* 2022; Baumgartner *et al.* 2024; El Khoury *et al.* 2025a, b).

Depending on the availability of organic matter, sulphate and iron, pyrite can form either as dispersed euhedral crystals or as aggregates in the form of framboids (Gabbott *et al.* 2004). Pyrite distribution and shape are often indicative of the biotic and abiotic conditions under which pyritization occurred (El Khoury *et al.* 2025b). For example, under most conditions, anatomical parts that resist decay (e.g. bone or shell) do not reduce sulphates and are unlikely to get pyritized in comparison to soft tissues (e.g. nervous and digestive tissues) (Gabbott *et al.* 2004; Farrell 2014). The incomplete pyritization observed in some fossils (Fig. 3A) could also reflect a closed system with a limited supply of the components necessary for pyrite formation, particularly iron or sulphates (El Khoury *et al.* 2025b). However, in marine settings such as those of the Cabrières Biota (Vaucher *et al.* in press), sulphate availability is generally not a limiting factor due to its abundance in seawater (Berner 2004; Algeo *et al.* 2015). Previous studies have suggested that carbonate cementation around decaying carcasses can inhibit the diffusion of sulphates to the decaying material, thereby preventing extensive pyritization (Gaines *et al.* 2012). However, no evidence of such carbonate cements is observed in the Cabrières Biota.




Fig. 2. BSE images of investigated fossils: algae (A, B), sponges (C–E), and vermiform organism (F–H). Fossil algae exhibit carbon-rich material (crm; A, B) and matrix-like material (mlm; A). The mlm lacks carbon and has a similar chemical composition to the surrounding matrix but a distinct texture. Longitudinal sponge spicules (ss; C) are enriched in euhedral iron oxides (eio; D). In some areas, dissolved iron oxides (dio; D, E) have left negative imprints where they were previously present. Euhedral iron oxides co-occur with small iron oxides (sio; D), which may also cover the surface of larger euhedral crystals (E). Vermiform organism fossils are extensively covered by euhedral, small, and framoid-like iron oxides (flio; F, G). Negative imprints are also observed where framoid-like iron oxides have dissolved (dflio; G). These framoid-like structures lack the typical framoidal morphology, characterized by aggregates of small euhedral crystals, and appear as spheres with irregular borders (H). The analysed algal filaments are those shown in Figure 1C; the analysed sponge is the one in Figure 1B; and the analysed vermiform organism is the one in Figure 1F.

Therefore, the limited pyritization (Fig. 3A) is more likely attributable to low environmental iron availability or a change in redox conditions. In such a scenario, anatomical features not replicated by pyrite retained their organic signatures. If decay was halted, due to

increased burial rates (Schiffbauer *et al.* 2014) or other factors such as specific mineralogical signatures in the sediment (McMahon *et al.* 2016; Anderson *et al.* 2018; Corthésy *et al.* 2024; 2025a), these parts could be preserved in carbon, as observed in some Cabrières fossils




Fig. 3. Elemental composition of the investigated fossils: mapping of the algae (A), vermiform organism (B), and sponge (C), and spectra from the sponge specimen (D). Elemental mapping of fossil algae (A) shows that the surrounding matrix is enriched in aluminium and silicon, while the fossil itself is enriched in carbon and iron, with silicon in some places. Note that silicon within the fossil does not correlate with aluminium, unlike in the matrix. Framboid-like structures in the vermiform organism fossil are iron-rich (B), similar to the composition of euhedral minerals observed in the sponge fossil (C), which also show enrichment in trace elements such as cobalt. Overall, the matrix exhibits the highest concentrations of aluminium and silicon, while carbon and iron concentrations within the fossils vary depending on the analysed region (D). The spectra in (D) correspond to the boxed areas of similar colour labeled 'spectrum' (spect.) in (A).

(Fig. 3A; refer to Vayda *et al.* (this volume) for further evidence of carbon enrichment). However, it is worth noting that the degree of compression in the carbon-rich material is not as pronounced as in Burgess Shale-type material (e.g. Butterfield 1990, 1995; Butterfield *et al.* 2007; Orr *et al.* 2009; Gaines *et al.* 2008); as the material still retains three-dimensionality (Fig. 2B) refer to Vayda *et al.* (this volume), for further information on the taphonomic pathways of algal filaments.

The presence of silicon in fossils (Fig. 3A; refer to Vayda *et al.* (this volume), for further evidence of local silicon enrichment in some algal filaments), together with pyrite crystals and carbon-rich material, is common in the fossil record. Similar associations are found in the Burgess Shale (Orr *et al.* 1998; Anderson *et al.* 2021), and experimental studies suggest that silicon can bind to organic material during early diagenesis

(Newman *et al.* 2019; Slatger *et al.* 2024; Corthésy *et al.* 2025a). In fact, under highly reactive conditions, silicon may precipitate on the surfaces of decaying organisms in the shape of amorphous, poorly crystallized minerals, enhancing their resistance to decay (Corthésy *et al.* 2025a). These silicon-rich minerals can be distinguished from those in the surrounding matrix based on their chemical composition (Corthésy *et al.* 2025a). In the Cabrières Biota, for example, silicon is not associated with aluminium in fossils, unlike the surrounding aluminosilicate-rich matrix (Fig. 3A, D).

The presence of reactive conditions during early diagenesis in the Cabrières Biota is further evidenced by the replacement of originally mineralized structures, such as sponge spicules, by pyrite (Figs 2C–E, 3C). Sponge spicules contain silicon; however, Cabrières spicules have been entirely replaced by

pyrite minerals (Figs 2C–E, 3C). Although sponge spicules are not among the most labile anatomical features, the pyritization of these elements is commonly observed in other exceptionally preserved biotas, such as the Ordovician Anji Biota (China; Wu *et al.* 2022). In the Anji Biota, it is suggested that the sulphate-reducing conditions favourable to pyrite precipitation promoted the dissolution of silicon within sponge spicules (Wu *et al.* 2022). This process resulted in the release of biogenic silica into the surrounding environment (Wu *et al.* 2022). Under these conditions, biogenic silica could have contributed, alongside environmental (abiotic) silica, to the silicon enrichment in other anatomical features and organisms, as observed in algae from the Cabrières Biota (Fig. 3A).

The proposed taphonomic scenario suggests that pyritization preceded silicification, whereas the alternative case, where silicification occurred first, is considered unlikely. If silicification had taken place prior to pyrite formation, the resulting mineral hardening of the organic matter (from the replication of organic matter by silica) would have inhibited decay, thereby

preventing sulphate reduction (by decaying organic matter) and subsequent pyritization. As such, the most plausible sequence involves the dissolution of sponge spicules and their replacement by pyrite under sulphate-reducing conditions (Fig. 4A–C), followed by the rapid silicification of other fossils and anatomical features using the biogenic silica released during this process (Fig. 4D).

In this model, silicification occurs when conditions become non-permissive for pyrite precipitation (e.g. if the iron becomes limiting or following a change in redox conditions; Jauvion *et al.*, 2020; Odin *et al.* 2024; Corthésy *et al.* 2025b). The extent of silicification would then depend on the availability of silicon and the stability of environmental conditions that support this process (Maliva & Siever 1988; Moore *et al.* 2020). This sequence would result in a range of preservation modes (i.e. pyritized fossils that may also exhibit silicified parts, and carbon-rich material in areas lacking both pyrite and silicon; Fig. 4D) mirroring the diversity of preservation observed in the Cabrières Biota (Fig. 3A, D).




Fig. 4. Chemical taphonomic pathway for fossils from the Early Ordovician Cabrières Biota. Dead organisms from a single assemblage containing silica-rich sponge spicules and algae start decaying on the seafloor (A). During early diagenesis, reducing conditions are established, promoting the release of silicon from sponge spicules and the reduction of seawater sulphates into sulphides by decaying organic material (B). These sulphides react with iron from the environment, leading to the precipitation of pyrite (coloured purple) within organisms, as long as the conditions remain favourable (e.g. sufficient iron availability or stable redox conditions) (C). Biogenic silicon, when available, may contribute to the silicification (coloured grey) of previously non-pyritized parts of dead organisms; carbonaceous material may persist in regions that were not enriched in iron or silicon (D). During late diagenesis, and under metamorphic activity and surface weathering conditions, pyrite crystals dissolve to varying extents and become oxidized, releasing a secondary source of iron into the environment (E). This newly released iron can reprecipitate, even in areas that were not originally pyritized, as poorly crystalline iron minerals, and coat the surface of earlier-formed euhedral pyrite crystals. This stage can also involve the leaching or removal of some carbonaceous material (F). In such regions, the matrix beneath the fossil becomes exposed, showing no significant chemical difference from the surrounding matrix. Nonetheless, structural differences in mineral arrangement may still distinguish the fossil-bearing zone from the surrounding sediment (refer to Fig. 2A for further details).

Following pyritization and silicification, fossils from the Cabrières Biota underwent significant weathering. Sulphur was completely leached from pyrite crystals as no sulphur was detected in any of the obtained spectra (Fig. 3D). Some pyrite crystals were either partially dissolved, resulting in altered framboids that lack distinct individual crystals and appear as wrinkled spheres (Fig. 2H), or completely dissolved (Figs 2D, E, G, 4E). Iron was subsequently mobilized and reprecipitated as poorly crystallized iron oxides (Figs 2D, E, 4F). These can be distinguished from original pyrite crystals by their lack of defined shape and tendency to coat the surfaces of both euhedral and framboidal minerals (Figs 2E, 4F).

Similar patterns of pyrite dissolution are observed in other exceptionally preserved fossil sites, such as the Anji Biota (Wu *et al.* 2022). In the Fezouata Biota, abundant poorly crystallized iron oxides often cover fossil surfaces (Saleh *et al.* 2021a). The association of cobalt with some Cabrières Biota fossils (Fig. 3C) is also evidenced in the Fezouata Biota and is thought to reflect intense weathering processes (Saleh *et al.* 2020b). This extensive weathering may also explain the lack of carbon enrichment in certain fossil parts from the Cabrières Biota (Figs 2A, 4F). In some specimens, these areas are chemically indistinguishable from the surrounding matrix (Figs 2A, 4F), with fossil structures only discernible through the specific organization of underlying matrix minerals where carbon has been leached [Fig. 1E; refer to Vayda *et al.* (in this volume) for further examples]. In brief, the oxidation of pyrite results in fossils displaying orange and red colours. When carbon is still present (Fig. 3A), fossils appear black (Fig. 1C), while when carbon has been leached and no pyrite formed, fossil casts exhibit more or less the same colour as the surrounding matrix (Fig. 1E).

The pronounced alteration affecting the Cabrières Biota may be linked to the metamorphic history of the region, including multiple episodes of fluid circulation and relatively elevated burial temperatures (Álvaro *et al.* 2008; Roger *et al.* 2020). Present-day conditions at these forested outcrops further contribute to their weathering through continued fluid movement.

Conclusion

Exceptionally preserved fossils of the Cabrières Biota result from a complex interplay of taphonomic processes. Pyrite precipitation, primarily driven by the decay of soft tissues under sulphate-reducing conditions, was likely the dominant initial preservation mechanism. Pyritization was followed by the partial

silicification of the organisms, potentially facilitated by biogenically sourced silicon. Tissues that were neither silicified nor pyritized were preserved as carbon-rich material, suggesting that elements such as iron and silicon were limited in the early diagenetic environment. The complete leaching of sulphur, dissolution of pyrite crystals, and loss of carbon in some specimens point to a substantial alteration due to late diagenesis, metamorphic processes, and surface weathering conditions, which have obscured aspects of the original modes of preservation. Understanding the details of this taphonomic pathway and how it may have biased our knowledge of this unique ecosystem requires further high-resolution taphonomic analyses. For instance, it would be interesting in the future to investigate whether specific anatomical features are preferentially preserved through particular modes of preservation (e.g. pyritization or silicification). It would also be important to quantify the biological structures preserved in this Lagerstätte, in a manner similar to previous studies of other sites with exceptional fossil preservation (Saleh *et al.* 2022c, 2023c; Whitaker *et al.*, 2022). Elemental and isotopic analyses aimed at investigating the origin of elements such as iron, as well as broader depositional factors like redox conditions, are equally important (e.g. Gaines *et al.*, 2012; Schiffbauer *et al.*, 2014; Hammarlund *et al.*, 2017, 2019; Qi *et al.*, 2018). However, at this stage, the preliminary results presented here, which point to a complex preservation pathway behind the exceptional fossil record of this Early Ordovician polar ecosystem, represent a crucial first step toward achieving such an understanding.

Acknowledgements. – We thank all participants in the fieldwork of April 2024. Jean-Paul Kundura and Marie-Hélène Kundura are thanked for donating material to the Claude Bernard University Collections. FS and NC are funded by an SNF Ambizione Grant (PZ00P2_209102). SL and HBD are funded by an SNF Sinergia Grant (198691). DP and VJ are funded by an SNF Ambizione grant (PZ00P2_193520). GJMP thanks funding from the Canton of Vaud (Switzerland). PaG is funded by the project ECO-BOOST of the French National Research Agency (ANR-22-CE01-0003). This paper is a contribution to the International Geoscience Programme (IGCP) Project 735—Rocks and the Rise of Ordovician Life (Rocks n'ROL) and to the INSU-Tellus 'LAGDOC' project. We thank the Editor and the two reviewers for their comments that helped us improve the manuscript.

References

- Algeo, T.J., Luo, G.M., Song, H.Y., Lyons, T.W. & Canfield, D.E. 2015: Reconstruction of secular variation in seawater sulfate concentrations. *Biogeosciences* 12, 2131–2151.
- Álvaro, J.J., Bauluz, B., Subías, I., Pierre, C. & Vizcaino, D. 2008: Carbon chemostratigraphy of the Cambrian-Ordovician transition in a midlatitude mixed platform, Montagne Noire, France. *Geological Society of America Bulletin* 120, 962–975.

- Anderson, R.P., Tosca, N.J., Gaines, R.R., Koch, N.M. & Briggs, D.E. 2018: A mineralogical signature for Burgess Shale-type fossilization. *Geology* 46, 347–350.
- Anderson, R.P., Tosca, N.J., Saupe, E.E., Wade, J. & Briggs, D.E. 2021: Early formation and taphonomic significance of kaolinite associated with Burgess Shale fossils. *Geology* 49, 355–359.
- Barling, N., Saleh, F. & Ma, X. 2023: A unique record of prokaryote cell pyritization. *Geology* 51, 1062–1066.
- Baumgartner, R.J., Van Kranendonk, M.J., Caruso, S., Campbell, K.A., Dobson, M.J., Teece, B.L., Verrall, M., Homann, M., Lalonde, S. & Visscher, P.T. 2024: Pyritic stromatolites from the Paleoproterozoic Dresser Formation, Pilbara Craton: Resolving biogenicity and hydrothermally influenced ecosystem dynamics. *Geobiology* 22, e12610.
- Berner, R.A. 2004: A model for calcium, magnesium and sulfate in seawater over Phanerozoic time. *American Journal of Science* 304, 438–453.
- Briggs, D.E., Bottrell, S.H. & Raiswell, R. 1991: Pyritization of soft-bodied fossils: Beecher's trilobite bed, Upper Ordovician, New York State. *Geology* 19, 1221–1224.
- Butterfield, N.J. 1990: Organic preservation of non-mineralizing organisms and the taphonomy of the Burgess Shale. *Paleobiology* 16, 272–286.
- Butterfield, N.J. 1995: Secular distribution of Burgess-Shale-type preservation. *Lethaia* 28, 1–13.
- Butterfield, N.J., Balthasar, U.W.E. & Wilson, L.A. 2007: Fossil diagenesis in the Burgess Shale. *Palaeontology* 50, 537–543.
- Caillaud, V., Birolini, E., Manzano, E.G., Michel, S., Regnier, S., Vaucher, R., & Servais, T. 2026: Acritarchs (organic-walled microfossils) from the Floian (Early Ordovician) of the Montagne Noire, France: biostratigraphy, palaeobiogeography and palaeoecology. *Lethaia*, in press.
- Corthésy, N., Saleh, F., Thomas, C., Antcliffe, J.B. & Daley, A.C. 2024: The effects of clays on bacterial community composition during arthropod decay. *Swiss Journal of Palaeontology* 143, 26.
- Corthésy, N., Saleh, F., Antcliffe, J.B. & Daley, A.C. 2025a: Kaolinite induces rapid authigenic mineralisation in unburied shrimps. *Communications Earth & Environment* 6, 4.
- Corthésy, N., Antcliffe, J.B. & Saleh, F. 2025b: Taxon-specific redox conditions control fossilisation pathways. *Nature Communications* 16, 3993.
- Daley, A.C., Gueriau, P., Lustri, L., Drage, H.B., Lynch, S., Monceret, E., & Monceret, S. 2026: Euarthropods from the Early Ordovician Cabrières Biota, including a new aglaspidid and freshly moulted trilobite material. *Lethaia*, in press.
- Drage, H.B., Legg, D.A. & Daley, A.C. 2023: Novel marrellomorph moulting behaviour preserved in the Lower Ordovician Fezouata Shale, Morocco. *Frontiers in Ecology and Evolution* 11, 1226924.
- Dupichaud, C., Lefebvre, B., Milne, C.H., Mooi, R., Nohejlová, M., Roch, R., Saleh, F. & Zamora, S. 2023: Solatan echinoderms from the Fezouata Shale Lagerstätte (Lower Ordovician, Morocco): Diversity, exceptional preservation, and palaeoecological implications. *Frontiers in Ecology and Evolution* 11, 1290063.
- El Khoury, A., Somogyi, A., Chi Fru, E., Saleh, F., Chraiki, I., Fontaine, C., Aubineau, J., Rollion-Bard, C., Harzhauser, M. & El Albani, A. 2025a: A battle against arsenic toxicity by Earth's earliest complex life forms. *Nature Communications* 16, 1–14.
- El Khoury, A., Saleh, F., El Albani, A., Fontaine, C., Rollion-Bard, C., Chraiki, I., Aubineau, J., Ngwal'ghoubou Ikouanga, J., Bhlisse, M., Zguaid, M. & Somogyi, A. 2025b: Pyrite morphology and sulfur isotopes refine taphonomic models for the 2.1 Ga Francevillian biota, Gabon. *Scientific Reports* 15, 1–11.
- Farrell, U.C. 2014: Pyritization of soft tissues in the fossil record: an overview. *The Paleontological Society Papers* 20, 35–58.
- Gabbott, S.E., Xian-Guang, H., Norry, M.J. & Siveter, D.J. 2004: Preservation of Early Cambrian animals of the Chengjiang biota. *Geology* 32, 901–904.
- Gaines, R.R., Briggs, D.E. & Yuanlong, Z. 2008: Cambrian Burgess Shale-type deposits share a common mode of fossilization. *Geology* 36, 755–758.
- Gaines, R.R., Hammarlund, E.U., Hou, X., Qi, C., Gabbott, S.E., Zhao, Y., Peng, J. & Canfield, D.E. 2012: Mechanism for Burgess Shale-type preservation. *Proceedings of the National Academy of Sciences* 109, 5180–5184.
- George, H., Bazzi, M., El Hossny, T., Ashraf, N., Abi Saad, P. & Clements, T. 2024: The famous fish beds of Lebanon: the Upper Cretaceous Lagerstätten of Haqel, Hjoula, Nammoura and Sahel Aalma. *Journal of the Geological Society* 181, jgs2023–210.
- Gougeon, R., Vaucher, R., Vidal, M., Monceret, E., Monceret, S., Birolini, E., Dupichaud, C., Lefebvre, B., & Saleh, F. 2026: Ichnology of the Lower Ordovician Landeyran Formation in Montagne Noire, southern France, and criteria for distinguishing simple trace fossils from body fossils. *Lethaia*, in press.
- Gutiérrez-Marco, J.C., García-Bellido, D.C., Rábano, I. & Sá, A.A. 2017: Digestive and appendicular soft-parts, with behavioural implications, in a large Ordovician trilobite from the Fezouata Lagerstätte, Morocco. *Scientific Reports* 7, 39728.
- Gutiérrez-Marco, J.C., Maletz, J., Vizcaino, D., Lefebvre, B. & Saleh, F. 2026: Lower Ordovician graptolites from the Cabrières region, Montagne Noire, France. *Lethaia*, in press.
- Hammarlund, E.U., Gaines, R.R., Prokopenko, M.G., Qi, C., Hou, X.G. & Canfield, D.E. 2017: Early Cambrian oxygen minimum zone-like conditions at Chengjiang. *Earth and Planetary Science Letters* 475, 160–168.
- Hammarlund, E.U., Smith, M.P., Rasmussen, J.A., Nielsen, A.T., Canfield, D.E. & Harper, D.A. 2019: The Sirius Passet Lagerstätte of North Greenland—a geochemical window on early Cambrian low-oxygen environments and ecosystems. *Geobiology* 17, 12–26.
- Harper, D.A., Hammarlund, E.U. & Rasmussen, C.M. 2014: End Ordovician extinctions: a coincidence of causes. *Gondwana Research* 25, 1294–1307.
- Harper, D.A., Zhan, R.B. & Jin, J. 2015: The Great Ordovician Biodiversification Event: reviewing two decades of research on diversity's big bang illustrated by mainly brachiopod data. *Palaeoworld* 24, 75–85.
- Harper, D.A.T., Candela, Y., Monceret, E., Monceret, F., Birolini, E. & Lefebvre, B. 2026: Brachiopods from the Cabrières biota (Early Ordovician, southern France): ecology and taphonomy. *Lethaia*, in press.
- Jauvin, C., Bernard, S., Gueriau, P., Mocuta, C., Pont, S., Benzerara, K. & Charbonnier, S. 2020: Exceptional preservation requires fast biodegradation: thylacocephalan specimens from La Voulte-sur-Rhône (Callovian, Jurassic, France). *Palaeontology* 63, 395–413.
- Kimmig, J. & Schiffbauer, J.D. 2024: A modern definition of Fossil Lagerstätten. *Trends in Ecology & Evolution* 39, 621–624.
- Laibl, L., Gueriau, P., Saleh, F., Pérez-Peris, F., Lustri, L., Drage, H.B., Bath Enright, O.G., Potin, G.J.M. & Daley, A.C. 2023a: Early developmental stages of a Lower Ordovician marrellid from Morocco suggest simple ontogenetic niche differentiation in early euarthropods. *Frontiers in Ecology and Evolution* 11, 1232612.
- Laibl, L., Drage, H.B., Pérez-Peris, F., Schöder, S., Saleh, F. & Daley, A.C. 2023b: Babies from the Fezouata Biota: early developmental trilobite stages and their adaptation to high latitudes. *Geobios* 81, 31–50.
- Lefebvre, B., Guensburg, T.E., Martin, E.L., Mooi, R., Nardin, E., Nohejlová, M., Saleh, F., Kouraïss, K., El Hariri, K. & David, B. 2019: Exceptionally preserved soft parts in fossils from the Lower Ordovician of Morocco clarify stylophoran affinities within basal deuterostomes. *Geobios* 52, 27–36.
- Lefebvre, B., Dupichaud, C., Vidal, M., Servais, T., Michel, S., Nardin, E., Birolini, E., Caillaud, V., De Oliveira Santos, V., Gougeon, R., Guenser, P., Monceret, E., Monceret, S., Nohejlová, M., Vaucher, R., & Saleh, F. Faunal distribution and palaeoenvironmental trends in the Lower Ordovician of the Montagne Noire: Insights from the Cabrières Biota. *Lethaia*, in press.

- Lei, X., Cong, P., Zhang, S., Wei, F. & Anderson, R.P. 2024: Unveiling an ignored taphonomic window in the early Cambrian Chengjiang Biota. *Geology* 52, 753–758.
- Li, L. & Reitner, J. 2026: Hexactinellid Sponges from the Early Ordovician Cabrières Biota, France: insights into Early Paleozoic Polar Ecosystems. *Lethaia*, in press.
- Lustri, L., Gueriau, P. & Daley, A.C. 2024: Lower Ordovician syn-ziphosurine reveals early euclerater diversity and evolution. *Nature Communications* 15, 3808.
- Ma, X., Edgecombe, G.D., Hou, X., Goral, T. & Strausfeld, N.J. 2015: Preservation pathways of corresponding brains of a Cambrian euarthropod. *Current Biology* 25, 2969–2975.
- Maliva, R.G. & Siever, R. 1988: Mechanism and controls of silicification of fossils in limestones. *Journal of Geology* 96, 387–398.
- Marin-Carbonne, J., Decraene, M.N., Havas, R., Remusat, L., Pasquier, V., Alléon, J., Zeyen, N., Bouton, A., Bernard, S., Escriu, S. & Olivier, N. 2022: Early precipitated micropyrith in microbialites: a time capsule of microbial sulfur cycling. *Geochemical Perspectives Letters* 21, 7–12.
- Martin, E.L., Pittet, B., Gutiérrez-Marco, J.C., Vannier, J., El Hariri, K., Leroisey-Aubril, R., Masrour, M., Nowak, H., Servais, T., Vandenbroucke, T.R. & Van Roy, P. 2016: The lower Ordovician Fezouata Konservat-Lagerstätte from Morocco: age, environment and evolutionary perspectives. *Gondwana Research* 34, 274–283.
- McMahon, S., Anderson, R.P., Saupe, E.E. & Briggs, D.E. 2016: Experimental evidence that clay inhibits bacterial decomposers: Implications for preservation of organic fossils. *Geology* 44, 867–870.
- Monceret, S., Monceret, E., Saleh, F. & Lefebvre, B. 2026: The Ordovician of the Cabrières area (Hérault, France): A brief overview of two centuries of palaeontological investigations. *Lethaia*, in press.
- Moore, K.R., Pajusalu, M., Gong, J., Sojo, V., Matreux, T., Braun, D. & Bosak, T. 2020: Biologically mediated silicification of marine cyanobacteria and implications for the Proterozoic fossil record. *Geology* 48, 862–866.
- Newman, S.A., Daye, M., Fakra, S.C., Marcus, M.A., Pajusalu, M., Pruss, S.B., Smith, E.F. & Bosak, T. 2019: Experimental preservation of muscle tissue in quartz sand and kaolinite. *Palaios* 34, 437–451.
- Odin, G.P., Leloup, J., Tu, T.T.N., Charbonnier, S. & Derenne, S. 2024: Taphonomic experiments on caridean shrimp elucidate the timing and influence of organic matter on diagenetic pyritization and sulfurization. *Chemical Geology* 644, 121858.
- Orr, P.J., Briggs, D.E. & Kearns, S.L. 1998: Cambrian Burgess Shale animals replicated in clay minerals. *Science* 281, 1173–1175.
- Orr, P.J., Kearns, S.L. & Briggs, D.E. 2009: Elemental mapping of exceptionally preserved ‘carbonaceous compression’ fossils. *Palaeogeography, Palaeoclimatology, Palaeoecology* 277, 1–8.
- Parry, L.A., Smithwick, F., Nördén, K.K., Saitta, E.T., Lozano-Fernandez, J., Tanner, A.R., Caron, J.B., Edgecombe, G.D., Briggs, D.E. & Vinther, J. 2018: Soft-bodied fossils are not simply rotten carcasses—toward a holistic understanding of exceptional fossil preservation: exceptional fossil preservation is complex and involves the interplay of numerous biological and geological processes. *BioEssays* 40, 1700167.
- Perez-Peris, F., Laibl, L., Vidal, M. & Daley, A. 2021: Systematics, morphology, and appendages of an Early Ordovician pilekiine trilobite *Anacheirurus* from Fezouata Shale and the early diversification of Cheiruridae. *Acta Palaeontologica Polonica* 66, 857–877.
- Polechová, M. 2016: The bivalve fauna from the Fezouata Formation (Lower Ordovician) of Morocco and its significance for palaeobiogeography, palaeoecology and early diversification of bivalves. *Palaeogeography, Palaeoclimatology, Palaeoecology* 460, 155–169.
- Polechová, M., Ebbestad, J.O.R. & Kröger, J. 2026: Molluscs of the Lower Ordovician Cabrières Lagerstätte, France – diversity and distribution. *Lethaia*, in press.
- Potin, G.J.M., Gueriau, P. & Daley, A.C. 2023: Radiodont frontal appendages from the Fezouata Biota (Morocco) reveal high diversity and ecological adaptations to suspension-feeding during the Early Ordovician. *Frontiers in Ecology and Evolution* 11, 1214109.
- Purnell, M.A., Donoghue, P.J., Gabbott, S.E., McNamara, M.E., Murdock, D.J. & Sansom, R.S. 2018: Experimental analysis of soft-tissue fossilization: opening the black box. *Palaeontology* 61, 317–323.
- Qi, C., Li, C., Gabbott, S.E., Ma, X., Xie, L., Deng, W., Jin, C. & Hou, X.G. 2018: Influence of redox conditions on animal distribution and soft-bodied fossil preservation of the Lower Cambrian Chengjiang Biota. *Palaeogeography, Palaeoclimatology, Palaeoecology* 507, 180–187.
- Rasmussen, C.M., Kröger, B., Nielsen, M.L. & Colmenar, J. 2019: Cascading trend of Early Paleozoic marine radiations paused by Late Ordovician extinctions. *Proceedings of the National Academy of Sciences* 116, 7207–7213.
- Roger, F., Teyssier, C., Whitney, D.L., Respaut, J.P., Paquette, J.L. & Rey, P.F. 2020: Age of metamorphism and deformation in the Montagne Noire dome (French Massif Central): Tapping into the memory of fine-grained gneisses using monazite U-Th-Pb geochronology. *Tectonophysics* 776, 228316.
- Saleh, F., Candela, Y., Harper, D.A., Polechová, M., Lefebvre, B. & Pittet, B. 2018: Storm-induced community dynamics in the Fezouata Biota (Lower Ordovician, Morocco). *Palaios* 33, 535–541.
- Saleh, F., Pittet, B., Perrillat, J.P. & Lefebvre, B. 2019: Orbital control on exceptional fossil preservation. *Geology* 47, 103–106.
- Saleh, F., Lefebvre, B., Hunter, A.W. & Nohejlová, M. 2020a: Fossil weathering and preparation mimic soft tissues in eocrinoid and somasteroid echinoderms from the Lower Ordovician of Morocco. *Microscopy Today* 28, 24–28.
- Saleh, F., Pittet, B., Sansjofre, P., Guériau, P., Lalonde, S., Perrillat, J.P., Vidal, M., Lucas, V., El Hariri, K., Kourais, K. & Lefebvre, B. 2020b: Taphonomic pathway of exceptionally preserved fossils in the Lower Ordovician of Morocco. *Geobios* 60, 99–115.
- Saleh, F., Antcliffe, J.B., Lefebvre, B., Pittet, B., Laibl, L., Peris, F.P., Lustri, L., Gueriau, P. & Daley, A.C. 2020c: Taphonomic bias in exceptionally preserved biotas. *Earth and Planetary Science Letters* 529, 115873.
- Saleh, F., Daley, A.C., Lefebvre, B., Pittet, B. & Perrillat, J.P. 2020d: Biogenic iron preserves structures during fossilization: a hypothesis: iron from decaying tissues may stabilize their morphology in the fossil record. *BioEssays* 42, 1900243.
- Saleh, F., Vaucher, R., Antcliffe, J.B., Daley, A.C., El Hariri, K., Kourais, K., Lefebvre, B., Martin, E.L., Perrillat, J.P., Sansjofre, P. & Vidal, M. 2021a: Insights into soft-part preservation from the Early Ordovician Fezouata Biota. *Earth-Science Reviews* 213, 103464.
- Saleh, F., Vidal, M., Laibl, L., Sansjofre, P., Gueriau, P., Pérez-Peris, F., Lustri, L., Lucas, V., Lefebvre, B., Pittet, B. & El Hariri, K. 2021b: Large trilobites in a stress-free Early Ordovician environment. *Geological Magazine* 158, 261–270.
- Saleh, F., Bath-Enright, O.G., Daley, A.C., Lefebvre, B., Pittet, B., Vite, A., Ma, X., Mángano, M.G., Buatois, L.A. & Antcliffe, J.B. 2021c: A novel tool to untangle the ecology and fossil preservation knot in exceptionally preserved biotas. *Earth and Planetary Science Letters* 569, 117061.
- Saleh, F., Vaucher, R., Vidal, M., Hariri, K.E., Laibl, L., Daley, A.C., Gutiérrez-Marco, J.C., Candela, Y., Harper, D.A., Ortega-Hernández, J. & Ma, X. 2022a: New fossil assemblages from the Early Ordovician Fezouata biota. *Scientific Reports* 12, 20773.
- Saleh, F., Lucas, V., Pittet, B., Lefebvre, B., Lalonde, S.V. & Sansjofre, P. 2022b: Iron from continental weathering dictated soft-part preservation during the Early Ordovician. *Terra Nova* 34, 163–168.
- Saleh, F., Ma, X., Guenser, P., Mángano, M.G., Buatois, L.A. & Antcliffe, J.B. 2022c: Probability-based preservational variations within the early Cambrian Chengjiang biota (China). *PeerJ* 10, e13869.
- Saleh, F., Antcliffe, J.B., Lustri, L., Daley, A.C. & Gibert, C. 2023a: Cambrian and Ordovician diversity fluctuations could be resolved through a single ecological hypothesis. *Lethaia* 56, 1–13.

- Saleh, F., Lefebvre, B., Dupichaud, C., Martin, E.L., Nohejlová, M. & Spaccesi, L. 2023b: Skeletal elements controlled soft-tissue preservation in echinoderms from the Early Ordovician Fezouata Biota. *Geobios* 81, 51–66.
- Saleh, F., Clements, T., Perrier, V., Daley, A.C. & Antcliffe, J.B. 2023c: Variations in preservation of exceptional fossils within concretions. *Swiss Journal of Palaeontology* 142, 20.
- Saleh, F., Antcliffe, J.B., Birolini, E., Candela, Y., Corthésy, N., Daley, A.C., Dupichaud, C., Gibert, C., Guenser, P., Laibl, L. & Lefebvre, B. 2024a: Highly resolved taphonomic variations within the Early Ordovician Fezouata Biota. *Scientific Reports* 14, 20807.
- Saleh, F., Lustri, L., Gueriau, P., Potin, G.J.M., Pérez-Peris, F., Laibl, L., Jamart, V., Vite, A., Antcliffe, J.B., Daley, A.C. & Nohejlová, M., 2024b: The Cabrières Biota (France) provides insights into Ordovician polar ecosystems. *Nature Ecology & Evolution* 8, 651–662.
- Saleh, F., Lustri, L., Gueriau, P., Potin, G.J.M., Pérez-Peris, F., Laibl, L., Jamart, V., Vite, A., Antcliffe, J.B., Daley, A.C. & Nohejlová, M. 2024c: Reply to: The Cabrières Biota is not a Konservat-Lagerstätte. *Nature Ecology & Evolution* 8, 2175–2178.
- Schiffbauer, J.D., Xiao, S., Cai, Y., Wallace, A.F., Hua, H., Hunter, J., Xu, H., Peng, Y. & Kaufman, A.J. 2014: A unifying model for Neoproterozoic–Palaeozoic exceptional fossil preservation through pyritization and carbonaceous compression. *Nature Communications* 5, 5754.
- Seilacher, A., Reif, W.E. & Westphal, F. 1985: Sedimentological, ecological and temporal patterns of fossil Lagerstätten. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* 311, 5–24.
- Servais, T. & Harper, D.A. 2018: The great Ordovician biodiversification event (GOBE): definition, concept and duration. *Lethaia* 51, 151–164.
- Servais, T., Cascales-Miñana, B. & Harper, D.A. 2021: The Great Ordovician Biodiversification Event (GOBE) is not a single event. *Paleontological Research* 25, 315–328.
- Slagter, S., Konhauser, K.O., Briggs, D.E. & Tarhan, L.G. 2024: Controls on authigenic mineralization in experimental Ediacara-style preservation. *Geobiology* 22, e12615.
- Stigall, A.L., Edwards, C.T., Freeman, R.L. & Rasmussen, C.M. 2019: Coordinated biotic and abiotic change during the Great Ordovician Biodiversification Event: Darriwilian assembly of early Paleozoic building blocks. *Palaeogeography, Palaeoclimatology, Palaeoecology* 530, 249–270.
- Stigall, A.L., Censullo, S.M., Hennessey, S.A., Bauer, J.E., Lam, A.R. & Wright, D.F. 2023: Diversification and speciation among Laurentian brachiopods during the GOBE: insights from basinal and regional analyses. *Estonian Journal of Earth Sciences* 72, 98–101.
- Van Iten, H., Birolini, E., Dupichaud, C., & Lefebvre, B. 2026: Cnidarians of the Cabrières Lagerstätte (Lower Ordovician, France): Diversity, taphonomy, paleoecology, and paleobiogeography. *Lethaia*, in press.
- Van Iten, H., Muir, L., Simões, M.G., Leme, J.M., Marques, A.C. & Yoder, N. 2016: Palaeobiogeography, palaeoecology and evolution of Lower Ordovician conulariids and Sphenothallus (Medusozoa, Cnidaria), with emphasis on the Fezouata Shale of southeastern Morocco. *Palaeogeography, Palaeoclimatology, Palaeoecology* 460, 170–178.
- Van Roy, P., Orr, P.J., Botting, J.P., Muir, L.A., Vinther, J., Lefebvre, B., Hariri, K.E. & Briggs, D.E., 2010: Ordovician faunas of Burgess Shale type. *Nature* 465, 215–218.
- Vannier, J., Wang, D., Han, J., Birolini, E., Monceret, E., Monceret, S., Kundura, J.-P., Kundura, M.-H., & Lefebvre, B. 2026: Worms from the Early Ordovician (Floian) Cabrières Biota (Montagne-Noire, southern France), *Lethaia*, in press.
- Vaucher, R., Martin, E.L., Hormière, H. & Pittet, B. 2016: A genetic link between Konservat-Lagerstätten in the Fezouata Shale (lower Ordovician, Morocco). *Palaeogeography, Palaeoclimatology, Palaeoecology* 460, 24–34.
- Vaucher, R., Pittet, B., Hormière, H., Martin, E.L. & Lefebvre, B. 2017: A wave-dominated, tide-modulated model for the Lower Ordovician of the Anti-Atlas, Morocco. *Sedimentology* 64, 777–807.
- Vaucher, R., Michel, S., Pas, D., Gougeon, R., Vizcaíno, D., Lefebvre, B., & Saleh, F. 2026: The Cabrières Biota inhabited a polar wave-influenced delta. *Lethaia*, in press.
- Vayda, P., Birolini, E. & Xiao, S. 2026: Ediacaran-type Vendotaenid macroalgae in an Ordovician Lagerstätte. *Lethaia*, in press.
- Vinther, J., Van Roy, P. & Briggs, D.E. 2008: Machaeridians are Palaeozoic armoured annelids. *Nature* 451, 185–188.
- Vinther, J., Parry, L., Briggs, D.E. & Van Roy, P. 2017: Ancestral morphology of crown-group molluscs revealed by a new Ordovician stem aculiferan. *Nature* 542, 471–474.
- Whitaker, A.F., Schiffbauer, J.D., Briggs, D.E., Leibach, W.W. & Kimmig, J. 2022: Preservation and diagenesis of soft-bodied fossils and the occurrence of phosphate-associated rare earth elements in the Cambrian (Wuliuan) Spence Shale Lagerstätte. *Palaeogeography, Palaeoclimatology, Palaeoecology* 592, 110909.
- Wu, X., Botting, J.P., Zhang, Y., Muir, L.A. & Ma, J.Y. 2022: Taphonomy of exceptional sponge preservation in the Hirnantian Anji Biota, China. *Journal of the Geological Society* 179, jgs2021–112.
- Zhu, M., Babcock, L.E. & Steiner, M. 2005: Fossilization modes in the Chengjiang Lagerstätte (Cambrian of China): testing the roles of organic preservation and diagenetic alteration in exceptional preservation. *Palaeogeography, Palaeoclimatology, Palaeoecology* 220, 31–46.