

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

# K–excess, an unexploited method for discovering red quasars

Thodori Nakos

Université de Liège  
Institut d’Astrophysique & Géophysique

J. Willis (Univ. Victoria), S. Andreon (Oss. Brera)  
J. Surdej (Univ. Liege)

# Outline

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

- 1 Scientific motivation
- 2 The data sets – Selection process
- 3 Properties of the quasar candidates
- 4 Current / Future work

# QSO selection in optical imaging surveys: The UV excess method (UVX)

Scientific  
motivation

The data sets  
– Selection  
process


Properties of  
the quasar  
candidates

Current /  
Future work

QSO SED: power law  
Stellar SED: convex  $\Rightarrow$

Quasars show UV flux  
excess  $\Rightarrow$

In  $(U - B)$  vs  $(B - V)$   
plane quasars occupy  
separate region!




# The UVX method (2)

Scientific motivation

The data sets  
– Selection process

Properties of the quasar candidates

Current /  
Future work



(QSO spectrum: Vanden Berk et al. AJ 2001)

# Selection biases

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

## UVX:

- For  $z > 2.2$  method fails
- Favors selection of optically (not IR) bright QSOs

## The “Universal law” of surveys

Each survey picks objects **respecting its selection criteria**  $\Rightarrow$   
objects **violating** these criteria are **NOT selected!!**  $\Rightarrow$

Bias-free surveys don't exist

# Red & Reddened Quasars

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

## Red & Reddened Quasars

- **Dust makes QSOs look redder (“reddened” quasars)**
  - Projection effects  $\Rightarrow$  obscuration (Type-II AGN)
  - Ly- $\alpha$  clouds  $\Rightarrow$  extinction
- **Intrinsically Red QSOs**

# Do we miss a red / reddened QSO population?

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

## Missed QSO population?

Missed QSOs because of selection effects??

Webster et al. (Nat. 1995), Francis et al (PASA 2000)

Benn et al. (MNRAS 1998), Richards et al. (AJ 2003)

## Warren, Hewett & Foltz (MNRAS 2000): KX

- Based on IR colors, possible to detect “red” quasars
- K-band plays key role in selection
- KX: infrared analog of UVX

# Implementing the KX-method (1)

XMM-LSS survey (Pierre et al. 2004, JCAP)

## Step 1: Perform quasar selection

Select quasar candidates using IR colors & morphology

- $R, z'$ -band data, 4 m tel., CTIO
- $K_s$ -band data, 2.5 m tel., Las Campanas

## Step 2: Multi- $\lambda$ properties of QSO candidates

- 2dF spectra
- $u^*g'r'i'z'$  data, CFHT
- XMM-Newton: X-ray data ([0.5–2] & [2–10] keV)
- Spitzer – SWIRE survey (IRAC 3.6, 4.5, 5.8, 8 and MPIS 24  $\mu m$ )

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

# Properties of the $Rz'K_s$ matched catalog

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

## Catalog matching:

- Independent reduction of  $Rz'$  &  $K_s$  data
- Catalogs extracted (SExtractor, Bertin 1996), correlated using  $K_s$  as a reference

Filter	Frames	Astr. pres.	Phot. pres.	Mag limit
$R$	190	0.3''	0.02	23.5
$z'$	190	0.3''	0.02	22.0
$K_s$	200	0.5''	0.07	18.0


# Selection of the quasar candidates

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work



Stellar spectra: Pickles 1998

# IR colors of the quasar candidates

Scientific  
motivation

The data sets  
– Selection  
process


Properties of  
the quasar  
candidates

Current /  
Future work

## QSO candidates:

point-like morphology  
galaxy-like colors

**85 sources**




# Reconstructing the observational SEDs

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work



# Evaluating the KX

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates


Current /  
Future work

**QSOs: 21**

**Contamin. Rate: 75%**

**Missed QSOs: 3**

**Success Rate: 85%**



Stellar photometry:

Dahn et al. AJ 2002

# Current/ Future work

Scientific  
motivation

The data sets  
– Selection  
process

Properties of  
the quasar  
candidates

Current /  
Future work

## Current

- **Evaluate method:** Compare KX, IR & optically-selected quasar candidate samples
- **Photometric redshifts:** Test existing QSO templates

## Future

Spectroscopic verification (CTIO proposal, March 2006)

- **Characterize contaminants**
- **Measure  $z$**   $\Rightarrow$  reconstruct *rest-frame* SED  
 $\Rightarrow$  Model effect of dust on QSOs