

The Stuck Brain: Constrained Connectivity and Temporal Rigidity in Tinnitus

Marie Detroz, MD,^{1,2} Steven Laureys, MD, PhD,^{3,4} and Audrey Maudoux, MD, PhD^{5,6}

Tinnitus, the perception of phantom sound in the absence of external stimuli, is more than an auditory phenomenon. It is increasingly recognized as a nervous system disorder implicating networks involved in attention, emotion, and self-referential processing. Neuroimaging research, especially with functional magnetic resonance imaging (fMRI), has increasingly moved from focusing on localized auditory cortex dysfunction to examining aberrant large-scale brain network connectivity (Maudoux and Lefebvre, 2014). In this context, the study by Altunkaya et al. makes a timely and methodologically notable contribution by applying constrained Independent Component Analysis (ICA) with both static and dynamic functional connectivity (dFC) analysis to resting-state fMRI data from individuals with chronic, non-bothersome tinnitus due to acoustic trauma (Altunkaya et al., 2025).

Their central finding is that tinnitus patients, compared to matched controls, spend significantly more time in a weakly connected brain state, a pattern interpreted as reduced network segregation and temporal rigidity. This supports a growing hypothesis: that tinnitus involves not just hyperactivity or maladaptive plasticity in auditory pathways but a failure of the brain to dynamically transition between functional states, becoming “stuck” in inefficient configurations (Xie et al., 2024).

Constrained ICA, as applied here, represents an important refinement over conventional ICA. By incorporating prior spatial information to guide component extraction, the authors improve anatomical interpretability while preserving a data-driven analytical framework (Lin et al., 2010). This hybrid approach allows them to identify five core resting-state networks, including the default mode network, salience network, and executive control network, and analyze static and time-varying connectivity among 23 regions of interest. These methodological choices are consistent with previous findings suggesting that tinnitus involves disrupted interactions among attentional, executive control, and self-referential systems (Husain and Khan, 2023).

A more novel aspect of the study is the use of dFC clustering to identify transient connectivity “states.” Unlike previous tinnitus studies focused on static or seed-based FC, Altunkaya et al. reveal that tinnitus patients occupy a low-connectivity state for longer durations than healthy controls.

Although the number of identified states was limited to two, the selection was guided by standard internal validation metrics (Silhouette and Calinski-Harabasz scores), and the authors rightly emphasize the importance of avoiding overfitting in small-sample studies (Altunkaya et al., 2025).

This increased dwell time in a less integrated state suggests impaired metastability, the brain’s ability to flexibly switch between configurations. Such findings resonate with network models of depression, pain, and insomnia, where altered state transitions and dwell times have also been observed (An et al., 2024; Liu and Wan, 2025; Xin et al., 2025). The implication is clear: tinnitus may not be solely a disorder of increased auditory gain but a trait-like disruption of dynamic brain state regulation.

Although the cohort included only non-bothersome tinnitus patients (based on Tinnitus Handicap Inventory [THI] scores), the presence of rigid connectivity dynamics in this group suggests that such alterations may not simply reflect distress or symptom severity. Rather, they may represent a trait-like neural signature, with potential relevance for prognosis or treatment stratification. As observed in the field, correlations between functional connectivity metrics, hearing loss, and THI scores were weak, likely due to small sample size and modeling limitations. Nonetheless, network-level rigidity may constitute a foundational biomarker, independent of current symptom severity, and future studies should examine whether these patterns predict progression to bothersome tinnitus or treatment response in more diverse populations (Piarulli et al., 2023).

Patients exhibiting increased rigidity and reduced network segregation may benefit from interventions aimed at enhancing neural flexibility, such as neuromodulation, neurofeedback, or mindfulness-based cognitive therapy. To better understand the mechanistic underpinnings of this “stuck brain” state, future work should incorporate advanced computational models capable of capturing directionality and information flow. While the current study relies on clustering and state transitions based on correlation matrices, approaches such as dynamic causal modeling, Granger causality, or time-resolved transfer entropy may help identify which brain regions or networks drive these persistent low-connectivity states. These models could enhance mechanistic understanding

¹Sensation and Perception Research Group, GIGA Consciousness, University of Liège, Liège, Belgium.

²Department of Otorhinolaryngology Head and Neck Surgery, University Hospital of Liège, Liège, Belgium.

³CERVO Brain Research Institute, University of Laval, Québec, Canada.

⁴Coma Science Group, GIGA Consciousness Research Unit, University Hospital of Liège, Liège, Belgium.

⁵Department of Otolaryngology, Robert Debré Hospital, Faculty of Medicine, Paris University, Paris, France.

⁶Institut Pasteur, INSERM A006, Institut de l’Audition, Unit Progressive Sensory Disorders, Pathophysiology and Therapy, Université Paris Cité, Paris, France.

and support the development of stratification tools for prognosis or treatment guidance in tinnitus.

Notably, the study by Panda et al. (Panda et al., 2023) in disorders of consciousness employed whole-brain causality models to reveal impaired posterior integration and thalamo-frontotemporal broadcasting. Applying similar techniques to tinnitus could clarify whether thalamocortical loops, insular hubs, or frontoparietal control systems constrain the brain's ability to flexibly transition between states. For instance, persistent thalamic gating failure could plausibly underlie both sensory intrusion (i.e., the phantom sound) and impaired state switching.

The study's generalizability is limited by its small, all-male sample (19 tinnitus patients and 19 controls). While demographic homogeneity reduces variance, it restricts relevance to broader populations, including women and those with bothersome tinnitus. Future research should replicate these results in larger, more diverse, and ideally longitudinal samples to assess predictive value.

Methodologically, the study highlights a key issue in dFC research: lack of standardization. There is no consensus on the optimal number of connectivity states or their interpretation. Harmonized protocols and cross-validation frameworks are needed, as recently proposed in dynamic brain network research (Allen et al., 2014).

In conclusion, the work by Altunkaya et al. offers valuable insights into the dynamic neural basis of tinnitus. By combining constrained ICA with dFC analysis, the study supports a shift from viewing tinnitus as a localized auditory issue to a disorder of impaired brain dynamics. The identification of a prolonged weakly connected state suggests a "stuck brain" profile, potentially serving as a biomarker for prognosis or treatment selection. This framework may guide personalized interventions, including neuromodulation strategies aimed at restoring network flexibility.

Author Disclosure Statement

No competing financial interests exist.

Funding Information

No funding was received for this article.

References

Allen EA, Damaraju E, Plis SM, et al. Tracking whole-brain connectivity dynamics in the resting state. *Cereb Cortex* 2014;24(3):663–676; doi: 10.1093/cercor/bhs352

Altunkaya S, Özmen G, Babür E. Investigating Neural Dynamics in Tinnitus Using Constrained Independent Component Analysis. *Brain Connect*; 2025.

An Z, Tang K, Xie Y, et al.; DIRECT Consortium. Aberrant resting-state co-activation network dynamics in major depressive disorder. *Transl Psychiatry* 2024;14(1):1; doi: 10.1038/s41398-023-02722-w

Husain FT, Khan RA. Review and perspective on brain bases of tinnitus. *J Assoc Res Otolaryngol* 2023;24(6):549–562; doi: 10.1007/s10162-023-00914-1

Lin Q, Liu J, Zheng Y, et al. Semiblind spatial ICA of fMRI using spatial constraints. *Hum Brain Mapp* 2010;31(7):1076–1088; doi: 10.1002/hbm.20919

Liu H, Wan X. Alterations in static and dynamic functional network connectivity in chronic low back pain: A resting-state network functional connectivity and machine learning study. *NeuroReport* 2025;36(7):364–377; doi: 10.1097/WNR.0000000000002158

Maudoux A, Lefebvre P. Tinnitus: Mechanisms induced from human functional neuroimaging studies. *Semin Hear* 2014; 35(02):131–144; doi: 10.1055/s-0034-1372529

Panda R, López-González A, Gilson M, et al.; Coma Science Group Collaborators. Whole-Brain analyses indicate the impairment of posterior integration and thalamo-frontotemporal broadcasting in disorders of consciousness. *Hum Brain Mapp* 2023; 44(11):4352–4371; doi: 10.1002/hbm.26386

Piarulli A, Vanneste S, Nemirovsky IE, et al. Tinnitus and distress: An electroencephalography classification study. *Brain Commun* 2023;5(1):fcad018; doi: 10.1093/braincomms/fcad018

Xie J, Zhang W, Yu C, et al. Abnormal static and dynamic brain network connectivity associated with chronic tinnitus. *Neuroscience* 2024;554:26–33; doi: 10.1016/j.neuroscience.2024.06.034

Xin W, Xue T, Cheng Y, et al. Reconfigurations of dynamic functional network connectivity after 1HZ repetitive transcranial magnetic stimulation in insomnia disorder. *Sleep Med* 2025;126:239–247; doi: 10.1016/j.sleep.2024.12.025

Address correspondence to:

Marie Detroz
Sensation and Perception Research Group
GIGA Consciousness
University of Liège
Avenue de l'Hôpital 1
Liege 4000
Belgium

E-mail: marie.detroz@chuliege.be