

## Intraoperative cytokines production during orthotopic liver transplantation

J. Pirenne, F. Noizat-Pirenne, D. De Groote, Y. Vrindts, M. Lopez, R. Gathy, P. Damas, M. Meurisse, N. Jacquet, P. Honoré, and P. Franchimont

Departments of Surgery, Transplantation, and Anesthesiology, and Laboratory of Radioimmunology, University of Liège, CHU Sart Tilman, 4000 Liège, Belgium. Medgenix Diagnostics, 6220 Fleurus, Belgium

**Abstract.** In summary, we established that a significant production of the monokines interleukin-6, tumor necrosis factor alpha, and interleukin-1 occurred during orthotopic liver transplantation whereas the lymphokines interferon gamma and interleukin-2 were not detected. Levels of interleukin-6 reached their maximum values before and especially at the end of the anhepatic phase. They remained high after the anhepatic phase, i.e. after reperfusion of the new livers. Tumor necrosis factor alpha and interleukin-1 reached their maximum values after the anhepatic phase. Not only were interleukin-6, tumor necrosis factor alpha, and interleukin-1 present in the serum but they could also be detected in the bile produced by these new livers. Mechanisms of monokine production during orthotopic liver transplantation is multifactorial in origin and further studies will have to evaluate the relative contribution of the various factors involved. The possibility of an association between peroperative monokines and transplant outcome and their potential clinical implication will have to be elucidated.

**Key words:** Cytokines – Interleukin-1 beta – Interleukin-6 – Tumor necrosis factor alpha – Liver transplantation

The liver plays a pivotal role in the cytokine network: cytokines act on the liver, liver cells produce cytokines, and the liver is a major clearance organ for circulating cytokines [1]. Orthotopic liver transplantation (OLTx) is a major abdominal surgical procedure in which a diseased liver is removed and a new liver is transplanted. Operative induced tissue injury, endotoxinemia, and kupffer cell activation after liver graft reperfusion are various conditions encountered during OLTx [2, 9, 11, 14]. Since each of these conditions has been shown to stimulate cytokine synthesis, OLTx should thus be accompanied by a significant production of cytokines. Moreover, the absence of

hepatic clearance during hepatectomy and during the anhepatic phase is likely to interfere with cytokine elimination [1]. OLTx should thus impose a considerable impact on both cytokine production and metabolism.

The aims of this study were twofold. The first was to determine peroperative cytokine production during OLTx by measuring cytokines in the serum during the different phases of OLTx. The second was to assess whether cytokines could be detected in the bile produced by the new liver.

### Materials and methods

**Patients.** This study included 7 recipients of hepatic allografts. Indications for OLTx were as follows: hepatocellular carcinoma,  $n = 2$ , primary graft nonfunction,  $n = 2$ , alcoholic cirrhosis,  $n = 1$ , secondary biliary cirrhosis,  $n = 1$ , and posthepatitic cirrhosis,  $n = 1$ .

**OLTx.** The principles of OLTx were removal of the diseased liver and placement of the graft followed by circulation reestablishment and biliary tract reconstruction. During hepatectomy and placement of the graft, a venous bypass was routinely used.

**Collection of serum samples and bile.** Serum samples of systemic venous blood were collected preoperatively in the right atrium through a swan ganz catheter immediately before the anhepatic phase (< AP), at the end of anhepatic phase (AP), after reperfusion of the transplanted liver (> AP), and at day 1 posttransplant. Bile was recovered from the transplanted liver > AP, and at day 1 posttransplant.

**Cytokine immunoassays.** The cytokines interleukin-1 beta (IL-1), interleukin-6 (IL-6), and tumor necrosis factor alpha (TNF) were measured in the serum and bile of all patients. Moreover, the lymphokines interleukin-2 (IL-2) and gamma interferon (IF) were measured in four patients. These cytokines were measured by using specific commercially available immunoassays from Medgenix Diagnostics (Medgenix Diagnostics, Fleurus, Belgium). The methods used, type of tracer sensitivity, precision, reproducibility, and accuracy are described in Table 1.

By performing recovery tests, we established in an earlier study that cytokines can be detected in the bile, using similar immunoassays (Cytokines measurement in the bile, manuscript in preparation).

*Offprint requests to:* Jacques Pirenne, M.D., M.S., CHU Sart Tilman, 4000 Liège, Belgium

**Table 1.** Characteristics of cytokine immunoassays

Cytokine	Type of assay	Tracer <sup>a</sup>	Sensitivity	Precision	Reproducibility	Accuracy	Int. Std. origin	Correspondence weight/units
IL-1	IRMA 1 step	Mab- <sup>125</sup> I	4 pg/ml	3.2%	7.2%	98%	MRC:86/552	1 ng/150 IU
IL-6	IRMA 2 steps	Mab- <sup>125</sup> I	5 pg/ml	5.6%	7.5%	89%	MRC:88/514	1 pg/0.25 IU
TNF	IRMA 1 step	Mab- <sup>125</sup> I	5 pg/ml	6%	7%	101%	MRC:87/650	1 pg/24 mIU
IL-2	RIA sequential saturation	Mab- <sup>125</sup> I	0.5 U/ml	8%	11%	82%	BRMP:ISDP 841	100 pg/1 IU
IF	IRMA 1 step	Mab- <sup>125</sup> I	0.2 U/ml	3.8%	9.9%	97%	NIH:Gg 23/901/530	50 pg/1 IU

IRMA, Immunoradiometric assay. 1 step, standards or samples + tracer incubated together. 2 steps, standards or samples incubated in a first step; washing and incubation of the tracer in a second step RIA, classical radioimmunoassay

<sup>a</sup> By a modification of the chloramine T method

**Table 2.** Preoperative values as compared with control values. Control sera were obtained from normal healthy volunteers

	Control Values	Preoperative values
IL-1 (pg/ml)	< 15 (n = 40)	Range: 0-3 Mean: 0.5
IL-6 (pg/ml)	0 (n = 49) 3-8.5 (n = 29) 24 (n = 1) 72 (n = 1)	Range: 0-205 Mean: 78
TNF (pg/ml)	3-20 (n = 72) 0 (n = 8)	Range: 7-30 Mean: 20

## Results

### A. Preoperative values of IL-6, TNF, and IL-1

Preoperative values of IL-6 and TNF were increased as compared with a group of 80 normal healthy volunteers (Table 2). Preoperative values of IL-1 were not increased as compared with 40 normal sera.

### B. Peroperative production of monokines IL-6, TNF, and IL-1

**Serum.** Peroperative production of monokines IL-6, IL-1, and TNF are shown in Fig. 1 a-c. These three monokines were elevated during OLTx as compared to preoperative values. IL-6 reached its maximal value at the end of AP. IL-6, however, was already elevated before AP and continued to display high levels after AP, i.e. after reperfusion of the new liver. TNF reached its maximal values after AP, i.e. after reperfusion of the new liver. IL-1 had a production pattern similar to TNF with maximum values reached after AP, i.e. after reperfusion of the new liver. The levels reached, however, were low. The levels reached by all these three monokines had returned to preoperative values by day 1 posttransplant.

**Bile.** High levels of IL-6, TNF, and low levels of IL-1 were detected in the bile produced by the new livers (Fig. 2 a-c).

Except for IL-1, biliary levels of these cytokines had significantly decreased by day 1 posttransplant.

### C. Perioperative production of IL-2 and IF

IL-2 and IF were not detected in serum in bile during OLTx.

## Discussion

There were high serum levels of the cytokines IL-6, and TNF, and low serum levels of IL-1 during OLTx. IL-6, TNF, and IL-1 were also detected in the bile produced by the newly transplanted liver. We believe that this cytokine production was triggered by intraoperative events occurring during OLTx procedure as these cytokines reached their maximal value during OLTx and subsequently returned to preoperative values by day 1 posttransplant. A similar observation has already been made by Tono et al. who found a transient but significant elevation of IL-6 in the bile soon after OLTx in rats [13]. The lymphokines IL-2 and IF were detected in the serum in the bile during OLTx. This demonstrated that the cytokine production we observed reflected merely a major stimulation of monokine synthesis by monocytes/macrophages in the absence of lymphocyte activation and detectable lymphokine production at this early stage of recipient immune activation.

At present, there is evidence supporting four possible mechanisms of monokine production during OLTx. These involve the stimulation of cytokine production by (1) tissue trauma, (2) endotoxinemia, (3) lack of hepatic clearance, and (4) kuppfer cell activation. Tissue trauma induced by surgery can lead to monokine synthesis which in turn can initiate the acute phase response that is generally observed after surgical operations [11]. The significance of both monokine production and acute phase response have been shown to correlate with the significance of the surgical operation. Since OLTx is a major abdominal surgical procedure, it should trigger considerable monokine production. Endotoxinemia has been shown to




Fig. 1a-c. Peroperative production of a IL-6, b TNF, and c IL-1

occur during OLTx, especially at the end of AP [9, 10, 14]. Monokines are among the various biologically active substances that are released following monocyte exposure to endotoxin. Thus, endotoxinemia would explain, at least partly, the high levels of monokines that we observed during OLTx, in particular IL-6 whose serum levels reached their maximal value at the end of AP.

A third hypothesis needs to be taken into account. The liver is the principal clearance organ of circulating cytokines [1]. Hepatic clearance is obviously absent during AP, and is already severely compromised before AP due to both surgical manipulation of the diseased recipient liver and to the presence of the venovenous bypass that we rou-

tinely use. Lack of hepatic clearance might account for impaired cytokine elimination and the increased serum levels that are observed before the new liver is functioning. Again, this would be, true mainly for IL-6 whose levels reached their maximum value before and especially at the end of AP. Reduction of hepatic clearance in patients with liver failure might also account for the observation that preoperative serum levels of IL-6 and TNF are increased when compared to normal values [1].

Maximum levels of the monokines IL-1 and particularly TNF were detected after AP, i.e. by the time the newly transplanted livers were revascularized. Interestingly, these monokines were present in the bile produced




Fig. 2a-c. Perioperative biliary a IL-6, b TNF, and c IL-1

by the transplanted livers. Based on these two findings, it is attractive to speculate that the new livers might be a major source of monokine synthesis. This fourth hypothesis is substantiated by several published observations. Kupffer cells are biologically capable of producing large amounts of IL-1, IL-6, and TNF when activated [6]. Caldwell-Kenkel et al. have demonstrated that activation of Kupffer cells occurs after increasing periods of cold ischemic storage and reperfusion of rat livers [2]. Rao et al. have shown that reperfusion following cold preservation not only results in significant activation of Kupffer cells but also in the subsequent release of IL-1 and TNF [12]. Finally, Colletti et al. have demonstrated that TNF is released following hepatic ischemia and reperfusion in a rat model [3, 4].

Although the appearance of IL-1 was demonstrated in serum and bile, this cytokine did not reach as high levels as IL-6 and TNF. A discrepancy between the levels of TNF and IL-1 has already been shown in other pathological circumstances such as sepsis [5]. Coproduction of an IL-1 inhibitor, or presence of the soluble form of the IL-1 receptor might neutralize IL-1 and explain our inability to detect IL-1 in circumstances in which this cytokine is likely to be produced.

High levels of endotoxin at the end of AP have been shown, more than any other factor, to have more serious prognostic implications in terms of graft failure, difficulty of convalescence, and ultimate outcome [9, 10, 14]. Given the fact that endotoxin is a major stimulus for monokine release, it is possible that the effects of endotoxinemia could be mediated, at least partly, by the liberation of cytokines. The monokines IL-6, TNF, and IL-1 exert important effects on hepatocyte metabolism [1]. Increased levels of cytokines during OLTx might thus have important implications with regards to function of the new liver and particularly to the pathophysiology of liver graft dys- and non-function. Because of the positive effect of TNF on cell membrane MHC expression, and on intercellular adhesion molecule expression, these observations might also be significant in terms of graft immunogenicity and pre-

disposition to the rejection process. Interestingly, it has been shown by Howard et al. that preservation injury results in a significantly higher incidence of rejection [8]. The hypothesis that this vulnerability to rejection might be mediated through the effect of cytokines is substantiated by a recent observation that intraoperative elevation of TNF seems to precede early postoperative rejection [7]. OLTx still has a significant perioperative morbidity and mortality [14]. Problems such as coagulopathy, cardiovascular collapse, renal failure, and respiratory insufficiency are still observed. These problems have been attributed, at least partly, to endotoxin [14]. It is possible, however, that high levels of the cytokines IL-6, TNF, and IL-1 would explain better some of these perioperative and early postoperative pathophysiological changes that are currently observed in the course of OLTx and that can jeopardize or even preclude transplant success. For example, TNF has been shown to be directly involved in the development of a severe pulmonary injury following hepatic ischemia and reperfusion [3, 4]. Overall then, perioperative assessment of monokines might be a useful prognostic premonitor in predicting short and long term outcome after OLTx. In order to assess a possible correlation between perioperative monokine production and both graft and patient outcome, we are now conducting a prospective long-term analysis including a larger number of patients.

In summary, we established that a significant production of the monokines IL-6, TNF, and IL-1 occurred during OLTx whereas the lymphokines IFN and IL-2 were not detected. Levels of IL-6 reached their maximum values before and especially at the end of AP. They remained high after AP. TNF and IL-1 reached their maximal values after AP, i.e. after reperfusion of the new livers. The mechanisms of monokine production during OLTx are multifactorial in nature and further studies are needed to evaluate the relative contribution of the various factors involved. The possibility of an association between perioperative monokine levels and transplant outcome and their potential clinical implication will have to be elucidated.

## References

1. Andus T, Bauer J, Gerok W (1991) Effects of cytokines on the liver. *Hepatology* 13:364-375
2. Caldwell-Kenkel JC, Currin RT, Tanaka Y, Thurman RG, Lemasters JJ (1990) Kupffer cell activation and endothelial cell damage after storage of rat livers: effects of reperfusion. *Hepatology* 13:83-95
3. Colletti LM, Remick DG, Burtch GD, Kunkel SL, Strieter RM, Campbell DA (1990) Role of tumor necrosis factor-alpha in the pathophysiologic alterations after hepatic ischemia/reperfusion injury in the rat. *J Clin Invest* 85:1936-1943
4. Colletti LM, Burtch GD, Remick DG, Kunkel SL, Strieter RM, Guice KS, Oldham KT, Campbell DA (1990) The production of tumor necrosis factor alpha and the development of a pulmonary capillary injury following hepatic ischemia/reperfusion. *Transplantation* 49:268-272
5. Damas P, Reuter A, Gysen P, Demonty J, Lamy M, Franchimont P (1989) Tumor necrosis factor and interleukin-1 serum levels during severe sepsis in humans. *Critical care Medicine* 17:975-978
6. Decker K (1990) Biologically active products of stimulated liver macrophages (Kupffer Cells). *Eur J Biochem* 192:245-261
7. Fugger R, Hamilton G, Steininger R, Mirza D, Schulz F, Mühlbacher F (1991) Intraoperative estimation of endotoxin, TNF and IL-6 in orthotopic liver transplantation and their relation to rejection and postoperative infection. *Transplantation* 52:302-306
8. Howard TK, Klintmalm GBG, Cofer JB, Husberg BS, Goldstein RM, Gonwa TA (1990) The influence of preservation injury on rejection in the hepatic transplant recipient. *Transplantation* 49:103-107
9. Miyata T, Todo S, Imventarza O, Ueda Y, Furukawa H, Starzl TE (1989) Endogenous endotoxemia during orthotopic liver transplantation in dogs. *Transplant Proc* 21: 3861-3862
10. Miyata T, Todo S, Selby R, Yokoyama I, Tzakis A, Starzl TE (1989) Endotoxaemia, pulmonary complications, and thrombocytopenia in liver transplantation. *Lancet* I:189-191
11. Nishimoto N, Yoshizaki K, Tagoh H, Monden M, Kishimoto S, Hirano T, Kishimoto T (1989) Elevation of serum interleukin 6 prior to acute phase proteins on the inflammation by surgical operation. *Clin Immunol Immunopathol* 50:399-401
12. Rao PN, Liu T, Snyder JT, Platt JL, Starzl TE (1991) Reperfusion injury following cold ischemia activates rat liver Kupffer cells. *Transplant Proc* 23:666-669
13. Tono T, Monden M, Yoshizaki K, Valdivia LA, Nakano Y, Gotoh M, Ohzato H, Doki Y, Ogata A, Kishimoto T, Mori T (1991) Interleukin 6 levels in bile following liver transplantation. *Transplant Proc* 23:630-631
14. Yokoyama I, Todo S, Miyata T, Selby R, Tzakis AG, Starzl TE (1989) Endotoxemia and human liver transplantation. *Transplant Proc* 21:3833-3841